[illegible]

[illegible] M.D.

[illegible]

28,499/B

18/6

12/6

Bt from Mr Brown 13/11/92

Heffer & Sons Cambridge 12/6 ... 17/5/34

Ex bibl. Min. Ebor. B

e Collegio Universitatis

[illegible]

William Munk M.D.

Digitized by the Internet Archive
in 2019 with funding from
Wellcome Library

https://archive.org/details/b30503218

79674

PAULI HERMANNI
Med. Doct. & Prof. Lugduno-Batavi
Celeberrimi

CYNOSURA MATERIÆ MEDICÆ,

SEU

BREVIS AC SUCCINCTA METHODUS NOTITIAM SIMPLICIUM MEDICAMENTORUM COMPARANDI,

DESUMTA

AB INTERNA PARTIUM CONSTITUTIONE,

UBI

SIMPLICIA JUXTA PARTES SUAS CONSTITUTIVAS IN CERTAS CLASSES DISTRIBUUNTUR, POST DENOMINATIONEM SUCCINCTA DESCRIPTIO TRADITUR, VIRES ATQUE VIRTUTES ENARRANTUR, ET MODUS DEXTRE USURPANDI MONSTRATUR,

IN LUCEM EMISSA

A

JOHANNE SIGISMUNDO HENNINGERO,
Med. Doct. & Prof. Publ. Ord.

ARGENTORATI,
Typis & Sumptibus JOH. FRID. SPOOR, Viduæ.

M DCC X.

1710

WELLCOME
HISTORICAL
MEDICAL
LIBRARY

PRÆFATIO AD LECTOREM.

QUam vaſtus ſit Materiæ Medicæ ambitus, & quanta cum difficultate notitia ejus acquiratur quivis in hac re vel leviter verſatus perſpicit, & querelæ eorum hujus cognitionem ſibi comparare allaborantium ſatis teſtantur. Tam late enim patet ac ipſa natura patet, & qua tota fere Phyſicæ pars ſpecialis continetur.

Uti vero in vaſto Oceano navigantibus contingit, ut navigationis viam ductu & indicio cynoſuræ dirigant, quo iter feliciter perficiant, ita ſi præſens hoc opus ſub titulo Cynoſuræ Materiæ Medicæ emiſero, non peccabo, quam facile enim viam monſtret, in hoc longo alias & anfractuoſo Materiæ Medicæ itinere, plures jam inſigni, cum emolumento experti fuere.

Optavi propterea ſerio, jam à pluribus annis, ut hoc laudatiſſimum & cedris dignum opus, à quodam aliquando luci publicæ committeretur, cum Studioſa Juventus maximam inde utilitatem ſit perceptura. Sed cum hactenus nil tale videre contigerit, nam opuſculum ſub nomine Lapis Lydius Materiæ Medicæ emiſſum, non niſi deſcriptum obtinere potui, imo potius conſpexerim plures hinc inde in libris emiſſis Beati Autoris opera pro ſua uti, interim tamen apud plurimos magnificis elogiis hoc opus exornari, non inconſultum fore duxi, ut hunc laborem in me ſuſciperem.

Cum enim in Collegiis & Prælectionibus de Materia Medica à me habitis, Auditoribus meis hunc Authorem de meliore nota commendaſſem, atque ex hac cauſa ſæpius deſcriberetur, inſuper in poſterum, DEO vitam concedente juxta Autoris ductum Materiam Medicam tradere conſtitutum atque propoſitum ſit, ut & mihi hac ratione inſtar cynoſuræ inſerviret & deſcribendi onere Auditores meos ſublevarem, atque præterea Bibliopola promte operam ſuam promiſerit, denique hoc inſtitutum ab Experientiſſimis quibuſdam hujus Civitatis Medicis non inutile æſtima,

tum

tum fuerit, eo promptius hoc ipſum in me ſuſcepi.

Confirmarunt ulterius me in hoc propoſito nonnulli ex CL. Dn. Auditoribus, qui egregios ut in hoc ſtudio profectus publice oſtenderent, in Solenni Auditorio per penſa hoc opus pertractandum & defendendum non ſine ſui laude & fructu in ſe ſuſceperunt.

Certo igitur Tibi perſuaſum habeas, me nec lucri captandi, nec gloriæ aucupandæ gratia hoc ſuſcepiſſe, ſed ſolum commodum & utilitatem Studioſæ Juventutis, unicum mihi fuiſſe ſcopum præfixum. Id quod ſi hoc ipſo obtinuero eſt de quo mihi ſummopere gratulabor. Vale.

Nomina Juvenum Inſigniſsimorum

qui in ſolenni conflictu Academico certa penſa non ſine laude defendenda in ſe ſuſceperunt.

1. JOH. JACOBUS Fried/ Argent.
2. JOHANNES GOLL, Argent.
3. JOH. DANIEL SALTZMANN, Argent.
4. ABRAHAM Kübler/ Argent.
5. GEORGIUS AUGUSTUS LUCIUS, Coburg.
6. JOH. JACOBUS Röderer/ Argent.
7. JOH. JACOBUS Marbach Argent.

PROEMIUM.

MAteria Medica dicitur nobis omne id, quod corpus noſtrum alterare & immutare poteſt, ſeu alimentoſum ſeu medicamentoſum, ex qua Medici ſelectiora eligunt.

Quatuor inprimis adminicula hic conſideranda veniunt. Primum ſuppeditabit ipſa ſimplicium Etymologia, ex qua ſæpe latens vis erui poteſt. v. c. in conſolida, tuſſilagine, hepatica, pulmonaria. Alterum petemus ex frequentiſſima ſimplicium inſpectione & uſu. Tertium deſumemus ab exploratione ſaporis & odoris; ſic odor fragrans calorem & vim aperiendi notat, conſiſtit enim hic in particulis ſubtiliſſi-

mis, quæ celeriorem Sanguinis motum, & per conſequens calorem producere aptæ ſunt natæ. Hic odor fragrans ſi ſuavis fuerit, denotat vim cardiacam, ſin vero fragrantiam odoris gravitas ſimul comitetur vis Uterina ſignificabitur, conſtant enim ſimilia particulis quidem ſpirituoſis, minus tamen volatilibus ſed fixioribus. Quoad ſaporem, quæcunque habent acrem, calida ſunt, & aperientia; quæ amarum, calida quidem ſunt, fixiori tamen ſale conſtabunt; quæ dulcem, ſunt temperata ſed viſcedine donata; quæ acidum & auſterum refrigerant & adſtringunt. [Novem ſpecies ſaporum recenſentur; Acerbum, auſterum, acidum, dulce, unctuoſum ſeu pingue, ſalſum, amarum, acre, inſipidum. Qua de re conferri meretur Davidis Abercrombii Nova Medicinæ tum Practicæ tum ſpeculativæ clavis, ſive ars explorandi Medicas plantarum ac corporum quorumcunque facultates ex ſolo ſapore. Londini 1685 edit.] Quartum denique à cujuſque plantæ nota eſſentiali & charactere deſumitur, unde etiam omnes plantæ, quæ eandem habent

notam

notam eſſentialem, ejuſdem ſunt facultatis: ſic omnes herbæ bacciferæ ſunt diureticæ, ſic omnes herbæ umbellatæ ſunt ſale volatili oleoſo præditæ.

Cognoſcuntur itaque virtutes plantarum omnium maxime Ratione, Experientia, & Analogiſmo.

PARS PRIMA.

REGNUM VEGETABILE EXHIBENS.

CAPUT PRIMUM.

DE RADICIBUS.

RADICES SALE VOLATILI OLEOSO CONSTANTES.

Acori vulgaris.
Indici.
Zedoariæ.
Galangæ majoris.
minoris.
Zinziberis.

Conveniunt in cruditate acida, Colica, obſtructionibus menſium, ſcorbuto, hydrope, vertigine, ſingultu, cordis palpitatione.

ACORI RADIX.

ACori veri [SEU CALAMI AROMATICI OFFICINARUM] C.B.P. radix eſt, oblonga, paululum compreſſa, in ſuperficie quaſi geniculata, foris ex flavo rubeſcens, intus alba, fungoſa, ſaporis acris ſuaviter amaricantis & aromatici, odoris fragrantis. [Refertur ad herbas aquaticas graminifolias non culmiferas flore ſtamineo, foliis iridis modo ſitis, hoc eſt acie folii cauli obverſa, aromaticis, odoratis, ſpica ſimplici eleganti juli ſpecie, ſeminibus in julum ſeu nucamentum coactis.]

Ob ⊖ volatile [acre] oleoſum [aromaticum] penetrantiſſimum ſuccurrit acidæ cruditati Ventriculi, in torminibus abdominis inde ortis, obſtructione Menſium è vitio Ventriculi, in ſcorbuto & hydrope pro confortante.

Primo vere effoſſa radix diutius fert ætatem ac aliàs.

Acorus in ſubſtantia rarius exhibetur, conditus vero frequentius, ſi vero datur crudus, datur à gr. v. ad xv. conditus vero à ℈j ad ʒj.

Ubi deſtillatur radix hæc cum ▽, ſi deguſtaveris aquam ingratiſſima eſt, ſi oleum, nauſeoſum, ſemper ingratiſſimum, contra fixum radicis in fundo remanens optime ſapit. Infuſum acori fit in ♎u Juniperino qui optimus, vel in Spiritu vini, ad pint. j. additur acori ℥j. Admiſcetur quoque morſulis.

Stimu-

Stimulatur acorus granis paradiſi. Temperatur partim dulcibus, partim minus acribus. Convenit in torminibus colicis ortis á cruditatibus acidis, item in ructibus moleſtis. e. c. ℞. acor. condit. ℈j Caſtor. gr. ix. l. x. Laud. opiat. gr. ij flaved. Cort. aurant. cond. ʒj add. Syr. Cortic. aurant. q. ſ.

Etiam in Morbis ſoporoſis acorus externe exhibetur. e. c. ℞. rad. pyretr. ſaxifrag. acori. ana ℥j ſummitat. origan. Mj. ſem. cardamom. min. ℥ß coq. in ſ. q. ▽æ, colatur. cap. ℥viij add. ⊖ gemm. ʒij f. Clyſter. Si non ſat acris, poteſt ſtimulari pulpa colocynth. ʒj linteolo involuta.

In Epithematibus contra Paralyſin ſive reſolutionem nervorum: ℞. acor. ℥j rad. pyretr. ℥iß helen. Zedoar. Zinzib. ana ℥ß. ſummit. hyperic. fol. majoran. ſpicæ. ā. p. ij affund. ſpir. Vin. vel ⩊ C. C. ſuccinat. vel ⩊ therebinth. ad emin. 3. vel 4 digitor. fiat infuſum Paralyticum externum. Hoc enim ſpirituum influxu ceſſante vellicantur nervi, ut rurſus ſpiritus influant.

ACORUS INDICUS.

[Acorus Aſiaticus radice tenuiore. Herm. Catal. de quo ibi ita: acorus hic omnibus partibus convenit cum Europæo, excepta ſola radice, quæ tenuior eſt, & compactior, ſuperficie crebrius geniculata, odore & ſapore gratiori. Sponte ſua luxuriat in Malabara & inſula Zeylan, Incolis Vazabu & Vazumbo audit.

Cum priore viribus convenit. vid. Garz. ab Horto aromat. hiſt. lib. 1. cap. 32.]

[CALAMUS AROMATICUS.

De quo ex prælectionibus MStis Hermanni Samuel Dale Pharmacolog. p.m. 327. ita loquitur: Calamus Aromaticus veterum eſt plantæ Ægyptiacæ (Alpin. Exotic. Caſſabel Darrirà dictæ; quæ tetrapetala eſt, ſiliculis oblongis, acuminatis, in baſi cauliculatis, ſeminibus nigris minutiſſimis refertis) cauliculi tenues, geniculati, foris pallide flaveſcentes, intus candidi, includentes medullam fungoſam, levem, ſimilem telis aranearum in ſe collapſis, ſaporis acris, ſuaviter amaricantis, odoris aromatici & fragrantiſſimi. Itaque non eſt radix ſed arundo aut cauliculi odorati.

Deſcribit hanc plantam Proſper. Alpin. de plant. Exotic. l.2.c.7.p.m.195. & de uſu Medico agit idem Alpin. de Medicina Ægyptiorum libro quinto. Aliàs de calamo odorato Dioſc. l.1. c.17. ingreditur compoſitionem theriacæ, ſed hodie noſtris officinis eſt incognitus, & loco ejus recipitur acorus verus.]

ZEDOARIA.

Eſt Zedoariæ Chinenſis radix, tuberoſa, brevis, compreſſa, foris cinerea, intus candida, quando cortex exterior eſt detractus, paucis fibris intertexta, ſaporis amaricantis aromatici, odoris fragrantis ad camphoram aliquantum accedens. Folia ad curcumæ folia accedunt, ſuntque palmæ magnitudine, fert florem in peculiari ſpica ut galanga, quos flores ſequuntur loculamenta terna.

Conſtat ⊖e volatili oleoſo acriore penetrantiore & ſpirituoſiore acoro, propterea ſe altius extendit, ut ſimul ſudorem moveat, phlegma contumax Pulmo-

num

num potenter diſcutiat, & colicæ convulſivæ hyſtericæ medeatur. Nauſeam tollit vel odor ſolus, vel ſi ſub Lingua teneatur, vel ſi externe Ventriculo applicetur vel ſola vel cum fermento. In vindicando ſanguine â malignitate tanquam panacæa habetur, aliis loco ſtimuli additur, eſt alexipharmacum egregium. In aſthmate etiam convenit, in nauſea poſt crapulam.

Sal volatile â curioſis deſtillatur, aqua ingrata quidem, ſed maxime ſudorifera, unde pro ▽ alexipharmaca uſurpo. [Conf. Hermann. Hort. Lug. Bat. ſub titul. Zinziber latifolium ſylveſtre. Zerumbet Garz. ab Hort. l.1. c.47. ubi hanc plantam prolixe deſcribit, & inter alia hæc de deſtillatione habet: Sale volatili oleoſo aromatico abundare hujus aromatis radicem, probat inter alia ejus deſtillatio. Tranſmittit enim per alembicum ▽ fragrantem, cum ſufficiente oleo, cui ſi recens fuerit deſtillata innatat ſal volatilis paucus forma nivis aut Camphoræ.]

Doſ. gr. vij ad ℈ß in ſubſtantia, condita vero ℈j. in infuſo ad ʒij, edulcoratur vel ſaccharo vel melle.

Zedoariæ optimus ſtimulus eſt camphora, quod quaſi à priori illis notum, qui recentem effoſſam examinarunt, qui dicunt, quod omnino ſpiret camphoram.

Specificatur in colica cum Caſtoreo, ut & in affectibus Uterinis.

℞. Zedoariæ ʒj Camph. ℈j ſacch. cand. ʒj vel ij M. f. pulv. pro novem doſibus, ſingulis diebus una, contra omnes malignitates.

℞. Zedoar. ℈ß. ſacchar. ʒß. M. f. pulv. ſumendus ex aceto contra nauſeam.

℞. Ze-

℞. Zedoar. ʒß. ⚗ ſerpill. gt. j. Laud. opiat. gr. j. Conſerv. roſar. ⊖lat. ʒj Syr. acetoſ. Citri q. ſ. M. f. bolus in quavis nauſea & Vomitu.

GALANGA MAJOR ET MINOR.

Galanga duplex eſt, major & minor. Major eſt planta Indica Banchabe dicta, radix tuberoſa, pollicem [& amplius, imo duos interdum] craſſa, nodoſa, inæqualis, in ſuperficie circularibus geniculis donata, [cortex] flaveſcens, [aut fuſcus] intus palleſcens, ſaporis acris aromatici, [aliquantum ingrati] odoris, [quodammodo] fragrantis.

Minor eſt plantæ indicæ Lagondi dictæ radix tuberoſa, nodoſa, majori ſimilis, ſed tenuior, digiti craſſitiem adæquans, odoris fragrantioris & jucundioris quam in galanga majore.

Galanga minor quæ efficacior eſt majori, relinquit poſt acredinem, intenſiorem ſtypticitatem, unde temperatior eſt, & in vertigine à vitio Ventriculi, in torminibus abdominis & doloribus poſt partum ſpecifica eſt, cæterum nervis & ſpiritibus commendatiſſima eſt, in ſingultu quoque & cordis palpitatione. [Ad claſſem iridum â nonnullis utraque planta refertur, quod ex foliorum ſimilitudine & zonis in radice circulariter cinctis probant.]

Minor nonnihil viſcidi habet in receſſu ſeu adſtringentiæ nonnihil, cerebro gratior eſt quam acorus, ad ſeptimum vel duodecimum annum poteſt conſervari. Fruſtula ſeu gemmæ radicis ſolent decerpi à Chinenſibus, iterumque plantari, etiam ſeminibus multiplicantur. In india orientali galangam conditam habent. Cum bucella panis & butyri capiunt pulve-

pulverem galangæ mulieres indicæ in doloribus poſt partum.

Datur à ℈ß ad ℈j in ſubſtantia, in infuſo velut acorus. juvatur zedoaria, helenio.

Specificatur in vertigine Doronico, cumino. Stimulatur Caſtoreo, ſale ſuccini, laudano opiato in doloribus poſt partum.

Pro juvene ſcotomia jejuno Ventriculo ſurripiente & exinde in vertiginem incidente.

℞. Galang. Zedoar. ana ℈j caſtorei gr. vij ☍ chamom. deſt. gt. ij f. pulv. exhibendus ad ℈ß ſingulis mane.

℞. Galang. q. v. M. cum ▽ plantag. fac epithema, impone regioni cordis, ad cordis palpitationem tollendam.

℞. Galang. ʒj ſem. aneth. ʒß f. infuſum ex vino ad ſingultum, cochleatim exhibendum.

℞. Galang. ℈j. croc. orient. gr. v. exhibe in vino vel ▽ cinamom in doloribus poſt partum.

℞. Rad. doronic. ſem. cumin. galang. min. ana gr. xv. add. pro ſtimulo gt. j. ☍ d. cinam. dari poteſt in vertigine. vel

℞. Galang. ʒß. doron. ʒj zinzib. chin. cond. ʒij ſem. cumin. ℈j ☍ d. cumin gt. iiij conſerv. flor. anthos ℥ß. ſyr. flor. tunic. q. ſ. f. Elect.

ZINZIBER.

Eſt radix tuberoſa, nodoſa, ramoſa, brevis, & paululum compreſſa, pallida, fibris crebrioribus dotata, ſaporis aromatici acerrimi fervidi, inſtar piperis, odoris [aliquantum] fragrantis.

Variæ ſunt ſpecies Zinziberis, quæ tantum differunt ratione fibrarum, quo vero plures fibræ, eo ignobilius, quo pauciores eo præſtantius, ſed hæc ultima ſpecies rarius in officinis proſtat, radix enim multas habet fibras & eſt lignoſior. [Planta ſimilis eſt iridi, unde à nonnullis appellatur, Iris latifolia tuberoſa, Zinziber dicta, flore albo.]

Cum Zedoaria convenit, excepto camphorato odore, hinc tanta vis alexipharmaca ipſi non ineſt. Gratius quidem eſt, ſed acrius & calidius & magis deſiccativum. Corrigit purgantium vehementiam præcipue agarici. Melancholicis & Cholericis largior uſus Zinziberis non conducit, quia humores nimis exſiccat, & ad motum ineptiores reddit.

Corrigitur acidis blandis, quia habet Naturam alcalinam, nempe rob ribium. berber. ſucco limon.

In morborum declinatione mire appetitum reſtaurat.

Recens effoſſa vim habet laxandi alvum, præcipue ejus ſuccus expreſſus, ſed ſicut in ſemine coriandri recens malignitas tractu temporis evaneſcit.

Zinziber in ſubſtantia datur à gr. v. ad xv. conditum vero ad ℈j. vel ʒß [& ultra] de cætero uſurpatur ut Zedoaria.

Feliciter quoque datur decoctum Zinziberis conditi ex aqua, in tuſſi ex materia viſcoſa in Pulmonibus hærente.

℞. Zinz. cond. ℥ß. conſerv. abſynth. ℥iij ∽ ⊕li vel 🜍is per campan. q. ſ. ad gratum acorem M. f. conditum, ſumendum in languido appetitu in declinatione Morbi.

℞. Agaric. ℈j. troch. alhand. gr. v. Zinz. gr. x. f. pulv. purgans pro j doſi. [Agaricus non facile in forma pulveris ob levitatem exhibetur, melius in infuſo vel vino medicato.]

℞. Gran. Till. gr. iij Zinzib. gr. vij. M. f. pulv. pro una doſi. Robuſtis exhibetur ſpecifice contra hydropem. Indi pilulas parant cum mucagine tragacanthi.

***

RADICES SALE VOLATILI DULCEDINE JUNCTA CONSTANTES.

Imperatoriæ.
Angelicæ.
Mei.
Leviſtici.
Nizin vel Niſi.

Locum habent in ophthalmia, colica, paſſione hyſterica, cardialgia, lypothymia, confortanda memoria, lochiiſque promovendis.

IMPERATORIA SEU OSTRUTIUM.

Eſt imperatoriæ majoris C. B. P. radix oblonga, pollicem craſſa, ſcirrhoſa [rugoſa] & geniculalata, foris fuſca, intus alba [fibris lateralibus exornata] ſpirituoſior penetrantior & amarior ſuperioribus, unde & ſalibus fixis conſtat. [Refertur ad plantas umbel-

bellatas ſeminibus nudis binis ſequentibus ſingulos flores.]

Radix de cætero coincidit cum angelica, ſtimulus eſt cæteris radicibus aromaticis. [Sapor acris aromaticus calefaciens & vellicans linguam, odor fragrans.]

In Colica hyſterica, convulſiva, tympanitide, tanta ſæpe agit cum violentia, ut freno opus habeat, nempe leviſtico, meo, alterave dulci & viſcoſa radice, in febribus intermittentibus præcipue tertiana & quartana ſpecificum eſt, ut & in morbis ſoporoſis.

Datur à gr. xv. ad ℈j. in ſubſtantia. à ʒj ad ʒij in infuſo.

Modus conjungendi, ſtimulandi, ſpecificandi videatur ex formulis.

℞. Imperat. ℈ß. angelic. ℈j. Mei gr. vij. M. f. pulv. in colica paſſione, addi poteſt 🝆 d. mac. gt. j. vel ij. Laud. op. gr. ij. poſſunt & addi Caſtorei gr. v.

℞. Imperat. angelic. ana ʒiij Zinzib. Zedoar. ana ʒj bacc. laur. exc. ʒij l. iij. flor. chamom. ʒj f. infuſ.

℞. Imperat. gentian. Chin. Chin. ana ℈ß ⊖ prunell. abſynth. Cent. min. ana gr. j. M. f. pulv. pro una doſi contra quartanam & omnes febres intermittentes.

ANGELICA.

Eſt Angelicæ ſativæ C. B. P. radix oblonga, aliquantum reſinoſa, ex capite craſſiori in fibras ſæpe pollicares, vel tenuiores diviſa, foris ex flavo rubeſcens, medulla interior alba, ſaporis acris ſuaviter amaricantis, cum quadam blanda dulcedine, odoris fragrantis. Conſtat ⊖e volatili 🝆ſo, adjuncta dulcedine quadam, quibus incluſæ particulæ anodynæ.

[Jun

[Jungitur plantis Gymnodiſpermis umbellatis.] Symbolizat cum Zedoaria & imperatoria, aſt temperatior eſt, & ob ſui blanditiem ad dulce vergentem, opus ſuum cum quadam vi anodyna abſolvit; eandem ob cauſſam ſingulare quid efficit in vitiis Pulmonum, [unde & Germ. Bruſtwurtz dicitur] & omnibus Uteri affectibus, ut & in Colica hyſterica, cordis & ſpirituum vitiis.

Zedoariæ, Zinziberi, imperatoriæ ſi jungitur, gratiam peculiarem conciliat eaſque temperat.

Jungitur ſuperioribus radicibus frequenter, ſtimulatur Camphora.

Datur ut ſuperiores radices in ſubſtantia, frequentius tamen in infuſo, non raro etiam in extracto feliciter datur.

Poteſt dari in Colica á vitiô Ventriculi ab acido acri humore orta ſeq. m.

℞. Extr. angelic. ʒß. ▽ flor. chamom. ℥ij [vel ▽ Carm. Schrœd.] Syr. ê Cortic. aurant. ʒij l. ℥ß. ſi Dolores urgeant, non ſufficit, ſed ſtimulo opus habet Laud. op. gr. j. l. ij. Si Colica hyſterica ſimul add. Tinctur. Caſtor. ʒj. Idem etiam in Colica convulſiva convenit, additur Clyſteribus contra Colicam vel ſimplicem vel compoſitam.

Infuſum Angelicæ.

℞. Angelic. Zedoar. galang. ana ʒij Caſtor. ʒj Camph. gr. xv. add. liq. Corn. Cerv. ſuccinat. ad eminent. 3. vel 4. digitorum. f. Eſſent. vel Tinctur. doſ. gt. XX. XXX. ad 50. ſi magis anodyn. expetas add. Laud. Op. ℈j.

Oleum Angelicæ egregie inſervit loco balſami Uterini, aqua deſtillata non adeo eſt ingrata, nec oleum.

MEUM.

Eſt Meum foliis anethi. C. B. P. radix oblonga, digiti craſſitie, foris ferruginoſa, intus pallida, & aliquantum reſinoſa, ſaporis & odoris ad angelicam & leviſticum accedentis. Ob vim anodynam ſuperior eſt, quam angelica, ſed ob acredinem remiſſior quam angelica, hinc ab angelica non niſi remiſſiori gradu differt. Ob ſui nimiam fragrantiam nonnullis moleſta eſt & odioſa, pectoris anguſtias & gravedinem capitis augens, hinc in doſi conſideratiore exhibenda. In pectoris Pituita acriori temperanda & evacuanda admodum conducibilis eſt. [Itidem annumeratur plantis umbellatis ſeminibus nudis binis ſequentibus ſingulos flores.]

Datur ut angelica, aſt in doſi parciori, juncta cum imperatoria.

Detenta ſub Lingua, arcet in multis paroxyſmum hyſtericum, nempe.

℞. Mei ʒj. imperat. ʒß. f. l. a. paſta cum ▽ appropriata. Clyſteribus in Urinæ ſuppreſſione cum congeneribus feliciter jungitur. v. c.

℞. Mei. Leviſtic. Angelic. ana ℥ß. Chærefol. Mj coque in ▽ Colatur. ℥viij add. 🝆 chamomill. vel amygd. dulc. ℥j ⊖ ʒj f. Clyſter.

LEVISTICUM.

Eſt Liguſtici vulgaris C. B. P. radix oblonga, pollicari craſſitie, in fibras craſſiores diviſa, foris fuſca,

intus

intus pallida, ſaporis acris aromatici ſubdulcis, odoris jucundi & fragrantis, nonnullis gravis & moleſti. [Semina oblonga, compreſſa, ſtriata, ſaporis acris aromatici ſubdulcis, odoris fragrantis. Eſt ſimiliter planta umbellata ſeminibus nudis binis flores ſequentibus.]

Differt â Meo acredine remiſſiore, dulcedine vero viſcoſiore & pertinaciore. Cæterum ſpecifice in ictero laudatur, dum vel ſua dulcedine acriorem Bilem mitigat, aut Bilem in inteſtinis cunctantem incidit & promovet, ob eandem rationem diureticum audit & antihyſtericum.

Cunctantibus à partu lochiis medetur, & olim â muliere quadam Anglicana pro ſecreto habitum.

Exhibetur velut Meum & angelica ℞. Leviſtic. ʒß exhibe ex ſucc. Leviſtici recenter expreſſi cochl. j. ad retenta lochia, tali modo exhibuit fœmina illa Anglicana. Raius in hiſt. plant.

℞. Leviſtic. ℥ß Mei. Zedoar. Zinzib. ana ʒij Cinam. ʒiß vel ij herb. abſynth. Meliſſ. ana Mj. Cort. aurant. Citr. ana ℥ß affund. vin. ſ. q. fiat infuſum antictericum, præprimis quando à Bile viſcidiore ortum habet.

NINZIN.

Eſt plantæ Japonenſis & Chinenſis dictæ Gin-Sen, vel Genſing, radix oblonga, digiti minimi craſſitie, ut plurimum bifida, aliquando multifida, circa finem filamentoſa, intus & extus flaveſcens, quæ ſpecies inter cæteras in M.N.C. contentas optima, ſaporis ſubdulcis, ſubacris, ſubamaricantis & aromatici gratiſſimi, odoris ſuavis.

De

[De hac radice Genſingh. vid. Andr. Cleyerus C. N. C. Dec. II. Ann. 4. obſ. 2. p. 3. ad 6. it. Chriſtian. Mentzelius C. N. C. Dec. II. Ann. 5. obſ. 39. p. 73. ad 79. ubi etiam figuras ex herbariis Sinicis & aliis authoribus producit. D. Mich. Bernh. Valentini in Muſ. Muſeor. l. 2. ſ. 2. c. 9. p. 163. allegat Noſtri D. Hermanni Colleg. Mat. Med. ubi putat eſſe ſpeciem Siſari Montani, quia ex ſemine ex Japonia accepto ſimilis planta excrevit, licet viribus radicem Ninſeng non æquaverit, id quod tamen diverſitati telluris adſcribit. Interim hoc certum, figuram in M. N. C. à Menzelio exhibitam â Siſaro montano non adeo abludere.]

Plukenetius Phythograph. Londini Fol. 1691. edita optimam habet figuram. Bontius vero valde erroneam.

Sua ſponte naſcitur in Japan & China, provenit etiam in regno Siam, naſcitur in declivioribus non in montanis, ut quidam volunt. Autumno [& hyeme] collecta planta longe eſt præſtantior, quam ſi vere colligas. Diu poteſt conſervari, cariſſima etiam eſt in India Orientali ℔j. 50. Imperial. Utuntur veluti Thée in India.

Temperatiſſima eſt & gratiſſima, inter angelicam, meum & leviſticum, & propter hujus temperiem convenientiſſimam, habet analogiam cum ſanguine humano, proinde polychreſta habetur, in ſpecie tamen commendatur in colica, Lypothymia, convulſione, confortanda memoria. In vertigine laudatur, quia motum inordinatum ſpirituum reſtituit, & ratione ſuavitatis compeſcit, unde horis matutinis maſticatur.

Datur

Datur ex infuſo ▽æ ferventis, ut & in pulvere à ℈j. ad ℈ij.

℞. Rad. Ninzin ʒj infund. ▽ fervent. ℥v. infuſum cape ad colicam Nephriticam, Ventriculi debilitatem & roborandam Naturam.

℞. Rad. Ninzin. ʒj angel. ʒß ſantal. Citr. ℈j f. infuſ. ▽ ferventiſſ.

℞. Rad. Ninzin. ℈ij vel ʒß flaved. Cort aurant. Citr. ana gr. vij. Laud. op. gr. j. Caſtor. gr. v. M. f. pulv. add. conſerv. chamom. f. bolus, in colica paſſione egregius.

℞. Rad. Ninzin. ʒß. conſ. alkerm ʒß ambr. gryſ. -ʌ- aurant. ſolut. gr. v. M. f. Elect. polychreſtum in capitis affectibus. Hoc utebatur Generalis Indiæ ſingulis mane.

*
* *

RADICES CONSTANTES SALE VOLATILI OLEOSO AROMATICO, MIXTIS PARTICULIS TERREIS ET SALINIS FIXIS.

Radix Valerianæ.
Serpentariæ Virginianæ.
Coſti albi.
Cyperi longi & rotundi.
Spicæ nardi & Celticæ.
Vincetoxici.
Helenii.
Ireos Florentinæ.
Curcumæ.

Petaſitidis.
Carlinæ.

Conveniunt in Hyſterica paſſione, Epilepſia, Ophthalmia, Apoplexia, Lethargo, ſcorbuto, febribus intermittentibus, tuſſi, Urinæ ſuppreſſione, Menſibus, fœtu mortuo lochiiſque pellendis, nec non ſtrumis tollendis. doſ. ʒß. in infuſo vel decocto à ʒij ad ℥ß.

VALERIANA.

Eſt radix Valerianæ hortenſis, phu folio Oluſatri Dioſc. C. B. P. oblonga, rugoſa, ſuperficie geniculata, digitum craſſa, foris ex cinericio fuſca, intus pallida, numeroſis fibris dotata, ſaporis acris, aromatici, odoris fragrantis & gravis, ad Nardum accedens. [Uſurpatur etiam in officinis Valeriana ſylveſtris major. C. B. P. n. IV. it. Valeriana paluſtris minor. C. B. P. n. VIII.] [Refertur ad gymnomonoſpermas, ſeminibus pappo criſtatis, floribus monopetalis in muſcariis umbellam mentientibus diſpoſitis, foliis conjugatis.]

Radix hæc effodienda, quando folia incipiunt prodire: feles maxime delectantur hac radice.

Acredine ſuperioribus radicibus eſt remiſſior, ſed amarior, odoreque gravior, & nonnihil adverſa, hinc tanquam antepileptica & Uterina celebratur, in imbecillitate viſus ſingulare quid obtinet.

Datur in ſubſtantia à ℈j. ad ʒß. in infuſo à ʒj ad ʒij. ob ſalia fixa commodius in decocto exhibetur, ob ſui adverſitatem in nonnullis movet Vomitum & ſedes. In Epilepſia datur cum ▽ flor. tiliæ, in Uteri vitiis

cum

cum ▽ Matricariæ, cum ſucco limonum data ante paroxyſmum, ſæpe tollit febres intermittentes.

℞. Rad. Valerian. ℥ij ſem. fænicul. ℥iß Camph. gr. x. M. f. infuſ. in debilitate viſus, & nebulis ante oculos, ex vino exhiberi poteſt tam interne, quam externe applicatum.

℞. Rad. Valerian. Serpent. Virginian. ana ℥j Camphor. ʒj Spir. vin. 🜄 ſat. q. f. f. Eſſ. ſ. Tinctur. Bezoardica, contra Epilepſiam ex Obſtructione Uteri, addita Caſtorei ʒiß fit Tinctura Uterina.

℞. Rad. Valerian. ʒß ⊖ Volat. C. C. gr. xv. Camph. g. ij. M. f. pulv. exhibetur ex ▽ Ceraſor. nigr. contra Epilepſiam. [Duobus potiſſimum in caſibus radix Valerianæ laudatur, 1. in viſus debilitate, qua de re legi merentur ea quæ habet Tabernæmont. in ſuo herbario. 2. in Epilepſia. vid. præter plures alios Panarol. Pentec. 1. obſ. 33. qui Epilepticorum optimum remedium nominat. Et Dan. Crüger. C. N. C. Dec. II. ann. 7. obſ. 78. p. 140.

SERPENTARIA VIRGINIANA.

Eſt ariſtolochiæ violæ fruticoſæ foliis, cujus radix ſerpentaria dicitur [it. Ariſtolochia, Piſtolochia, ſeu Serpentaria Virginiana caule nodoſo] radix tenuis, fibroſa, vincetoxico vulgari candidior, plures fibræ ex uno capite prodeuntes, ſaporis acris incidentis & amaricantis, odoris & ſaporis vincetoxico & valeriana fragrantioris & gratioris. [Figuras exhibet Plukenet. Phytogr. Tab. 15. 78. 148. conf etiam Hermannus noſter Paradiſ. Batav. p. 260.]

Subtilior & gratior valeriana & penetrantior eſt, hinc in febribus intermittentibus & quartana hodie ce-

lebratur, jungitur Chinæ Chinæ, gentianæ aliiſve, ſtimulatur ⊖e abſynth. ammoniac. additis etiam ſi opus ſit purgantibus.

Sociatur quoque tanquam ſingulare medicamentum contra Apoplexiam & Lethargum, ut & ſcorbutum, ſi adſint febres intermittentes.

℞. Rad. ſerpent. Virg. ℈j Cort. Chin. Chin. ʒß. Macer. in vino abſynth. ad eminent. iij digitorum, capiatur totum ante paroxyſmum, præmiſſis tamen vel purgantibus, vel Vomitoriis.

℞. Rad. ſerpent. Virg. Gentian. ana ℈j ⊖ abſynth. gr. xv. ad febrem tertianam, ſi Vomitus conjunctus exhiberi poteſt cum ſucco limonum vel cum gr. j. Laud. op. vel pil. de cynogloſſa, de nocte vero cum Nuce Moſchata toſta, ſeu ex conſerv. abſ.

℞. Rad. ſerpent. Virginian. Gentian. ⊖ abſynth. ana ℈ß jalap. ℈ß troch. alhand. gr. v. Diagryd. gr. iij cape ex vin. abſynth.

℞. Serpent. Virgin. ʒj. Chort. Ch. Chin. ʒij fol. ſenn. ℥j. affund. vin. pint. ß. vel Cereviſ. q. ſ. quando ægri capiunt de ſenna ℥ß non purgat, ideoque ad ℥j vel ℥iß. radix enim ſerp. virg. aufert vim purgantem ſennæ. [Alexipharmaca eſt, Morſum canis rabidi ſanat, ab Hydrophobia defendit. Certiſſimo & præſentaneo remedio eſt adverſus lethiferos morſus ſerpentis Boicininga dicti. Unde & forte ſerpentaria dicta.]

COSTUS.

Eſt radix oblonga, craſſitie pollicari, dura, lignoſa, cortice craſſo, pallido, medulla denſa, buxei coloris, ſaporis acris, amari, aromatici, odoris fragrantis accedentis ad iridem florentinam.

Tres

Tres ejus dantur ſpecies, arabicus, niger & buxeus, inter quas tantum vulgaris albus officinalis eſt, violaceo odore inficiens urinam. [Nonnulli Canellam albam, pro vero Coſto arabico habent, quem propterea Coſtum corticoſum, vel ventricoſum nominant. Sed ſpeciem Coſti non eſſe Paull. Hermann. in Colleg. MSt. oſtendit, allegante ita Mich. Bernh. Valentin. Muſ. Muſeor. l. 1. ſ. 2. c. 17. p. 175.]

Dulcis ſpeciebus cæteris uſitatior, ſalia ſua utraque fovet in blanda quadam glutinoſitate, à qua temperatur, & proinde inter expectorantia refertur, ſuumque opus tam per diaphoreſin, quam per diureſin abſolvit.

Ob odorem cerebro haud moleſtus Cephalicis ac Uterinis incidentibus adnumeratur. Externe vero 🜖 Coſtinum Paralyſi medetur.

Datur in pulvere à ℈j ad ʒß in infuſ. ad ʒij pro ℥viij vini.

In difficili expectoratione cum tuſſi & dolore capitis.

℞. Rad. Coſt. helen. ir. florent. ana ʒij imperat. ʒj paſſul. min. ℥j f. infuſ. ex vino, vel bland. decoct. ex 🜄 Mulſa.

℞. Raſ. lign. ſaſſafr. ℥j Cort. tamariſc. ℥ij rad. ir. florent. coſt. helen. ana ʒvj bacc. junip. ℥j f. decoctum ad aquas hydropicas per diureſin evacuandas.

CYPERUS LONGUS ET ROTUNDUS.

Cyperus longus, eſt Cyperi odorati radice longa, ſive Cyperi officinarum C. B. P. radix oblonga, tenuis, nodoſa & geniculata, contorta, fractu contumax, foris ſubnigra, intus albicans, ſaporis ſuavis ſubacris & aro-

matici paululum exſiccantis, odoris fragrantis, ad nardum accedentis.

Cyperus rotundus, eſt Cyperi rotundi orientalis majoris C. B. P. radix rotunda, olivæ magnitudine & figura, foris rufa, ſcabra, intus alba, pluribus ex uno capite tenuibus quaſi filis dependentibus, ſaporis & odoris ejuſdem cum Cypero longo.

In Germania etiam reperitur Cyperus rotundus [Cyperus rotundus vulgaris C. B. P. n. iij.] ſed nullarum eſt virium. Cyperus longus præſtantior eſt rotundo, congruit cum galanga, ſed non tam acris eſt & plus habet adſtringentiæ, hinc Ventriculum à relaxatis fibris debilitatum mire confortat, & eandem ob cauſſam ingreditur gargariſmata ad mobiles dentes firmandos.

[Eſt planta gramini folia, flores ſtaminei diſpoſiti in juba, caule triangulo.]

Datur in ſubſtantia ad ʒß in decocto & infuſo velut aliæ radices.

℞. Rad. Cyper. galang. cubeb. ſummit. abſynth. ana ʒß. ſantal. r. ℈ß ℅ cinamom. gt. iiij, irroretur ♎u ⊕li, extrahitur ad debilem Ventriculum cum Vomitu à relaxatis fibris.

℞. Rad. Cyper. ir. florent. ana ʒij Nuc. cupreſſ. ℥ß. gland quern. ʒj flor. balauſt. Mj. malicor, ʒiij alum. ʒj coq. ex ▽ ferrariorum q. ſ. add. ſyr. myrtin q. ſ. f. gargariſma ad firmandos mobiles dentes, in cauſſa ſcorbutica, & uvulam relaxatam confirmandam.

SPICA NARDI.

Eſt vel Indica vel Celtica. Spica Nardi Celtica eſt ſpecies Valerianæ, vid. Cluſ. [rarior. plantar. hiſtor. l. 4. c. 39. p. 57.] Spica

Spica Nardi Indica ſeu potius Gangetica; [Nardus Indica, quæ Spica Nardi & Spica Indica officinis C. B. P.] eſt radix oblonga, digiti craſſitie & longitudine, comoſa, quaſi ex marcidis foliorum nervis vel nervoſis filamentis convolutis congeſta, colore ex fuſco rubeſcente, ſaporis amari, acris aromatici, odoris Spica Celtica gratioris ad Cyperum accedentis.

Spica Celtica ſunt Nardi Celticæ vel rectius Valerianæ Celticæ [Nardi ſpuriæ Narbonenſis C. B. P.] radices fibroſæ, capillares, nigricantes, cum foliis parvis ex viridi flaveſcentibus, ſaporis acris amaricantis, odoris fragrantis graviuſculi.

Nobilior eſt Spica Indica ac Celtica, virtute ſuperans Cyperum & Valerianam, ob odorem penetrantiorem & graviorem & ⊖ia penetrantiora. Utero & capiti eſt dicata.

Exaltatur ejus vis Caſtoreo & Camphora, aſarum ſpecifice corrigitur Spica Nardi.

Datur in infuſo & decocto velut aliæ radices.

℞. Spic. utriuſque ana ʒ ß. aſari ʒij in robuſtis ʒ iij vel iiij coquatur in vino vel Cereviſia leviter, ad Menſes, fœtum mortuum & lochia pellenda.

℞. Spicæ utriuſque, Valerian. ana ʒj Camphor. gr. v. Caſtor. gr. viij. f. pulv. antihyſtericus cum vino enulato, cum ▽ chærefol. matricariæ in obſtructionibus Meſenterii; ubi eſt Cachexia cum tabeſcentia Corporis una cum Curcumæ ʒj præſcribitur.

Indi habent ☍ Nardinum egregium nervinum.

VINCETOXICUM.

Sunt Aſclepiadis flore albo C. B. P. radices albicantes, tenues, fibroſæ, plures ex uno capite prodeuntes,

tes, ſaporis acris, amaricantis, nonnihil nauſeoſi, odoris graviuſculi. Una tantum ſpecies eſt officinalis. [Refertur ad plantas ſiliquoſas unicapſulares polyſpermas, ſiliquis poſt ſingulos flores monepetalos quinquevariam diviſos, ſingulis univalvibus ſuperiore parte hiantibus, unico alveolo ſemina plura papposa lanugine criſtata & multiplici ſerie imbricatim invicem incumbentia continente, inſtructis.]

Valerianæ & Nardo accedit, aſt viribus longe eſt remiſſius, fovet quoque aliquid nauſeoſi, hinc Vomitum levem nonnullis movet, præcipue ſi ſit recentius.

In incidendo Pulmonum muco, in tollendis ſtrumis cæteriſque tumoribus per Urinam educendis virtutem habet. [Vincetoxicum ſcrophularum eſſe remedium appropriatum, dum maxima vi penetrandi præditum, ſuaque ſubtilitate lympham incraſſatam attenuat, ut per glandulas commode tranſcolari denuo poſſit, prolixe probat Joachim. Georg. Elſner. C. N. C. Dec. I. ann. I. obſ. 57. p.m. 153. hanc ob cauſſam etiam in hydrope curanda laudatur, videri meretur Georg. Tob. Dürr. C. N. C. Dec. II. ann. 7. obſ. 58. p. 105. & Caſp. Hoffmann. ap. Laurent. Scholz. conſil. 337. p.m. 951.]

Datur in ſubſtantia ad ʒj decocto & infuſo ab ℥ß ad ℥j.

℞. Vincetoxic. ℥j ▽ Mulſ. ℥iij vel iiij coquatur ad formam ſyrupi, poteſt exhiberi infanti in tuſſi ſtomachali, ubi evacuatio per vomitum locum habet.

℞. Vincetoxic. ſerpent. virgin. Valerian. ana ℥j raſ. lign. ſaſſafras. bacc. Junip. ana ℥ß. Macera in vino, poſt leniter coquatur. ad ſtrumas & Urinas promovendas.

Contra

Contra coſtarum dolorem Poterius [Cent. 1. cap. 83.] ſequenti modo uſus fuit vincetoxico.

℞. Rad. vincetoxic. ℥iß. folior. myrti Mj. ſem. hyperic. rhabarb. elect. ana ʒiß f. decoct. in ſ. q. 🜄æ communis, in cujus colaturæ ℥vj diſſolv. ſyr. capill. ♀is ℥iß.

HELENIUM.

Eſt Helenium vulgare C. B. P. radix oblonga, craſſa, foris fuſca, intus pallida, pinguis, ſaporis acris amaricantis aromatici, odoris fragrantis.

[Annumeratur plantis floribus compoſitis diſcoides, pappeſcentibus non lacteſcentibus.]

Helenii ⊖ia pinguedini & reſinæ incluſa ſunt, hinc in Colica, Paralyſi, ſcabie, tuſſi, tremore membrorum, præcipue a Mercurialibus orto conducit, relinquit ob pinguedinem lubricantem vim anodynam & apertam alvum.

Ob analogiam quam habet cum Bile, ſupplet ejus defectum, & vitia emendat.

Datur commode cum 🜄 propria deſtillata, quæ teſte Febure [Chym. part. 2. ſect. 1. pag. m. 290.] ⊖e volatili copioſo eſt impregnata & liquorem ſ. ⊖ C. C. æmulatur: ſeu vino enulato.

N. ſat fragrans 🜄 ex helenio deſtillatur, ſi non 🜄 in ſuperflua quantitate addatur.

In Chloroſi ſpecificum medicamentum audit, condita radix vel per ſe, vel in tabulis vel in morſulis, vel in decocto pro expectorante. In ſcorbuto ſtimulatur Cortice Winterano. In Colica flavedine corticum aurantiorum. Externe tam ejus decoctum, quam unguentum laudatur in ſcabie. Vis plurima

concentrata in ejus reſina videtur, quæ & in Uteri ruptura ex partu difficili ſpecifica audit.

℞. Rad. helen. flaved. Cort. aurant & Citri ana ℥j coq. in vino, decocto add. Cort. Winteran. ʒiß, datur in ſcorbuto, cum dolore colico & Paralyſi.

℞. Helen. condit. Zinz. Chin. cond. ana ℥iiij paſſul. Corinth. ℥ij f. decoctum ex ▽, in tuſſi chronica commendatur.

℞. Rad. helen. bardan. fœnicul. ana ℥iiij f. ex ▽ decoct. contra tremores artuum ex mercurialium medicamentorum ſumtione.

℞. Gumm. helen. ℥ij thereb. ℥iij ☍ hyperic. ℥iiß Myrrh. ſang. Dracon. ana ʒj M. f. Balſ. quo inungantur pudendorum rupturæ poſt partum.

IRIS FLORENTINA.

Eſt Iridis albæ Florentinæ C. B. P. radix oblonga, tuberoſa, digitum craſſa, paululum compreſſa, ſaporis amaricantis, acris, ſubpinguis, odoris fragrantis violacei.

[Annumeratur plantis enneapetalis tricapſularibus, foliis enſiformibus cauli obverſis, radice tuberoſa.]

Iris Florentina Enula mitior eſt & jucundior, ſed pinguior, guſtu tamen in receſſu nonnihil faſtidioſa, præcipue ſi recens ſit, hinc alvum puerorum magis lubricat, & certis in conditionibus Vomitum movet. In tuſſi quoque convenientior & moderatior eſt, movens ſimul Urinam, eamque violaceo odore inficiens, paſſiones torminoſas & pavoroſas infantum compeſcere & ad ſomnum diſponere dicitur, ſi cum Croco exhibeatur.

Ob-

Ob ſubſtantiam levem, farinaceam & poroſam abſorbet, abſtergit, hinc oſſibus denudatis & capillis humentibus inſpergitur.

℞. Ireos florent. ℈j. Croc. or. gr. v. ſem. fæn. ℈ß M. f. pulv. pro 2 doſ. exhibetur cum ▽ Mulſa vel cæteris paribus etiam cum oxymell.

Exhiberi poteſt infantibus in torminibus abdominis cum alvo adſtricta & noctes ſimul inquietas habentibus.

℞. Rad. ir. flor. gr. xv. oxymell. ʒiij datur infanti pro leni purgante.

℞. Rad. ir. flor. ʒß exhibe ex ſyr. violar. ℥ß ſacch. alb. ʒj ▽ Borragin. ℥iij. dari poſſet in raucedine.

℞. Rad. ir. florent. ariſtoloch. rotund. ana, in oſſibus denudatis, ſi caries ſimul adſit. ℞. Rad. ir. florent. ʒj. Euphorb. ℈j [addi poſſ. 🝆 d. caryoph. gt. iiij.]

CURCUMA.

Vocatur & Crocus terreſtris. [Indicus] Eſt radix oblonga, parva, tuberoſa, nodoſa, intus & extus crocei coloris, ſaporis ſubacris, ſubamaricantis, pinguis, odoris ad Zinziber accedentis, aſt debilioris. [Quibuſdam dicitur radix Zinziberis facie intus flava.]

[In horto Acad. Lugd. Batav. radicem ita deſcribit: conſtat radice tuberoſa, tereti, in tranſverſum acta, Zinziberis facie, craſſitie digiti, ex multis propaginibus, cum circularibus in ſuperficie Zonis geniculatim conflata, aliquot craſſiuſculas fibras ex geniculis demittente, ponderoſa, foris pallida & modice ſcabra, intus crocea, ac temporis decurſu in puniceum inclinante, ſolidam quaſi ex condenſato croceo ſucco preſ-

preſſius compacta, guſtus acris amaricantis aromatici pinguis, odoris fragrantis.]

Duæ ſunt ſpecies prima radix longa, altera radix tuberoſa [& rotunda] illa, maxime in uſum venit.

Figura, quæ extat in Bontio, nihil valet, eſt enim Cannæ Indicæ, in iis etiam locis naſcentis, ubi Curcuma provenit, quæ forſan cauſa erroris fuit. [in Hort. Lugd. Batav. ita noſter: folia curcumæ, foliis cannæ indicæ junioris adeo ſimilia, ut à rudioribus non facile ab invicem diſcerni queant. Hoc errore ductum video Bontium, qui Hiſt. Nat. & Med. l. 7. c. 30. loco curcumæ delineavit cannam indicam in India frequentem, & hortis noſtris ſatis familiarem.]

[Sub eodem genere ſummo merentur reponi, & ſimili omnino fructificatione gaudent, Bangala Indorum, cardamomum majus & minus, Coſtus Arabicus, utraque Galanga, Zedoaria, Zerumbeth Garciæ, Zinziber: Hermann. ibid. Notæ ſunt radix tuberoſa, folia lata oblonga, alternatim poſita, flores tetrapetali difformes, vaſcula ſeminalia tricapſularia.]

Matureſcit foditurque radix poſtquam flores exaruerunt.

Curcuma pinguior eſt, quam iris florentina, aſt virtute aromatica gratior & penetrantior. [Donatur ſale oleoſo volatili, partibus blandis viſcoſis immerſo, id. ib.] Hinc tanquam polychreſta ab Indis inter ciborum aromata eſt recenſita ſeu recepta.

In ictero & vitiis Bilis ſcilicet defectu & inertia ejus, felicius datur quam helenium.

Ex noſtris obſervationibus in facilitando partu ſingularis eſt.

Datur

Datur ut iris florentina & helenium, frequentius in ſubſtantia, quam infuſo. Extra ſub forma unguenti applicata medetur nervorum doloribus. Sternutatoriis quoque admiſcetur.

℞. Curcum. ℥ß. Croc. or. gr. v. ⊖ volat. C. C. gr. vij. M. f. pulv. exhiberi poteſt in ictero, orto à Bile viſcoſiore & minus acri. vel

℞. Curcum. ℥ß ⊖ abſynth. ℈j M. f. pulv. contra icterum, exhibendus in ▽ Chelidonii majoris.

℞. Curcum. ʒj Croc. or. ℈ß. Laud. op. gr. j. M. f. pulv. ad part. facilitandum, cui addi poterunt Caſtorei gr. v. & radix Coſti nigri ad ʒß.

PETASITES.

Eſt Petaſitidis vulgaris & majoris C. B. P. radix oblonga, digitum craſſa, ramoſa, reſinoſa, foris fuſca, intus albicans, ſaporis acris aromatici, amaricantis, odoris fragrantis.

[Refertur ad gymnoſpermas, floribus compoſitis, pappeſcentes & lacteſcentes.]

Salia ſua penetrantia & ſpirituoſa in reſina fovet, quare aſthmaticis & hyſtericis convenit.

In Morbis malignis & peſte excellens eſt alexipharmacum, [unde & Peſtilentz-Wurtzel appellatur] & propterea Theriacalia, Tincturas Bezoardicas ut & aquas & aceta prophylactica ingreditur.

Stimulatur Zedoaria, Imperatoria, maxime vero Camphora, vis conſiſtit in reſina, vel cortice reſinoſo, qui ad ʒj ex vino vel aceto exhibetur, datur etiam in extracto per Spir. Vin. facto.

℞. Petaſit. angel. Zedoar. Valerian. ana ʒj affund. Spir. Vin. Camphorat, vel acet. deſt. ad eminent. ij

vel iij digitorum, f. Eſſent, vel tinct. alexipharmaca, cochleatim, ſingulis diebus capienda, contra peſtem.

℞. Petaſit. helen. ana ℥ij liquirit. ℥j. ficuum n. viij f. cum aqua & vino decoct. pectorale, ubi metus malignitatis ſimul ſubeſt.

CARLINA.

In Sylvis Miſniæ & Hercyniæ provenit. Eſt Carlina acaulos magno flore C. B. P. radix oblonga, digiti craſſitie, ut plurimum in ſuperficie carioſa & foraminulenta, colore foris rufo, intus albido, ſaporis acris, odoris gravis & fragrantis.

[Refertur ad herbas pappoſas non lacteſcentes, flores perfecti compoſiti diſcoidei.]

Raro præſcribitur à Medicis, forſan ob ignorantiam ejus uſus.

Carlina Meo, Leviſtico & Pyrethro quodammodo reſpondet, ſed ingratior eſt & penetrantior. Ob nimiam fragrantiam multis moleſtum Capitis dolorem vertiginem & nauſeam infert. Helmontius etiam vult ſomnum impedire. [De ea agit Helm. in tumul. peſtis ſub finem: cujus verba hæc ſunt: quicquid in peſte agendum eſt, id feſtinanter curandum eſt. Ideoque ne tenax intus hoſpitium, Peſtis ſibi ſumat, procurandum: ut imago non dudum intus fluctuet peſtilens, ſed tota ſuppellex foras per ſudorem proliciatur & excidat. Carlina enim Imperatori (Carolo Magno, unde & Carolina ſeu Carlina dicta) ab Angelo oſtenſa dicitur pro peſte exercitus, forte ideo dicitur Carduus angelicus, quod primus ortus imaginis peſtilentis circa ſtomachum ſopores, nauſeas febrem vomitum & cephalalgias ſuſcitet. Ixia (ita vocatur ſuccus Carlinæ

næ vel viſcum radici illius herbæ adhærens,) vero ſomnum arceat, multoque magis ſopores contra naturam. Ideoque extractum Carlinæ recentis non fore infœcundum recenti peſti ſperant.]

Interim Hyſtericis quibus odor non infenſus eſt mire conducit, ut & in peſte præſervanda. Potentiſſime movet ſudorem, eoque nomine Herpetem cutaneoſque affectus ex fundamento curat, per ipſum enim ſudorem expellit acidum illud corroſivum herpetis.

Jungitur Pyrethro in infuſis & decoctis ad Paralyſin, præcipue Linguæ.

Decoctum etiam commendatur ad dolores Hypochondriacos. Vomitum quoque quibuſdam moviſſe teſtis eſt Hoffman. in Clave, [inquit: Carlinæ decoctum quibuſdam etiam Vomitum movet in brodio carnis factum, ut non raro obſervavi in praxi.] In decocto datur doſi parciori, in infuſo largiori.

℞. Carlin. ℥j angelic. ʒij Caſtor. ʒß. Camphor. ℈ß. M. f. pulv. pro vj doſibus ad paſſionem hyſtericam.

℞. Carlin. Pulveriſ. ʒß. ad ℥j ſingulis diebus contra herpetem, ſeu cape ejus decoctum. vel

℞. Carlin. ʒij Cort. lign. ſaſſafr. ʒiij f. pulv. ad vj doſes.

In Cachexica icteritia: ℞. Carlin. ℥j ⊖ vol. C. C. gr. ix. f. pulv. exhibendus in ▽ deſtil. Chelid. Maj. [De radicis Carlinæ vi defatigante conferri meretur Eman. König. C. N. C. Dec. II. ann. I. obſ. 169. p. m. 401.]

RADICES

***

RADICES CONSTANTES SALE AROMATICO BLANDO SEU MITIORI, CONSTANTES VIRTUTE DIURETICA APERIENTE, ET BLANDE SUDORIFERA.

Fœniculum.
Apium paluſtre.
Petroſelinum.

Hæ radices rariſſime ſudorem movent, utplurimum Urinas.

FOENICULUM.

Eſt fœniculi vulgaris Germanici C. B. P. radix oblonga, recta, digitum craſſa, pallida, ſaporis ſubdulcis, ſeu cortices hujus radicis, ſubacris, aromatici & tenuis, odoris fragrantis.

[Pertinet ad umbellatas ſeminibus nudis binis ſequentibus ſingulos flores, ut & ſequens apium paluſtre, & petroſelinum.]

Fœniculum ob ſalia dulci viſcoſitate temperata, Urinam movet & nonnunquam ſudores, ſpecificum eſt contra Variolas & Morbillos. Confert quoque ad promovendam Chylificationem, & lactis generationem, nempe incidendo, & per Urinam educendo acida viſcida, à quibus generatio lactis impeditur. N. Cave ne ad lactis generationem medicamenta adhibeas nimis Urinam pellentia, tum enim materiam lacti dicatam ad renes & per Urinam protrudunt.

Tollit quoque tremores membrorum à Mercurio ortos, velut helenium & bardana.

In

In uſum trahendus potius cortex radicis quam interior ſubſtantia, datur potius in decocto quam in ſubſtantia.

In Variolis & Morbillis juvatur ſcordio, ſtimulatur ⊖e C. C. in diureſi vero ligno & baccis juniperi.

℞. Rad. fœnicul. ℥iiij ſem. fœnic. ℥j f. decoct. ad lactis generationem in mammis. vel

℞. Rad. fœnicul. ℥ij ſaxifrag. ℥ß. flor. ſambuc. Mj ſummitat. aniſ. vel fœnicul. p. j.

℞. Rad. fœnicul. vel potius Cort. ℥ij ſcord. Mß vel j. lent. excort. ℥ß. caric. ping. n. v. M. f. decoct. ex quo exhib. ⊖ C.C. ℈ß. contra Variolas & Morbillos, poteſt addi Caſtor. ad gr. vij. quod in Morbillis ſingulariſſimum.

℞. Cort. fœnicul. ℥iß. vel ij bacc. junip. ℥iij lign. junip. raſp. ℔j macera, & coque leni △e, cui adde ſyr. radic. 5 aperient. ad gratiam, ⏝us ⊖is dulc. quantum videtur.

APIUM PALUSTRE.

[Eſt apii paluſtris & apii Officinarum C.B.P. radix pollicem craſſa, albicans, à craſſiori capite in numeroſas fibras divaricata, ſaporis ſubacris, odoris fragrantis,]

Apium fovet aliquod volatile fumoſum, capiti adverſum, veluti Sium aquaticum quod Epilepſiæ obnoxiis aliquando nocet, diureſin potentius excitat quam fœniculum, & propterea pellendo materiam lactis ad renes, ejus redundantiam minuit & coagulum reſolvit.

Datur in decocto cum congeneribus, & applicatur quoque externe ſub forma Cataplaſmatis.

Semper debet eſſe recens, aliàs nihil valet.

℞. Rad. apii ℥ij petroſel. ℥j ſem. apii petroſel. ana ℥j fol. apij Mij ſem. cumin. ℥iß coque in vino & aceto in formam cataplaſmatis, quod applica mammis ad lactis redunduntiam, add. ſperma ceti, ſi lac ſit coagulatum.

PETROSELINUM.

[Eſt Apii hortenſis ſeu Petroſelini vulgo C. B. P. radix oblonga, recta, craſſa albida, fibris lateralibus donata, ſaporis ſubacris & aromatici, odoris fragrantis.]

Recentiſſimum debet eſſe. Multi Epileptici non ferunt ejus odorem. [Hinc etiam Tabernæmontanus mulieribus gravidis, puerperis & lactantibus Petroſelini uſum prohibet, ne infantes Epilepſia, vel motibus convulſivis corripiantur. vid. l. 1. ſect. 3. c. 25. p. m. 266.]

Petroſelini radix validior eſt Apio, de cætero cum hoc convenit.

Datur ut apium.

℞. Rad. Petroſel. ℥iij ſem. petroſel. ℥ß. fol. petroſel. Mj. rad. urtic. min. ℥i f. ex ▽ decoct. diureticum, quod ſi nimis & cum dolore pellat urinam, add. rad. ſymph. & althææ ana ℥ij vel iij.

***

RADICES SALE VOLATILI AROMATICO SPIRITUOSO ET PENETRANTE ET PAUCO ACIDO VOLATILI CONSTANTES.

Et proinde tam ſudoriferæ, quam diureticæ, acidum rancidum ſcorbuticum corrigentes.

Armo-

Armoracia ſeu raphanus ruſticanus.
Arum.
Scylla.
Pyrethrum Italicum & Germanicum.

ARMORACIA SEU RAPHANUS RUSTICANUS.

[Eſt raphani ruſticani C. B. P. radix oblonga, pollicis craſſitudine ac etiam craſſior, colore albo, guſtu acerrimo, ſinapi haud abſimili, odoris nullius.

Pertinet ad plantas ſiliquoſas bicapſulares cum ſepto intermedio, flore tetrapetalo.]

Armoracia ratione ſui ſalis penetrantis & ſpirituoſi, juncto pauco acido volatili, ⊖ia ſylveſtria quævis, acido viſcido putridove muco immerſa tam per diureſin, quam per diaphoreſin educit, & proinde in ſcorbuto ab acido rancido oriundo, it. in Paralyſi & morbis ſoporoſis ſpecifica eſt. & in appetitu tam proſtrato quam aucto ſingulariſſima eſt.

In ſoporoſis affectibus plantis pedum ſub Cataplaſmatis forma applicatur. Additur quoque diureticis loco ſtimuli. Juvatur in ſcorbuto cochlearia, trifolio fibrino, naſturtio, cortice Winterano.

Stimulatur aro, ſem ſinapi, erucæ, aliiſque acribus.

Temperatur ſero lactis, Cichoreo, ſucco aurantior. Citri, paſſulis &c.

Nunquam datur in decocto, aliàs ſalia volatilia evaneſcunt. Datur ejus raſura in Electuariis & in infuſo, cum ſero lactis vel vino. Vel datur ejus ſuccus ad ℥j.

℞. Rad. armorac. raſp. Mß. fol. cochlear. Mj. Cort. Winter. ʒij Vin. alb. q. ſ. vel ℔ j f. infuſum, poteſt exhiberi in caſu ſcorbutico cum dolore artuum à frigore orto. doſ. ℥v. l. vj bis de die.

Raphanus ruſticanus ſi menſæ apponitur, tum corrigitur raſura amygdalarum dulcium, & ſaccharum ſuper inſpergitur.

℞. Rad armorac. trifol. fibr. Naſturt. rec. ana Mj. Cort. Winteran. ʒiij f. infuſum ex vino vel ſero lactis.

℞. Raſ. rad. armorac. p. ij flaved. Cort. Citr. ʒij ſucc. cochlear. ℥ß. ſyr. aurant. q. ſ. f. Elect.

RADICES SALE VOLATILI ACERRIMO ABSQUE OLEO GAUDENTES.

ARUM.

Eſt Ari vulgaris C.B.P. radix rotunda, tota alba, nucis avellanæ magnitudine. Aron ob ſal acre valde ſubtile volatile, cruditates acidas, auſteras & terreſtres viſcidaſve, & ex his natos Morbos potenter diſſolvit, & per diureſin educit. Hinc ſtomachicum & thoracicum excellentiſſimum eſt, junctum acidis volatilibus ſeu ſpirituoſis fit antiſcorbuticum, & ſanguinem congrumatum feliciter diſſolvit.

Quando acetum affundimus, radix effervefcit, ad alcalicum igitur magis accedit, quam cæteræ plantæ.

Ejus extractum [ſeu potius radix cum Spir. Vin. camphorato ſubacta] extra membris applicatur Paralyticis.

Datur ob nimiam ſui acredinem nunquam recens ſed ſiccatum, donec ejus acredo ſit temperatata. Præparatur etiam aceto, quod tamen nimis partes acres invertit. Omnium optima præparatio eſt cum vino, quod

quod ratione blandæ aciditatis non adeo caſtrat particulas ſalinas.

Additur tanquam ſtimulus medicamentis terreis, acida abſorbentibus. Datur præparata à ʒß ad ʒj.

℞. Rad. ari præpar. ʒß f. pulv. ſumendus ex vino rhenano in anorexia.

℞. Rad. ari præpar. C. C. p. p. ana ℈j M. f. pulv. ad tollenda tormina ventris ab acido orta, addi poteſt Laud. op. gr. j.

℞. Rad. ari ʒß acet. deſtil. ℥j datur in caſu ab alto ad ſanguinem coagulatum, addi poteſt ſperma Ceti & oculi 69. Pulveres ad caſum ingreditur loco ſtimuli.

℞. Rad. ari p. p. ʒj ireos flor. ʒß. flor. 🜍 benz. antihect. Pot. à ℈ß. M. f. pulv. contra vitia Pulmonum doſ. ℈j.

℞. Rad. ari p. p. ʒß ▽ deſtil. ari cochl. j. exhibe ad excitandos ægros in Morbis ſoporoſis.

Diuretica fit maxime, ſi ipſis conjungitur.

SCILLA.

[Eſt Scillæ vulgaris radice rubra C. B. P. radix bulboſa magna, pugnum craſſa, cœpacea ex pluribus tunicis ſeu lamellis compacta, inferius multis craſſis fibris prædita, ſaporis acris amari, odoris nullius.

Refertur ad plantas tricapſulares hexapetalas, floribus in ſpicis longa ſerie diſpoſitis.]

Scilla præter acredinem cum aro communem poſſidet amaritiem ingratam, & nonnihil nauſeoſam, hinc præter virtutes aro communes movet Vomitum & alvum. Velut arum in Pituitoſis affectibus opportune exhibetur.

Cruda eſt valde acris, exurens fauces, hinc ſiccatur in Sole & clibano, præparatur ut arum, in ſubſtantia nunquam datur.

Datur in infuſo cum melle, ▽ Mulſa, item ex aceto. Hinc proſtat acetum, mell. & oxymell. ſcillitic. loch de ſcilla, in qu bus omnibus exhibetur. [vid. Geſnerus in epiſtolis ſuis hinc inde.]

Pro vomitu citando, acuitur ♁iatis, aſaro, temperatur ☍ amygd. dulc. [in aſthmate acetum ſcyllitic. optime jungitur cum ſyrupo violarum.]

℞. Oxym. ſcill. ℥j Vin. ſcill. ℥ij M. f. vomitorium blandum, addi poteſt ☍ amygd. dulc. ℥ß.

℞. Oxym. ſcill. ℥j M. f. vomit. cui correctorii loco additur ▽ Cinamom. ℥ß.

RADICES SALE ACRI FIXO ABSQUE OLEO CONSTANTES.

PYRETHRUM.

Pyrethri duæ ſunt ſpecies, Italicum & Gallicum. Eſt Pyrethri flore bellidis C.B.P. radix tenuis, oblonga, foris fuſca, intus pallida, [ſaporis acris fervidi, odoris nullius.]

Utrumque pyrethrum ob acredinem ſuam Linguæ pertinaciſſime inhærentem & fontes ſalivales aperientem, Salivam copioſe educit, hinc in odontalgia ab obſtructione & Catarrhis orta ſpecificum. Ob eandem penetrantem acredinem nervos vellicantem, eorumque obſtructiones aperientem, in Morbis ſoporoſis & aphonia efficaciſſimum eſt.

Datur ejus ſubſtantia ad ʒß additur quoque decoctis, infuſis clyſteribus. jungitur ſaxifrag. aro, ſcillæ. Carlin. Colocynth. ciclam. &c.

℞. Rad.

℞. Rad. pyrethr. ʒß ſucc. abſynth. q. ſ. f. turunda intrudenda cavo dentis, contra dolores ortos vel à Catarrhis, vel vermi us. vel

℞. Pyrethr. ʒj ☍ origan. gt. v. Camph. gr. x. mucilag. tragacanth. q. ſ. f. turunda cavo dentis intrudenda.

℞. Rad. Pyrethr. utriuſque ſaxifrag. rec. ana ℥j coq. in ▽ ſeu vino, affundæ ▽æ deſtill ari & ſcillæ q. ſ. vel infunde ari & ſcillæ ana ʒj. f. apozema, detinendum in ore, in loquela amiſſa.

℞. Rad. Pyrethr. utriuſque ℥j ſummit. orig. Mj. pulp. colocynth. in pet. lig. ʒj coq. in ▽ add. colatur Hier. picr. ℥ß Tinct. Caſtor. ℈ij ⊖ vol. ſuccin. ʒj vel ⊖ gemm. ℈ij f. Clyſter acris ad excitandos Apoplecticos.

RADICES SALE ACRI ABSQUE OLEO TAM FIXO QUAM VOLATILI CONSTANTES.

Pimpinella Saxifraga.

Eſt Pimpinellæ ſaxifragæ majoris umbella candida C.B.P. radix candicans tenuis, emittens ex capite craſſiori nonnullas fibras, ſaporis acris, odoris nullius. Sponte proveniens, habet majorem acrimoniam præ illa, quæ in hortis colitur.

Saxifraga pyrethro cui propemodum accedit remiſſior eſt, aſt ob acredinem quodammodo piperatam ſalubre ciborum aroma.

Præterqnam quod in Morbis ſoporoſis cum pyrethro convenit, diureticum inſigne eſt, & juncta Uterinis anodyna, dolores poſt partum tollit, eorumque materiam peccantem educit per Urinam.

Contra tremores artuum à ☿ialibus ortos laudatur una cum fœniculo. Datur in decoct. infuſ. & Clyſter. in ſubſtantia ad ʒj. Schenkius

Schenckius obſervat quod mammis applicata copioſe lac generarit, ſed non multum fidei adhibetur huic obſervationi. Hujus ▽ deſtillata conveniens vehiculum medicamentorum diureticorum.

Radix ſaxifragiæ exhibetur cum extracto Caſtorei in ſanguine congrumato & viſcido ex lochiis retentis.

In Hydrope & difficultate Urinæ ſequens convenit remedium: ℞. Rad. 5. aperienr. ʒiij ſaxifrag. ℥ij cort. fraxin. rec. ℥j bacc. junip. ℥ß. f. decoct. ex vino.

Radix Graminis.

Non graminis cujuſvis, ſed graminis canini arvenſis ſeu graminis Dioſcoridis C.B.P. Germ. Queckgraß. [Radix tenuis admodum, longa, geniculata, alba, ſaporis ſubdulcis, cum aliquali in receſſu acrimonia, odoris nullius.]

Temperatior videtur ſaxifragia, ideo non tantum lene eſt diureticum, ſed in atrophia & affectione hypochondriaca deobſtruens eſt, laudatur quoque tanquam anthelminticum.

Ingreditur infuſa & decocta, nunquam vero exhibetur in ſubſtantia. Aqua deſtillata appropriatum menſtruum eſt & vehiculum diureticorum, deobſtruens anthelminticum, ſtimulatur ſaxifragia. In vermibus jungitur filici, oſmundæ, ſemini ſantonici aliiſve.

℞. Rad. Granum. ℥ij urtic. min. c. tot. Mj. f. decoct. in atrophia & macie, ubi materia peccans educenda per Urinam.

℞. ☿ dulc. gr. v. ſi ſit puer 6. vel 7. annorum ▽ gramin. deſt. cochl. ij. Contra vermes puerorum.

℞. ☿

℞. ☿ viv. ℥j affund. ▽ deſt. gramin. pone in digeſtione per 24. horas, decanta ▽ & exhibe contra vermes, ſingulis diebus.

℞. Thereb. Venet. ℥ß. vitell. ov. q. ſ. Cum ▽ gramin. ℥vj f. Emulſ. diuretica in iſchuria renum, addi poteſt Tinct. Caſtor. ʒß Mell. ℥j. ad ℥ß Thereb. addi poteſt ſucci limonum ℥ß.

***

RADICES SALE BLANDO BALSAMICO CONSTANTES.

Salſaparilla.
China.

Sunt Smilacis aſperæ Peruanæ ſive Salſæparillæ C.B.P. radices tenues, calami craſſitie, longæ, ſarmentoſæ, plicatæ & rugoſæ, ex flavo pallidæ, intus candidæ, ſaporis & odoris nullius evidentis niſi farinacei.

Chinæ duæ ſunt ſpecies 1. Chinenſis 2. Africana ſeu Pſeudochina. Eſt Smilacis aſperæ Chinenſis Lampatam dictæ radix craſſa, tuberoſa, lignoſa, aliquantum reſinoſa, pallide rubeſcens, intus candida, ſaporis terreſtris, leviter adſtringentis.

Salſæparillæ & Chinæ effectus probat ſal blandum, quod in farinacea ſalſæparillæ ſubſtantia, & in ſubſtantia chinæ nonnihil reſinoſa latet, & in utriuſque radicis decocto manifeſte ſe prodit. Hoc cum tenuiſſimarum partium, & ſanguini noſtro naturali valde analogum ſit, ſalia ſeu ſpiritus non tantum exaltat, ſed & acida quævis pertinaciſſima etiam, remotiſſimis partibus inhærentia, à quibus Lues ♀ea, arthritis, ſcorbutus, atrophia, phthiſis aliique Morbi

 oriun-

oriuntur, infringit, & cum blanda vi anodyna per diaphoreſin expellit.

Radix Chinæ penetrantior & ſpirituoſior quam ſalſæ parillæ, utraque prius maceranda, in decoctis, lento △e factis vaſe probe clauſo, exhibetur, ſpecificatur foliis chamæpitis, & in variis morbis exhibetur.

℞. Rad. Chin. ſalſæparil. ana ℥iiij Chamædr. Chamæp. ana Mj. ſi dolores intenſiores add. flor. chamom. p. iij rad. liquir. ʒij vel iij datur in arthritide & podagra.

℞. Rad. Chin. ℥j coq. in ▽æ ℔ij ad medietatem f. apozema, add. fol. chamæp. pugillos aliquot, exhiberi poteſt in morbis chronicis contumaciſſimis.

℞. Rad. Chin. ℥j raſ. C.C. turion. ſ. conos abietis. ℥iß coq. in ▽ ſ. q. f. decoct. in ſcorbuto cum Paralyſi.

℞. Rad. Chin. ſalſæparill. ana ℥ij Cort. lign. Guajac. ℥iß vel ij lign. ſaſſafras ℥j add. ☿ viv. ♁ij crud. in petia ligat. ana ℔ß. coq. invicem f. apozema in Morbis ♀eis, ſingulis diebus bis vel ter ſumendum.

RADICES CONSTANTES SALE VOLATILI, BLANDO CUM TEMPERATA TENUI VISCOSITATE.

Orchis ſeu Satyrium.

Orchis eſt vel rotunda ſeu teſticulorum forma, vel palmata manuſve referens. Utraque ejuſdem virtutis. Sunt radices rotundæ, ſucculentæ albidæ compreſſæ paululum, viſcoſæ, ſubdulces, odoris ſeminalis.

Utriuſque orchidis vim aphrodiſiacam ſeu ♀em ſtimulantem docet Experientia & Ratio, à ſubſtantia ſubdulci & viſcida ſubminiſtrari ſemini generando materiam, ab ejus ſale blando volatili, ſperma ſeu Uri-

nam redolente excitari orgaſmum in ſemine, & titillaonem in partibus genitalibus.

[Refertur ad plantas tricapſulares flore monopetalo rictu & labio vario conſtante.]

Datur in ſubſtantia vel cruda ad ʒj it. condita, vel in eſſentiam redacta cum ♒u formicarum, addita eſſentia ambræ. Juvatur Eryngio, nuce juglande condita, exterior cortex nucis juglandis optimus eſt, conſtat enim ⊖e aromatico, cujus interior pulpa plane expers eſt; priapo cervi. Stimulatur formicis alatis, ▽ Magnanimitatis, it. ſemine ſinapi, erucæ, ſpeciebus diambr. ♒u roſar. ambrato.

℞. Rad. ſatyr. cond. ℥j Elect. diaſatyr. Nuc. jugl. cond. flaved. cort. aurant. cond. ana ℥ß Eryng. cond. conſerv. roſar. r. ana ℥j ſem. eruc. ʒij ambr. gryſ. gr. v. fol. ☉ri n. iiij ſyr. gran. cherm. & Cinam. ana q. ſ. f. Elect. ad ♀em languidam excitandam.

℞. Coagul. leporis ℥j ſatyr. cond. ℥ij ℴℴ majoran. gt. xII. ambr. gryſ. ʒj ſyr. cherm. q. ſ. f. Elect. cap. ad ʒj c. Eſſ. ſatyr. præcipue ad conceptionem promovendam, commode autem capiuntur illa medicamenta mox poſt fluxum menſium & quidem ante introitum lecti.

In Italia recipiunt radicem recentem ſatyrii, exprimunt ſuccum, & miſcent cum vitello ovorum capiuntque ſingulis horis matutinis.

[De Mumia Satyrii aphrodiſiaca conferri meretur Henr ab Heer obſ.9. p m. 133. ſatyrium inter analeptica medicamenta à nonnullis refertur.]

***

RADICES SALE BLANDO FIXO, TENUI VISCOSITATI INCLUSO CONSTANTES.

Scorzonera.
Eryngium.
Bardana.
Cichorium.

SCORZONERA.

[Scorzonera tum latifolia, tum anguſtifolia C.B.P. eſt radix oblonga, digiti craſſitie, foris pallida, intus alba, ſaporis ſubdulcis ac grati, odoris nullius.]

Alia ſcorzonera ad nos transfertur ex Hiſpania, cui noſtra viribus plane non convenit, caret enim virtute aromatica.

Quæ in noſtris officinis proſtat, gaudet ſale blando tenui viſcoſitati involuto, quo acria quævis & acida mulcet & temperat, unde in pthiſi, macie, ictero, febribus malignis, variolis & morbillis laborantibus tam medicamentum, quam alimentum eſt.

Datur in ſubſtantia cocta cum brodio carnis pro alimento, condita vero pro medicamento. Jungitur egregie C. C. ſalſæparillæ cichorio. Decoctis acrioribus additur tanquam temperans.

℞. Scorzon. Siſar. paſtinac. maj. ana coq. in brodio, exhibetur hecticis & tabe conſumtis, temperat enim ſanguinem acriorem.

℞. Scorzon. ℥ij Cichor. ℥j pomum integr. Citri ſimul decoque pro potu ordinario in febribus.

℞. Rad.

℞. Rad. ſcorzon. ℥iiij f. decoct. c. 🜄 quo capiat caſtor. gr. v. l. Eſſent. Caſtor. aut Tinctur. Myrrh. gr. xv. contra variolas & Morbillos.

℞. Scorzon. ℥ß hord. mund. ℥j coq. in ſ. q. 🜄 fontan. add. Syr. roſar. vel violar. q. ſ. ut & -∩- 🜍is per camp. q. ſ. ad gratiam f. julapium refrigerans in Morbis quibuſvis calidis.

ERYNGIUM.

[Eſt Eryngii vulgaris C.B.P. radix longa, pollicem craſſa, foris nigricans, intus alba, in medio nervo ſolidiore flaveſcente repleta, ſaporis ſubacris dulcis, odoris nullius. Refertur ad plantas ſeminibus ſolidis & floribus pluribus in capitulum digeſtis.]

Eryngii vis potior in cortice conſiſtit, ob levem amaritiem potentior eſt ſcorzonerâ, unde incidendo & fluxile reddendo ſerum, id per Urinam elicit, cæterum ſcorzoneræ viribus reſpondet.

Non facile poteſt in pulverem teri, niſi ſiccetur, ſed tum abit omnis vis, hinc radix in decocto & condita datur. Veneris quoque incentivum blandum eſt, & ad gonorrhæam ſimplicem commendatur à Welſchio [ſyllog. obſerv. Marcell. Cumani obſ. 44. p.m. 64.]

℞. Rad. Eryng. ℥ij bacc. junip. ℥j f. decoct. ex quo ℞. ⊖ geniſt. gr. xv. contra difficultatem Urinæ cum ardore.

℞. Rad. Eryng. Scorzon. ana ℥ij Cichor. bardan. ana ℥j ſalſæparill. ℥ß ſem. fœnicul. ℥j f. decoct. antiphthiſicum, addi poteſt colaturæ -∩- ⊖✳ci vel ⊖ volat. ☍ſ. Sylvii ſecundum judicium Medici M. c. Syr. d. Symph. Fern. v. c. ad ℔j decocti ℥j ſyrupi.

BARDANA.

Eſt Lappæ majoris, Arcii Dioſc. C.B.P. radix oblonga, craſſa, foris nigricans intus alba, ſaporis ſubdulcis, acris & amaricantis, odoris fere nullius, pertinet ad plantas capitatas pappeſcentes, floribus fiſtuloſis.]

Quod potentior ſit Eryngio judicatur ex eo, quia majorem habet amaritiem, præſtat ſi radix recens uſurpatur. Cortex radicis acrimonia majori gaudet, quam interior ſubſtantia radicis.

Primo anno enata digiti craſſitie, eſt effodienda, altero enim anno cauleſcit radix, ſuntque vires in floribus & caule.

Bardanæ radix Eryngio potentior, hinc non tantum Urinam movet, eamque ſæpe turbidam reddit, ſed etiam ſudorem movet, & ſanguinem congrumatum reſolvit, propterea in Pleuritide, phthiſi, atrophia, arthritide pro ſpecifica habetur.

Datur in ſubſtantia ejus raſpatura recentior à ʒj ad ij, in pulvere vix habet uſum. Datur & in decoctis. D. Herm. ſæpe ægrotis ab alto delapſis præſcribit. ∇ deſtillata vel potius decoctum eſt vehiculum conveniens in qua raſura radicis bardanæ contra pleuritidem exhibenda eſt.

℞. Rad. bardan. ʒij herb. Chamædr. Chamæp. ana Mj. coq. in lacte, exhibe contra arthritidem.

℞. Raſ. bardan. ʒij cape ex decocto proprio bardanæ in pleuritide.

℞. Raſ. bardan. ℈j. dent. apr. mand. luc. piſc. ana ℈ß. fl. 🜍is. gr. v. ♁ diaph. gr. x. vel xv. M. f. pulv. ſumendus ex decoct. bardanæ bis vel ter de die in Pleuritide.

CICHO-

CICHORIUM.

Eſt Cichorii ſylveſtris ſeu officinarum C.B.P. radix oblonga, circuloſa, dum recens lacte turgida, ſaporis ſubacris ſubadſtringentis, amaricantis. Radix Cichorii ſativi minus medicamentoſa.

[Jungitur plantis ſeminibus ſolidis ſeu non pappeſcentibus & lacteſcentibus floribus planipetalis.]

Conſtat ⊖e diuretico admodum blando, & ferme tanquam polychreſtum tum medicamenti tum alimenti loco variis in Morbis laudatur, præcipue ubi reſolvendum, temperandum, abſtergendum eſt. Hinc in inflammatione faucium, ictero, obſtructionibus vaſorum lacteorum, in ſanguine acri & acido temperando, in nimia ſiti, ingrato ſapore oris, feliciter adhibetur, additur quoque medicamentis purgantibus juſto acrioribus aliiſve, ut & antiſcorbuticis & antimelancholicis.

Datur in ſubſtantia vel ſaccharo condita, vel decocta in aqua vel brodiis carnium, junctis appropriatis, datur & ſuccus cum ▽ deſtillata. Taraxaco tanquam genuino analogo jungitur.

℞. Rad. Cichor. Mj. paſſul. min. ℥ß vel j. coq. in ▽ Colatur. infund. ferment. panis petiis except. q. ſ. repone donec fermentet, fit potus egregius pro ſiti fallenda.

℞. Rad. Cichor. ℥ij tarax. chelid. maj. fragar. ana Mj. coq. in vino vel ▽ in ℔j Colatur. infund. fol. ſenn. ℨvij f. apozema alterans & temperans univerſaliter ſanguinem.

RADICES

⁂

RADICES SALE BLANDO DIURETICO CONSTANTES.

Aſparagus.
Ruſcus.
Ononis.
Rubia tinctorum.

ASPARAGUS.

[Sunt Aſparagi ſativæ C.B.P. Radices ſpongioſæ, longæ rotundæ ex uno capite in plures fibras ſeſe diffundentes, ſaporis ſubdulcis ſubacris, odoris nullius. Eſt planta baccifera foliis capillaceis tenuiſſimis, alternis. Floribus ſubrotundis parvis, propendentibus, radice fuſi formi.]

Blandum ſuum ſal diureticum in blanda mucagine fovet & contra icterum, alioſque Morbos chronicos cum appropriatis exhibetur.

Vis conſiſtit in cortice radicis recentis, ut & julis ſeu novellis germinibus quæ ratione ſpiritus vegetantis quo turgent, ♀em ſtimulare creduntur.

Ob ſui mucaginem in quam reſolvuntur germina nocent iis in quibus temperamentum eſt acidiuſculum, ſunt & viſcoſiores.

℞. Rad. aſparagi. ℥j gramin. ℥ij bacc. junip. ℥iiij f. decoct. diureticum ex vino & aqua.

Ex decocto aſparagi datur etiam ☿ vivus vel dulcis contra lumbricos, infantibus & adultioribus, ſed decoctum & ▽ graminis præſtat in hoc caſu.

RUSCUS

RUSCUS.

[Eſt Ruſci C. B. P. radix craſſa, obliqua, implexa, fibras crebras longas albas demittens, ſimiles fibris aſparagi, annumeratur plantis bacciferis.]

Ruſcus viſcoſitate qua aſparagus turget, caret, & propterea tutum & blandum eſt diureticum, imo potentius quam aſparagus, ut in ſpecie antihydropicum audiat. Datur in decoctis.

℞. Rad. ruſc. ℥iiij coq. in 🜄 communi, decocto adhuc calenti infunde fol. ſenn. ℥j. f. apozema. Eſt medicamentum Riverii contra arthritidem.

℞. Rad. ruſc. ℥iiij ir. florent. ʒij f. decoct. ex 🜄, ſi radix ir. flor. non ſufficit, poſſunt addi fol. ſennæ, exhiberi poteſt in hydrope, ubi parcius mittitur Urina.

ONONIS.

[Sunt Anonidis ſpinoſæ flore purpureo C. B. P. radices albidæ, fractu contumaces, lignoſæ, ſaporis & odoris nullius. Pertinet ad leguminoſas papilionaceas, ſiliquis unicapſularibus, ſingulis ſequentibus ſingulos flores tetrapetaloides.]

Ononis ſal ſuum diureticum ligneæ quaſique terreæ ſubſtantiæ incluſum habet, hinc præter virtutem diureticam habet quoque vim abſorbentem & abſumentem ſarcocelen & ramices carnoſas.

Datur cortex, in quo vis major quam in interiori ſubſtantia, ex decoctis & infuſis. Stimulatur ſale proprio vel geniſtæ.

℞. Rad. ononid. ℥iiij herb. ononid. Mij f. infuſ. ex vin. Malvat. deſtilletur, deſtillatum novis radicibus & foliis affunditur, & ponitur in digeſtione, & cochlea-

tim cum ⊖e proprio geniſtæ à gr. vij. ad xv. contra calculum exhibetur.

℞. Cort. rad. ononid. ʒj exhibe ex decocto proprio per integrum menſem ad curandam ſarcocelen.

RUBIA.

[Eſt rubia tinctorum ſativa C.B.P. radix oblonga, tenuis, ſucculenta, extus & intus rubicunda, ſaporis ex dulci amaricantis & adſtringentis, odoris nullius.

Refertur ad ſtellatas, ſeminibus poſt ſingulos flores nudis binis, foliis ad ſingula caulium genicula pluribus, radiatim decuſſatis.]

Si inculta naſcitur parum rubedinis habet, & ad uſum tinctorum minus apta, ſed ad uſum Medicum optima. Rubia partium tenuiorum quam ononis, & proinde virtutis potentioris, viſcerum ſimul obſtructionem reſerat, & ſanguinem congrumatum potenter diſſolvit, quam ob cauſam vulneraria cenſetur. Datur ad ʒj in pulv. in dec. ad ℥ij.

℞. Rad. rub. ocul. ♋. ana ʒj. ſperm. Cet ℈j. ⊖ vol. ſuccin. ℈ß. M. f. pulv. pro ij vel iij doſibus contra caſum ab alto, ubi metus ſanguinis congrumati, ſimul gr. j. vel alterum moſchi detineatur in ore.

℞. Rad. rub ℥iiij Cin. ſtip. fabar. Mß coq. in ▽ f. apozem. diureticum, in difficultate Urinæ ſeu iſchuria à ſanguine congrumato.

* * *

RADICES AMARÆ SALE FIXO CONSTANTES.

Conveniunt virtute cum ⊖ibus volatilibus ☍ſis, verum rigidiora ſunt & potentiora, ut in morbis chronicis

nicis pertinaciſſimis, & auſtera acida putrida validius infringunt, ac per diureſin & diaphoreſin expellunt.

GENTIANA.

[Eſt gentiana major lutea C. B. P. Radix longa, craſſa, foris fuſca, intus ex flavo rufeſcens, ſaporis intenſe amari, odoris nullius. Pertinet ad plantas bicapſulares flore monopetalo uniformi.]

Gentiana ob amaritiem intenſiorem & ⊖ fixum rigidius & potentius in ſpecie confert in febribus intermittentibus, & ab his orta ventriculi debilitate, expellit quoque lumbricos, quibus amara adverſa ſunt, & ratione virtutis alcalinæ, contrariantur acidis putridiſve in quibus vermes nidulantur.

Sub ſpecie turundæ fiſtulis intruditur, quia laxior radicis ſubſtantia à ſuſceptis ex fiſtula humoribus intumeſcens eam ſenſim dilatat, & quia ⊖ acre callum ulceris fiſtuloſi exedit ſeu abſumit: eandem ob cauſam pulvis ejus fonticulis ubi fluere deſinunt inſpergitur, vel ſub forma globuli inditur.

In theriacam recipitur: ex ſolo melle & hac radice fac theriacam: in hydrophobia laudatur ſpecifice; ob nimiam ſui diaphoreticam & diureticam exſiccantem vim obeſis & pinguioribus convenientior eſt quam emaciatis.

℞. Rad. gentian. ʒß ⊖ abſynth. gr. xv. in robuſt. ad ℈j commodiſſime in pilulas redigitur cum extract. inſpiſſato abſynthii q. ſ. f. pil. ☉ fol. involvend. Conveniunt in omnibus febribus intermittentibus ſeu tertianis ſeu quartanis.

℞. Rad. gentian. ſerpent. virg. Cort. Chin. Chin. ana ℈ß. ⊖ prunell. ℈ß. vel gr. xv. M. f. pulv. vel prædicto modo poteſt redigi in pilulas.

Cum ſaccharo ♄ni poteſt exhiberi in ▽ graminis contra vermes.

℞. Rad. gentian. ℥iij ſerpent. virgin. ℥ß. Cort. Ch. Chin. ℥iij flor cent. cent. min. p. j. pip. long. ℥ß. M. f. vin. medicatum. cui in alvi adſtrictione adde rad. Jalapp. (folior. ſennæ) ℥ß. helleb. n. ℥iij paſſul. exacin. ℥j f. infuſum vel vinum medicatum.

⁂

RADICES SALE FIXO CUM PAUCO SALE VOLATILI CONSTANTES INCLUSO RESINÆ.

Peucedanum.
Fraxinella ſeu Dictamnus albus.
Ariſtolochia.
Filix utraque.
Oſmunda.

Huic ⊖i fixi conjuncta eſt reſina, quæ ad ſtrobilos pini & abietis accedit. Unde quoque judicatur participare de blanda quadam volatili aciditate, qua ſtrobili pini gaudere videntur, idcirco convenit in Uteri obſtructione, in ſedandis doloribus, incidendo pulmonum muco.

PEUCEDANUM.

[Eſt Peucedani Germanici C. B. P. radix craſſa, longa, ſuperius coma prædita, extus nigreſcens, intus albeſcens, ſaporis pinguis acriuſculi, odoris ſulphurei.]

Datur ut ſuperiores radices,

℞. Rad.

℞. Rad. Peuced. ſtrobil. pin. & abiet. ana ℥ij herb. chamæp. Mß. f. decoct. ex ▽ vel vino, ad quoſcunque dolores, & obſtructiones Uteri tollendas: remedium fuit Plateri.

FRAXINELLA.

[Eſt dictamni albi vulgo ſive fraxinellæ C. B. P. radix, ſeu potius radicis cortex craſſiuſculus, albicans, cannæ in modum convolutus, ſaporis amari.]

Fraxinella plus habet ⊖is volatilis quam Peucedanum, quod arguit bituminoſus & aromaticus odor, propterea etiam cortex, in quo vis concentrata eſt ad partum facilitandum, excludenda lochia, maxime laudatur, ob intenſam ſui amaritiem contra lumbricos exhibetur. Datur ut cæteræ radices.

℞. Rad. fraxinell. ʒij N. M. ℈ß. M. f. pulv. exhibetur ex vino vel ▽ puleg. ad facilitandum partum, loco ſtimuli add. Caſtor. gr. v. borrac. gr: v.

℞. Rad. fraxinell. ʒiij ariſtoloch. ℥j caſtor. ʒj f. infuſum ad paſſionem hyſtericam tollendam, it. ad remoras menſium & fœtum expellendum.

ARISTOLOCHIA.

Ariſtolochia rotunda cæteris ariſtolochiis præſtantior, ſal fixum paucumque volatile in ſubſtantia pinguiori fovet, propterea in tuſſi, aſthmate, facilitando partu, menſibus pellendis, ſcabie, incidit, aperit & anodyna eſt. Recipitur quoque inter vulneraria, & in clyſteres ad Morbos ſoporoſos, it. in cardialgia cum ovo ſorbili data ſpecifica audit.

Ariſtolochia longa ⊖ia ſua in ſubſtantia laxa & nonnihil ſpongioſa gerit, unde ſimul imbibit, abſorbet

abſtergit, hinc veluti gentiana fiſtulis dilatandis intruditur ſub ſpecie turundæ.

Ariſtolochia tenuis remiſſiorem habet amaritiem, aſt ſi ſit recens intenſiorem acredinem volatilem & penetrantem, quo nomine cæteris ariſtolochiis additur tanquam ſtimulans.

Ariſtolochia cava ⊖ fovet in ſubſtantia farinacea ferme in mucilaginem fatiſcente, & à multis præfertur ariſtolochiæ rotundæ, præcipue quoad vim anodynam & mulcentem, ſi nempe ſit recens.

Ariſtolochia fabacea remiſſior eſt & debilior quam cava, ob ſui ſubſtantiam farinaceam majorem habet vim exſiccantem & abſorbentem quam cæteræ Ariſtolochiæ, unde illam ſolent eligere Chirurgi in denudatis oſſibus.

℞. Ariſtol. ir. flor. ana ℥j f. pulv. tenuiſſimus oſſibus denudatis inſpergendus, ubi vero jam nigreſcunt oſſa, add. Euphorb. ℈j. in principio gangrænæ frequenter uſurpatur.

℞. Ariſt. rot. ʒj. cape ex vino enulato contra ſcabiem cum aſthmate & doloribus.

℞. Ariſt. rot. ℈ij Myrrh. gr. xv. borrac ℈ß. Caſtor. gr. vij. Croc. orient. gr. v. f. pulv. in partu difficili: ſumi poteſt ex vino.

℞. Rad. ariſt. rot. ʒj. ſumatur ex ovo ſorbili calido contra cardialgiam, quod mire deprædicatur.

℞. Ariſt. cav. ir. flor. helen. ana ℈j f. pulv. capiendus ex oxym. ſcyll. ▽ puleg. contra aſthma ſcorbuticum cum doloribus.

℞. Rad. ariſt. cav. gentian. ana ℈j ☿ dulc. rit. par. gr. xv. f. pulv. pro ij vel iij doſ. ſumi poteſt ex ▽ graminis contra lumbricos.

FILIX.

FILIX.

[Filix mas ſeu non ramoſa dentata C.B.P. & femina ſeu ramoſa major pinnulis obtuſis non dentatis C.B.P. eſt radix craſſa, ramoſa, fibroſa, foris nigricans intus pallida, multis appendicibus involuta & implexa, ſaporis primum ſubdulcis inde amariuſculi ſubadſtringentis, odoris nullius.

Jungitur plantis epiphylloſpermis, floribus muſcoſis & ſeminibus pulverulentis averſa foliorum parte adnatis, exiguis veſiculis per maturitatem in binas partes diſſilientibus incluſis, foliis diviſis.]

Filix utraque ⊖ ſuum fixum particulis terreis & viſcidis incluſum habet, laudatur tanquam aperiens in affectu hypochondriaco aliiſque morbis chronicis; In pellendo calculo & nephritide ut & in vermibus expellendis ſingularis habetur. Decoctum ejus ſuper vermes effuſum eos necat, quia oleoſum eſt & vermibus aërem non concedit. Expreſſa mucago membris ambuſtis mire medetur.

Contra lumbricos datur in ſubſtantia, ſtimulatur ☿dulci ſem. cinæ ſeu ſanton. it. in morbis chronicis datur in decocto.

Vis plurima conſiſtit in julis ſeu novellis germinibus.

℞. Rad. filic. oſmund. polypod. ana ℥ij herb. capill. Mj paſſul. corinth. ℥ß. ⊖ ♃ri ℥ß. coq. ex ▽ communi, decocto calenti add. Epithym. Mj. caryoph. arom. ʒß. f. apozema contra morbos chronicos, add. ſyr. capill. ♀is. q. ſ.

℞. Rad. Filic. femin. ʒj ☿ dulc. gr. x. f. pulv. cap. ex ▽ gramin, pro ij doſ excipe melle & fac. bolum.

OSMUN-

OSMUNDA.

Filix ramoſa non dentata florida C.B.P. Filici analoga eſt, aſt temperatior & blandior, fovens in mucagine nonnihil ⊖is volatilis, quare ſpecifice commendatur in rachitide.

Mucago ejus ad curandam herniam in pueris extra applicata approbatiſſimum remedium: Vis omnis conſiſtit in julis & in radicis medulla.

℞. Jul. Oſmund. coq. c. lacte, cape ſingulis diebus contra rachitidem.

℞. Oſmund. medull. ℥iij Cort. Winter. ʒiij herb. capill. Mj. coq. ex ▽et vin. f. decoct. contra rachitidem.

RADICES PURGANTES.

Conſtantes particulari textura & figura, poris membranarum tum ventriculi & inteſtinorum, tum particularum ſanguinem conſtituentium è directo contraria, vehentes ſimul auram ſubtilem ſanguini impermiſcibilem & hoſtilem.

* * *

SALE VOLATILI GAUDENTES.

Aſarum.
Cyclamen.
Hermodactyli.
Rhabarbarum.
Rhaponticum.

ASARUM.

[Eſt Aſari C.B.P. radix tenuis, fibroſa, ex fuſco rufeſcens, ſaporis acris nauſeoſi, cum aliquali adſtrictione,

ne odoris ad Nardum accedentis, unde & Nardus ſylveſtris appellatur.]

Aſarum gaudet Θe purgante acri, fixo potius quam volatili, & ſtyptica terra, quod confirmat odor, ſapor, modus exhibendi & præparandi, in receſſu habet quandam vim adſtrictoriam.

Datur in ſubſtantia à ʒß ad ʒj & ℈iiij, movet vomitum & ſedes. Vomitus autem volatilitatis, ſedes fixitatis argumentum ſunt. Datur in infuſo & decocto à ʒij ad ʒiij in extracto à ℈j ad gr. xxv. in infuſo rarius vomitum, ſedes vero frequentius movet, fortiori igne coctum vel toſtum vim purgativam amittit, ac diureticum & Uterinum fit.

Melancholicis ex aſaro peſſima accidunt ſymptomata, ſeroſis & bilioſis plurimum contuliſſe experientia probat: hinc in hydrope, hydrocephalo, ictero, febribus intermittentibus conducit. Riverius & Timæus extollunt in doloribus iſchiaticis, coxæ, alii vero in aſthmate tanquam ſpecificum remedium. Ob odoris nardini graveolentiam virtutis uterinæ indicem, Menſes, fœtum, lochia pellit & promovet. Pro ſcopo purgante datur ʒj, pro Utero movendo in minori quantitate gr.vij. ad ℈ß.

Toti huic radici non eſt confidendum, in quibuſdam officinis planta extat inſipida, plures non purgavit radix etiam in ſylveſtribus proveniens, contra in quibuſdam adeo vehementer, ut ultra ℈j non liceat aſſumere.

Conjungitur, ſtimulatur, corrigitur variis medicamentis appropriatis, uterinis & thoracicis additur pro ſtimulo.

℞. Rad. galang. ℈ß. aſari ʒß. Θ ☐ri gr. v. l. vj. M. f. pulv. capiendus ex oxymell. ℥j.

H ℞. Rad.

℞. Rad. aſari. ʒij gentian. ℥ß. centaur. min. pug. j. coq. leniter cum Cereviſ. ex hoc decocto cape ⊖ abſynth. ℈j contra febres intermittentes, præprimis contra quartanam.

℞. Rad. aſari ʒj calam. arom. ℥ß. curcum. ʒj. caryoph. ℈ß. f. infuſum cum vino antiictericum & hydropicum.

℞. Caſtor. ℈ß borrac. Ven. gr. xv. rad. aſari. gr. v. M. f. pulv. excipiendus conſerva florum ſtœchad. vel quocunque ſyrupo ſpecifico, f. bolus in Obſtructionibus Menſium.

CYCLAMEN.

Eſt cylaminis orbiculato folio inferne purpuraſcente C.B.P. radix tuberoſa, craſſa, orbiculata, carnoſa, foris nigra, intus albicans, ſaporis acris pungitivi exurentis, ingrati, odoris nullius. Conſtat particulis ⊖inis volatilibus, quæ avolant cum ſeneſcit, cum terreſtribus, quibus includuntur catharcticis.

Planta eſt tricapſularis, pentapetala, (petalis ad ungues junctis) ſeminibus anguloſis in capſulis rotundis.

Radix recens eſt valde acris, ubi vero exſiccata, nulla acredo percipitur. Purgat cum inflammatione Ventriculi, faucium & ani, ubi nulla purgantia in hydrope effectum dedere uſurpatur.

Particulis purgantibus vehementioribus gaudet quam aſarum, & proinde robuſtioribus datur pro purgando ſero & bile. Ob ſui penetrantiam ſuccus abdomini hydropicorum applicatus aquas educit, unde unguentum de arthanita ingreditur, quod ſequentibus ſtimulatur 1. ☍ colocynth. per infuſionem facto.

℞. Un-

℞. Unguent. d. arthanit. ℥j 🜍 colocynth. q. v. inunge totam abdominis regionem. 2. 🜍 ex granis Tilli, quæ ſuppetunt 🜍 acerrimum pauciſſimis notum, penes Medicos Indos ſecretiſſimum. Additur quoque Clyſteribus ad excitandos ſoporoſos. Quidam ad ſuppoſitoria recipiunt cum aloë, ad promovendum fluxum hæmorrhoidum.

Licet cum violentia ſumma expurget humores, tamen tardioris eſt operationis, propter particulas terreſtres ut propterea raro in uſum trahatur. Stimulatur aſaro, diagrydio, ☿ro. Plebeji robuſti capiunt ejus ſubſtantiam ad ʒj pro decocto ℥ß. Sed tutior modus exhibendi eſt extracto facto cum 🜋 vini vel aceto à ℈j ad gr. xxv. xxx.

℞. Extr. cyclam. ℈j 🜔 ☿ri gr. vij. vel ℈ß. diagryd. gr. v. oxymell. q. ſ. f. bolus. in hydrope, ubi alvus purganda.

℞. Extr. cycl. ʒß. oxym. ℥ij corrigitur 🜄 cinam. ℥ß. f. hauſtus in hydrope.

HERMODACTYLI.

Eſt Colchici radice ſiccata alba, C.B.P. radix rotunda, tuberoſa, avellanæ magnitudine, compreſſa, albida, ſaporis dum recens acerrimi, dum annoſa ſubdulcis farinacei paulum viſcoſi vel lenti, odoris nullius.

Radix non niſi ſiccata apud nos habetur.

Hermodactylus recens 🜔em volatilem purgantem in farinacea mollecula ſubſtantia fovens, pituitam & ſerum per vomitum & alvum purgat, ſiccatus vero vel toſtus, & proinde 🜔ibus volatilibus orbatus, propter annoſitatem ſæpe omni virtute catharctica orbatus eſt.

In Ægypto ob blandam viſcidam farinaceam ſubſtantiam cedit præſertim feminis emaciatis in nutrimentum, veluti ari radix & colocaſia [conf. Proſp. Alpin. de plant. exotic. l.2. c.18.]

In oriente datur recens, cum Zinzibere ſumunt hermodact. ʒß Zinzib. ℈ß. Zinziber additur propter excrementitiam ejus viſcoſitatem tormina cauſſantem. Omnes Medici Indorum, ſolo Zinzibere purgantia ſua corrigunt.

Siccatus qualis in officinis proſtat, non niſi cum cæteris purgantibus fortioribus præſcribitur, contra arthritidem & podagram.

Hermodactylus commendatur in gutta ſerena tanquam ſpecificum, ſeq. modo.

℞. Hermod. ʒß. exhibe ex 🜄 fœnicul. ad gutt. ſerenam, quæ fit à nervo optico obſtructo vel compreſſo à materia viſcida.

℞. Hermod. ℈j. turbith. gialap. ana gr. viij Diagr. gr. iij Zinz. ℈ß. M. f. pulv. ad podagram, ubi alvus eſt adſtrictiſſima.

RHABARBARUM.

Eſt Lapathi maximi Sinenſis radix prægrandis, oblonga, tuberoſa, circularibus zonis rugoſa, tuberibuſque inæqualis, communiter ponderoſa, foris fuſca intus crocea, nonnihil fungoſa, ſaporis amaricantis & lenta viſcedine ſubadſtringentis, nauſeoſi odoris pontici debilis.

Conſtat 🜔e purgante volatili, pauca reſina, & terra. hinc ob 🜔 volatile pinguedine quaſi temperatum blande purgat bilem & pituitam, & ſpecificum eſt in ictero, & affectionibus à bile oriundis.

Vim

Vim purgantem in volatilitate conſiſtere exinde patet. 1. quia extractum cum ▽ factum purgat. 2. quod ſiccatum ætate omnem vim purgantem amittit & acredinem, 3. quod aqua ejus deſtillata eſt nauſeoſa, acris & purgativa, quod jam Borrichius in act. Hafn. obſervavit. 4. ejus odore ſolo quidam ad ſedes moventur 5. per coctionem amittit purgantem virtutem.

Subducit quoque copioſe Urinam, eamque bene coloratam & pinguem reddit. Ob reſinam blandam anodynum eſt & balſamicum in vulneribus, ſuas operationes cum vi anodyna abſolvit, ob terram auſteram relinquit in receſſu vim adſtringentem.

Tota ſui ſubſtantia ſingulare antidyſentericum infringens, purgans & mitigans acidum, relinquens in receſſu vim anodynam & adſtrictoriam.

Conveniens eſt Medicina Melancholicis, quod acidam ventriculi cruditatem tollit, nam quando acidum affunditur rhabarbaro effervescit, ut & delapſis ab alto, quod ſanguinem congrumatum diſſolvit.

[Rhabarbari elogium & de ſingulari ejus virtute contra hydropem vid. ap. Scholz. Epiſt. med. xv. p. m. 18.]

Conducit in gonorrhæa, urinæ ardorem temperando, malignitatem expellendo, fluxumque ſiſtendo. Toſtum & volatilitate orbatum pulchre abſorbet & adſtringit, hinc in gonorrhæa, hæmorrhagiis, aliiſve fluxibus ſiſtendis feliciter adhibetur.

Vertiginoſis aliquando nocet, [conf. Joh. Mauric. Hoff. C.N.C. Dec.III. an. 2. obſ. 208. de vertigine à rhabarbaro] quia ejus fumus & ponticus odor cerebrum petit & turbat.

Crudum datur ad ʒj. ʒiß. in infuſo ad ℥ß. Ob debilem purgandi vim jungitur & ſtimulatur variis. Corrigitur ejus acredo & amarities refrigerantibus aqueis, v.c. 🜄 Cichor. Endiv. Plantag. ſero lactis, ſucco violarum, roſarum. Toſtum ſeu fortiori 🜂e coctum & rurſus ſicccatum datur ad ʒj.

℞. Rhabarb. ʒj 🜔 ♀r. gr. x. vel xii. M. f. pulv. in ictero ex vino calido ſumendus.

℞. Rhabarb. ʒj Elect. è ſucc. roſar. ʒiij 🜔 abſynth. Crem. ♀ri ana ℈ß. ſyr. d. cich. c. rhab. q. ſ. M. f. bolus in omnibus affectibus bilioſis exhibendus.

℞. Rhab. ʒß. 🜔 abſynth. ℈ß. N. M. ℈ß. caryoph. g. ij l. iij M. f. pulv. exhib. ex 🜄 Cinam. pro purgante in febribus cum diarrhæa.

℞. Rhab. ʒij fl. roſ. rubr. p. ij infund. in ſeri lactis ſ. q. pone in digeſt. per aliquot horas ℞ Colatur ℥iij add. ſyr. roſ. ſol. ℥ß. M. f. pot. laxans in dyſenteria benigna.

℞. Rhab. ʒß Laud. op. g. j. additur propter tormina, Conſerv. roſ. r. ʒj ſyr. roſar. q. ſ. f. bolus in Dyſenteria conveniens.

℞. Rhab. ʒß. extr. tormentill. inſpiſſ. gr. xv. terr. ſigill. l. C.C. præp. l. unicorn. foſſil. ℈ß. theriac. ℈j ſyr. roſ. q. ſ. f. bolus in Dyſenteria maligna.

℞. Rhab. toſt. ʒj ℴℴ Nuciſt. expr. ʒß. Corall. r. præpar. ℈j Laud. op. gr. j M. f. pulv. excipiendus ſi placet conſerv. roſar. r. ſ. nymph. f. bolus in quocunque fluxu ſiſtendo.

℞. Rhab. toſt. ʒj. N. M. toſt. ſacch. ♄ni ana ℈j oſſ. ſep. præp. ʒß. reſin. thereb. q. ſ. f. pil. in gonorrhæa ſimplici & pollutione nocturna ex debilitate, add. terr. aluminoſ. dulc. ℈j, ad malignitatem expellendam adde

de ☿ virid. quantum videtur, nempe pro qualibet doſi gr. iij ad v. in principio, in progreſſu vero ſingulis diebus gr. j. ſub finem gonorrhæa ſi fluxus non vult ſiſti add. borrac. calcinat. ℈j & cum balſ. peruv. f. pil.

℞. Rhab. reſid. ex tinct. ʒß. N. M. toſt. ℈ß. Laud. op. gr. j. conſ. roſ. ⊕lat. ʒj. M. f. bol. adſtringens.

℞. Rhab. ʒß thereb. ʒj ſacch. ♄ ℈ß ☿ virid. gr. j. vel ubi major malignitas adeſt gr. iij cum vitello ovi & ▽ appropriata fiat hauſtus contra gonorrhæam.

RHAPONTICUM.

Eſt Rhaponticum folio Lapathi majoris glabro, Rha & Rheum Dioſcoridis C.B.P. Eſt Lapathi Thracici maximi radix craſſa, oblonga, ramoſa, brachiata, foris fuſca, intus crocea laxa & rara, tranſverſim ſecta ſpiras oſtendens, ſaporis adſtringentis cum levi quadam acredine & lentore, odoris pontici debilis.

Rhabarbarum & Rhaponticum conſimiles flores & ſemina habent, & facie plantarum nihil differunt, ſed tantum ratione radicis, idcirco præſumitur unam eſſe plantam, diverſitas radicis oriri poteſt à ſolo.

Conſtat particulis ⊖inis purgantibus ſed debilibus valde in Thracico, potentioribus in Chinenſi, incluſæ mucilaginoſis balſamicis & terreſtribus.

Rhabarbaro mitius in purgando, potentius vero in adſtringendo : laudatur præterea ob blandum ⊖ incidens in orthopnœa.

Datur in ſubſtantia ad ʒj in decocto & infuſo ad ℥j.

℞. Rhapont. Ocul. ♋ ana ℈j ſperm. Cet. gr. vij ⊖ vol. C. C. ℈j M. f. pulv. ad ſanguinem congrumatum reſol-

reſolvendum, ubi multo præſtantius quam rhabarbarum propter levem dulcedinem.

℞. Rhapont. ℥iß. helen. ir. florent. ana ℥j. imperat. ℥ß. glycyrrh. ʒiij fol. hyſſop. marrub. ana Mj. ſem. fœnic. aniſ. ana ʒij f. decoct. ex ▽ colatur. add. oxym. ſcyllitic. q. ſ. ut & ♌ aſthmat. q. ſ. f. apozema in gravi & difficili reſpiratione.

*
* *

PURGANTIA SALE FIXO RESINÆ INCLUSO CONSTANTIA.

Gialappa.
Mechoacanna.
Turbith.

GIALAPPA.

Eſt convolvuli Mexicani Jalappa dicti radix oblonga, craſſa, mechoacanna nigricantior [unde & Mechoacanna nigra dicitur] & circuloſior, ſapore acuto incidente, muſcoſo, [nauſeoſo] odore nullo, ſciſſa in taleolas ad nos adfertur.

Planta eſt monopetala ſeminibus anguloſis in capitulis ſubrotundis. ut & Mechoacanna.

Gialappa quamvis univerſaliter quoſvis humores purgat tuto & citra ullam violentiam, ut & pueris exhiberi poſſit, tamen ſerum & aquas ſpecifice expurgat. Ejus vis purgans conſiſtit in reſina, cui ⊖ia inſeparabili nexu ſociata, eaque quaſi temperata. hinc reſina & extractum frequenter exhibetur.

Subſtantia datur à ℈j ad ʒß. reſina à gr. vij. ad ℈ß. extractum à gr. xv. ad ℈j.

Tinctu-

Tinctura Galappæ proſtat in officinis Batavis, ſed quia quantitatem ♎ vini affuſam neſcimus non tuto præſcribitur.

Galappa in ſubſtantia multo præſtantior eſt in hydrope quam reſina, & extractum [Borellus Cent. 2. obſ. 23. appellat arcanum in hydrope] alias extractum mitius ipſa reſina, quia ⊖ia jam in eo ſunt diſſoluta.

Semper cum aliis jungitur purgantibus, [per ſe ſola tuto poteſt exhiberi] ſalibus, morbo appropriatis, corrigentibus, adjuvantibus & ſtimulantibus, ut ⊖e ♀ri.

℞. Galap. gr. xv. Crem. ♀ri. gr. x. diagryd. gr. v. ☍ fœnic. gt. ij. M. f. pulv. purgans catholicus.

℞. Galap. gr. xv. diagryd. gr. v. ⊕tr. depur. gr. vij. ☍ cinam. gt. j. M. f. p. in alvi adſtrictione.

℞. Galap. gr. xv. ☿ dulc. ℈ß diagryd. vel troch. alhand. gr. iij. ☍ junip. gt. ij. M. f. p. purgans in hydrope.

℞. Galap. ʒj diagryd. ʒß. Caryoph. arom. Zinz. ana ℈ß. gumm. ammoniac. q. ſ. in aceto vel ▽ Cinam. ſol. f. pil. uterinæ ad ℈j propinandæ.

℞. Reſin. galap. gr. vij. diagryd. gr. iij. ☿ dulc. gr. x. ♀r. ⊕lat. gr. vij. M. f. pulv. vel cum conſerv. roſ. pallid. f. bolus.

℞. Extr. Galap. ℈j troch. alhand. gr. iij. ☍ cinam. gt. j. M. f. pil. deaurentur, ad uſum Cephalicum, vel cum conſerva florum tunicæ f. bolus.

MECHOACANNA.

Eſt Bryonia ſeu Mechoacanna alba C. B. P. Eſt convolvuli Mexicani radix craſſa, ſubſtantiæ laxioris quam Galappæ, [alba unde & Rhabarbarnm album

 dicitur

dicitur, carnoſa, circuloſa, ſaporis farinacei, odoris nullius.] Effoditur in regione Mechuacan, ubi in taleolas ſciſſa ad nos transfertur.

Mechoacanna Galappæ analoga eſt, ſed quia reſinam habet pauciorem ideo remiſſior eſt jalappa.

[Inſigniter Mechocannam dilaudat Gabelchov. C.1. Cur.13. p.27. ſqq.]

Arthriticis in ſpecie conducit. Placentæ ſeu trochiſci ipſa radice præſtant, radix enim admodum fallax eſt, quia una pars ejus cum vehementia, altera vero nihil quicquam purgat. Ratio eſt, quia plurima pars reſinæ in qua eſt vis purgandi ad partem vulneratam in effoſſione tendit. Infantibus etiam commodiſſime dari poteſt, quia eſt inſipida & inodora, contra lactis grumeſcentiam datur.

Datur in ſubſtantia à ʒß ad ʒj extract. à gr.xv. ad ℈ij. reſina datur ut Jalappæ.

℞. Mechoac. ʒß diagryd. gr. v. ⊖ ♀ri gr. iij. ☍ caryoph. gt. j. M.f. pulv. in Melancholia exhibendus.

℞. Mechoac. ʒß amygd. dulc. ℥ß ſacch. in ▽ roſ. ſolut. q. ſ. f. l. a. panis Martius. in arthritide.

℞. Mechoac. ℈j ocul ♋. ʒß. ⊖ ♀ri gr. iij. M. f. pulv. contra grumeſcentiam lactis infantum, puerperis etiam exhibendus, addi poteſt ſperm. ceti gr. vij. vel ix.

TURBITH.

Eſt etiam ſpecies convolvuli indici, foliis altheæ ſimilibus. Radix oblonga, craſſa, reſinoſa, foris ferruginea, intus albida, ſaporis ſubacris, nauſeabundi reſinoſi. Tota planta lacte turget, concreſcente in reſinam, quia ſubſtantiam habet valde poroſam, reſina non

non æqualiter diſtribuitur, ita ut nihil reſinæ in quadam parte ſæpe ſubſit, ideoque non purgat.

Suſpectum eſt, quod ſæpius tormina excitat, quæ deducunt à reſina pertinaciore, ego vero à particulis irregularibus, in ima enim parte reſinoſitas tanta non eſt.

Turbith potentius in ſerum viſcidamve lympham agit quam Jalappa, & propterea in podagra & arthritide, ubi potentioribus purgantibus opus eſt, conducit. Vis conſiſtit in reſina ejuſque cortice, in quo reſina latet.

Cave ne temere exhibeas gravidis quia Uterum valde commovet.

Datur in ſubſtantia à ℈j. ad ℈ij reſina & extractum, quod omnium optimum, velut in Jalappa. infuſum à ʒij ad ℥ß.

℞. Turbith. vegetab. reſinoſ. ℈j. troch. alhand. ſeu diagr. gr. iij ⊖ ♀ri gr. v. ℅ caryoph. gt. j. f. pulv. pro educenda pituita.

℞. Extr. turbith. ℈j. diagr. gr. iij. caryoph. arom. gr. v. ⊖ ♀r. gr. v. balſ. Peruv. q. ſ. f. bolus.

℞. Rad. turbith. ʒß. glycyrrh. ℈ß ⊖ vol. ſuccin. ℈ß. helleb. alb. gr. v. f. pulv. ſternutatorius ad excitandos Apoplecticos,

In officinis proſtant ſpecies diaturbith. doſ. à ℈ß ad ʒß.

℞. Rec. ſpec. diaturb. ʒß Crem. ♀ri. ℈ij f. pulv. in hydrope.

*
* *

RADICES SALE FIXO ACRI CONSTANTES ET VEHEMENTISSIME PURGANTES.

Cucumis aſininus.
Eſula major & minor.
Helleborus niger.
albus.
Bryonia.

CUCUMIS ASININUS.

Eſt cucumis Sylveſtris aſininus dictus C.B.P. radix craſſa, oblonga, alba, ſaporis acris nauſeoſi, odoris dum recens viroſi. Planta eſt pomifera flore monopetalo marginibus tanquam partitis, fructu ovali, hiſpido hirſuto.

Serum & aquam per Vomitum ſedes & Urinas copioſiſſime, ſed tarde & cum violentia excernit, hinc rarius exhibetur hydropicis pro ſcopo purgandi per inferiora, ſed tutius datur extractum cum ♒ vini ♃iſato vel acet. deſt. factum à ℈ß ad gr. xv. & ℈j, additur ☿ dulc. ſtimulatur troch. alhand. ſimul corrigitur ☍ d. caryoph.

℞. Extract. cucum. aſinin. gr. xv. quod perperam elaterium vocatur, extr. troch. alhand. gr. iij ☍ d. caryoph. gt. j. Extr. abſynth. q. ſ. f. pil. contra hydropem.

℞. Rad. 5. aperient ana ℥j rad. cucum aſinin. ℥iij M. f. decoct. cui add. ſyr. 5. rad. aper. pro ſcopo diuretico poſſunt etiam ⊖ia addi.

[De

[De elaterii virtute & effectibus in hydrope conſ. Martin. Liſter. Exercitat. de hydrope p. 17. & hinc inde.]

ESULA.

Major & minor. Major eſt Tithymalus paluſtris fruticoſus C. B. P. radix oblonga, pollicem craſſa, foris fuſca, intus pallida, lacte acri exedente & nauſeoſo turgida, ſi recens: eſt Planta tricapſularis tricoccos, conceptaculis poſt ſingulos flores ſingulis.

Minor eſt Tithymalus foliis pini, forte Dioſcoridis Pityuſa C.B.P. radix oblonga, Eſula majori tenuior, foris fuſca, intus ex flavo albeſcens, lacte acri turgida.

Eſula minor virtute fortior majori, utraque ſupra & infra educit humores hydropicorum tanta cum vehementia & moleſtia, ut robuſtioribus tantum exhibenda ſit, in cachexia, febribus, hydrope, ſcabie. Jungitur alterantibus pro ſtimulo. Additur etiam Clyſteribus acribus ad excitandos Apoplecticos.

Vis omnis conſiſtit in cortice, qui à ℈ß ad ℈j in ſubſtantia datur. Feliciori cum ſucceſſu exhibetur extractum cum 🜅 vini vel aliis menſtruis ſubacidis factum. Extractum ex radice datur tantum ad gr. xv. & ℈j. Si in ſubſtantia datur corrigitur oleo aniſi. v. c. ℞. Cortic. eſul. min. ℈j Elæoſarch. aniſ. Citr. ana ℈ß. troch. alhand. gr. v. f. pulv. antihydropicus ex vino abſynthite.

℞. Extr. eſul. ℈ß extr. ligu. al. gr. vij. troch. alhand. gr. v. 🝆 caryoph. gt. j. conſerv. flor. perſic. ℈j. l. q. ſ. f. bolus in hydrope.

Quidam Medicus exhibuit infuſum in hydrope feliciter, quod tamen potentiſſimum & cum judicio dandum

℞. Rad. Eſul. min. ℥ß. Helleb nigr. ʒij Colocynth. in pet. ligat. ʒj. Caryoph. ℈j affunde vini ſ. q.

℞. Eſul. helleb. n. ana ʒvj helen. ℥j gentian. ℥ß. rhabarb. ʒvj fol. ſen. ℥j ⊖ ♀ri. ʒiij include nodulo, affunde vinum generoſum q. ſ. f. alterans & ſimul purgans, in Cachexia, febribus, hydrope, ſcorbuto.

HELLEBORUS NIGER.

Eſt Helleborus niger flore roſeo C.B.P. n.IV. Sunt radices tenues fibroſæ nigricantes ex uno capite prodeuntes, ſaporis acris pungitivi, amari nauſeoſi. Eſt planta multiſiliquoſa, pentapetala.

Viſcidos & Melancholicos ſeu acidos aduſtos & coagulatos humores validiſſime tam ἄνω quam κάτω expurgat, ut in morbis capitis, ſoporoſis, maniacis, Chronicis, Catarrhoſis, Hypochondriacis præ cæteris eligatur.

Fibras plurimi rejiciunt, reſervantque tantum cuſpidem ſuperiorem, ſed in hac tanta non eſt vis, quam in fibris.

Raro pro baſi præſcribitur, ſed potius tanquam ſtimulans pro cæteris purgantibus. In ſubſtantia datur à ℈j ad ʒß in robuſtioribus. In infuſo à ʒj ad ʒij ex vino cydoniato, oxymelle, tanquam correctorio. tutiſſimum eſt extractum, vel cum 🜋 vini vel aceto factum, quod eſt nonnihil reſinoſum.

Aquam deſtillatam purgativa acredine eſſe imbutam experimentis Zvvelfferi. [Pharm. Reg. p. 35.] Olai Borrichij & aliorum conſtat.

℞. Helleb. nigr. ſacch. cand. ana ℈j M. f. pulv. exhibe in cochleari oxymell. pro robuſto, vel add. ⊖ ♀ri gr. v. 🝆 de caryoph. gt. ij. M. f. pulv. pro maniaco.

℞. Extr.

℞. Extr. hell. n. gr. xv. ☿ dulc. gr. vij. diagr. gr. iij Camph. gr. v. Eſſ. Caſtor. q. ſ. f. pil. purgantes, valde conveniunt in chloroſi.

HELLEBORUS ALBUS.

Eſt Helleborus albus flore ſubviridi C.B.P. radix oblonga, pollice aliquando craſſior, foris fuſca, intus alba, numeroſis albicantibus fibris ſtipata, ſaporis acris, paululum amaricantis, ſubadſtringentis, ingrati & nauſeoſi.

Helleborus albus nigro multo vehementior eſt, & aliquando convulſiones parit, proinde ejus ſubſtantia non niſi robuſtiſſimis, maniacis, furioſis, Melancholicis, quartanariis concedenda eſt.

Corrigitur vel per acida, vel per vinum Malvaticum, in quo vel coquitur vel maceratur, iterumque ſiccatur, & ſic interne exhibetur, ſemper tamen genium ſuum retinet.

Externe ob ſui acredinem pulveribus ſternutatoriis admiſcetur ad excitandos Apoplecticos, & ſanandos Catarrhoſos.

Subſtantia robuſtiſſimo datur à ℈ß ad gr. xv. in infuſo ad ʒj melius tamen in extracto.

℞. Helleb. alb. ℈ß ⊖ ♀ri gr. vij. ☿ dulc. ℈ß L. O. gr. j. f. pulv. in mania exhibendus cum ſyr. de fumar.

℞. Galang. ʒij affund. vin. helleborat. ℥iiij f. Eſſentia cochleatim danda in mania.

℞. Extr. helleb. alb. gr. xv. ⊖ ♀ri gr. v. conſerv. fumar. ʒj M. f. bol. pro ij doſibus, robuſtiſſimo vero pro una doſi exhiberi poteſt.

℞. Helleb. alb. infundatur in vino Malvatico, eximatur radix & novum vinum ſuperaffundatur, hoc ſecun-

ſecundum vinum helleboratum confert in ſpecie maniacis ad cochl. j. vel ij in quo etiam varia medicamenta aliis morbis appropiata exhiberi poſſunt, v.c. Eſſent. caſtor. vel caſtoreum ipſum in Uteri vitiis, ſeu Eſſent. abſynth. in quartana.

℞. Helleb. alb. fol. majoran. ſummitat. origan. ana ℈j M. f. pulv. ſternutatorius ad excitandos Apoplecticos, in graviori caſu add. gr. j. vel ij Cantharid.

BRYONIA.

Eſt bryonia aſpera ſeu alba baccis rubris C. B. P. radix prægrandis, longa, craſſa carnoſa, albida, quaſi fungoſa, intus circulis donata, ſaporis acris mordacis amaricantis, nauſeabundi, odoris viroſi dum recens eſt, mitior fit dum ſeneſcit. Decocta ejus acria ſunt ſatis, extractum eſt optimum.

Eſt planta baccifera ſcandens, flores monopetali quinquefariam diviſi.

Recens & vegetativo ſuo ſucco adhuc turgida validior eſt quam ſiccata, hæc ſerum & aquas magis per inferiora, illa vero ἄνω & κάτω potenter exterminat. Speciale quid obtinet in ſanguinis grumis diſſolvendis, item in aſthmate & hydrope pectoris, in hydrope Uteri, ut & ob inſignem ſui amaritiem in vermibus.

Sicca infuſis diureticis additur pro ſtimulo, ſuccus radicis à ʒij ad ℥ß exhibetur cum oxymell. contra aſthma.

℞. Raſur. bryon. rec. ℥ij Camph. ʒij f. maſſa, excipiatur petiis in podagra.

℞. Succ. bryon. ʒij 🜖 thereb. ſpic. q. ſ. f. liniment. pro partibus convulſis.

In

In Clyſteribus contra hydropem. ℞. Rad. bryon. ℥j. irid. flor. ℥ß. cort. ebul. ſambuc. ana ℥ß. bacc. junip. ſem. cumin. fœnic. ana ʒij coq. ex ▽ & Urin. colatur. add. hier. picr. ℥ß. vitell. ov. n. j.

Externe radix recens ingreditur Emplaſtra ſeu Cataplaſmata contra tumores pedum œdematoſos & podagricos. ℞. Bryon. rec. ℥ij flor. cham. melil. ſambuc. ana Mj. f. catapl. ex ▽ & Urina.

℞. Bryon. ſicc. rad. ari ana ʒiß flor. ♃is ʒj f. pulv pro aſthmaticis a ʒß ad ℈ij ſumendus ex ſyr. violar.

℞. Rad. bryon. ℥ß. coq. cum oxymell. contra aſthma.

℞. Rad. bryon. ℥ß. abſynth. Mj. flor. viol. p. j. coq. in vino & aqua, applicetur totum parti dolenti in hemicrania.

*
* *

RADICES SALE PURGANTE BLANDO CONSTANTES ET VULGO LAXANTES.

EBULUS.

Eſt Sambuci humilis C. B. P. radix oblonga tenuis albida, ſaporis amaricantis, ſubacris & nauſeoſi.

Eſt planta baccifera trypirena.

Serum & ▽ tam per ſedes quam per Urinas blande evacuat, præcipue cortex unde antihydropicus audit. Subſtantia lignea interior adſtringit potius, & propterea in fluore albo & nimio Menſium fluxu ex vino capitur.

Externe valet contra tumores pedum œdematoſos ℞. piper. ʒj. Cort. Ebul. ℥j helen. bacc. junip. ana

℥ß. flor. chamom. melilot. ana Mj. coq. in ▽ & Urin. pint. j. f. Catapl.

Interne. ℞. Cort. Ebul. ℥ij bacc. junip. ℥j flor. ſambuc. p. j. infund. vin. ſ. q. ad ▽as per Urinam & alvum educendas.

POLYPODIUM.

Eſt Polypodii vulgaris C.B.P. radix oblonga, minimi digiti craſſitie, foris rufeſcens intus herbacea, ex viridi palleſcens, multis nodis & tophis quaſi acetabulis geniculata, ſaporis ſubacris dulcis, quodammodo adſtringentis & nauſeoſi.

Planta eſt Epiphyllospermos.

Optimum eſt quernum, præter virtutes cum filicibus & oſmunda communes poſſidet vim blandam laxantem, ideo Melancholicis, Hypochondriacis ſcorbuticis, tuſſientibus ſpeciatim conducit, præcipue ubi eorum alvus eſt adſtricta, externe ad oris & faucium vitia ſcorbutica.

In decoctis & infuſis frequentius datur circiter ad ℥j.

℞. Rad. Polypod. ℥j. Enul. camp. ℥iiij coq. in hydromell. & vin. ana ℔ij f. apozema, quod à Poterio in aſthmate commendatur, præcipue ubi alvus eſt adſtrictior, addi poſſunt fol. Nicot. ʒij.

℞. Polypod. ℥ij paſſul. min. ℥ß. ⊖ ♀ri ʒj coq. in ▽ Co latur adhuc calenti add. Mannæ ℥vj ſem. Coriandr. fœnic. cinamom. ana ʒij f. apozema purgans antihypochondriacum.

℞. Rad. Polypod. ℥ij Malv. hort. flor. roſar. ana Mj. herb. prunell. ſalv. ana Mß. coq. in ▽ ℔iiij add.

aceti

aceti circiter. ℔ß. & mell. roſat. q. ſ. f. gargariſma contra omnis generis inflammationes faucium, addi poteſt Alum. ℨj prunellæ. ℨij.

***

RADICES PARTICULIS DULCIBUS SEU BLANDE VISCOSIS CUM DULCEDINE CONSTANTES.

Dulcia concipiuntur corpuſcula rotunda, mollecula, ad latera leviſſimis flexilibus & obtuſis fibris inſtructa, linguæ meatibus cæteriſque poris citra ullam renitentiam ſe inſinuantia. Suntque vel alimentoſa, dum in chylum facile mutantur & poris partium aſſimilantur, vel medicamentoſa, & ob ſui texturam villoſam acria quævis, rigida & aſpera obtundunt, temperant, emolliunt & flexilia reddunt, ſanguinis nimium motum ſiſtunt, eumque nonnihil incraſſant, hinc in faucium aſperitate, ſiti nimia, defluxionibus acrioribus & ſanguinis æſtu convenientiſſima ſunt.

GLYCYRRHIZA.

Eſt Glycyrrhiza ſiliquoſa vel Germanica C. B. P. radix oblonga, ſarmentoſa, digiti craſſitie, coloris flavi, ſaporis ſubacris mellei dulciſſimi, tenuis odoris mellei; debilis amarities latet in cortice, hanc excipit levis acredo, unde colliguntur ⊖ia blanda.

Planta eſt ſiliquoſa papilionacea unicapſularis.

Copioſe colitur in agris Bambergenſibus, jam vero omnium optima ex inſulis Canarienſibus transfertur, quæ pugni craſſitiem adæquat.

Ob ſui dulcedinem acredine leni temperatam mulcet & incidit, ut proinde in faucium aſperitate,

 tuſſi

tuſſi ſicca, in Pleuritide dyſuria & ſtranguria, ut & in phthiſi in ſpecie laudetur; & ubi ſaccharata ob acetoſitatem non conveniunt glycyrrhiza præſcribitur: temperat quoque diaphoretica & diuretica, quæ ob ſui acredinem ſuſpecta habentur, infantum quoque alvum emollire & ſolvere creditur.

Datur in quavis formula. in ſubſtantia ad ʒj in infuſo & decocto ℥j.

℞. Rad. glycyrrh. ℥iiij Cinam. ℥ij coq. ex ▽ Hordei add. ferment. piſtor. petiis exceptum, repone donec fermentet, f. potus ad ſitim tollendam, addi quoque poſſunt paſſulæ minores.

℞. Rad. glycyrrh. Doronic. ana ℥ß tuſſilag. ℥ij ir. flor. ℥ß. herb. ſalv. fol. tuſſilag. ana Mj. jujub. n. vj. vel vij. ſebeſten. n. v. caric. ping. n. vj. ſem. cubeb. ℥ß f. c. ▽ & vino decoctum pectorale.

℞. Glycyrrh. alth. ana ʒß f. pulvis in ▽ chærefol. tuſſilag. & papav. exhibendus in Pleuritide.

DORONICUM.

Sunt Doronici radice ſcorpii C. B P. radiculæ oblongæ, rotundæ, tuberoſæ, nodoſæ, digiti minoris craſſitie, ex viridi palleſcentes, plures ex uno capite tenuibus quaſi filis dependentes, ſaporis ſubacris, ſubdulcis ad glycyrrhizam accedentis, odoris nullius: minorem habet dulcedinem quam glycyrrhiza, ſed ſaporem magis aromaticum. Hodie nobis eo nomine non venit, quo veteribus, venenatam enim qualitatem non habet.

Planta eſt diſcoidea, radiata pappeſcens.

Habet in dulcedine ſi recens ſit aliquid aromatici, quod cuminum redolet, ſeu ſemina nigellæ, & propterea

pterea ob hoc blandum aromaticum junctum dulcedini diſcutit vertiginoſos vapores, & involvit ſæpe exiſtentes in confuſo & inordinato motu.

Datur in ſubſtantia à ʒß ad ʒj in decocto vero more congenerum, ſed rarius datur in decoctis & infuſis. Jungitur viſco querno. Stimulatur galanga, cubebis, cumino, ⊖e Volat. ſuccin. Caſtoreo.

℞. Rad. Doronic. ℈j galang. ℈ß. ⊖ Volat. ſuccin. gr. v. caſtor. gr. vij. f. pulv. cum conſerv. fl. anthos & ſyr. fl. tunic. q. ſ. f. bolus. in vertiginoſis addi poteſt Laud. op.

In ſanguine congrumato ℞. Doronic. ʒj ex ▽ Chærefol. add. ⊖ Volat. C.C. Oc. ♋. ſperm. Cet.

* * *

RADICES AQUOSIS CRASSIS SEU MUCILAGINOSIS CONSTANTES.

Mucilaginoſa concipiuntur corpuſcula oblonga, mollia, minus mobilia ac aquea, multis obtuſis ad latera fibrulis inſtructa, quibus fluida quævis & acria quævis comprehendunt, ſtringunt, incraſſant & obtundunt, hinc virtute dulcibus accedunt, temperata ſunt, potentius acria obtundunt, fluida incraſſant, ſanguinis nimium motum ſiſtunt, rigida emolliunt &c.

ALTHÆA.

Eſt Althæa Dioſcoridis & Plinii C.B.P. radix oblonga, digiti craſſitie, in ramos divaricata, foris pallide cinerea, intus pallida, ex ſiccitate rugoſa, ſaporis mucilaginoſi & lenti, odoris nullius evidentis.

Planta eſt Pentapetala [monopetala, foliis diviſis] ſemilunaribus in rotunditate caſeolum exprimente ſeminibus diſpoſitis.

Agit in omnia ⊖ia, tam ſalſa quam acida, partim ea ▽eis fundendo, partim ⊖ina ſpicula involvendo.

Præter virtutes dulcibus cæteriſque mucilaginoſis communes, ſingulari proprietate tanquam anodyna confert affectibus Urinariis, ſtranguriæ, dyſuriæ, prohibet calculoſum acidum, & promovet generati calculi excretionem. Externe emollit, maturat & dolorem ſopit, quam ob cauſam ingreditur Cataplaſmata, Emplaſtra, Clyſteres &c.

Datur in ſubſtantia ad ℨj in decocto & infuſo ab ℥ß ad ℥j.

℞. Rad. alth. ℈j ſacch. cand. ℨß. M. f. pulv. in dyſuria add. nuc. moſch. ℈ß.

℞. Rad. alth. ℥ij fol. malv. Mij. paſſul. min. ℨvj coq. in ſ. q. ▽ f. apozema, emolliens, laxans ſi velis habere, add. fol. ſenn. ℨvj Crem. ♀ri. ℨiß. & colaturæ permiſce ſyr. d. ſpin. cervin. f. apoz. emolliens & laxans.

In fluxionibus acribus ſalſis. ℞. Rad. alth. lil. alb. ana ℥j tuſſilag. ℥ij herb. ſcabioſ. heder. terreſtr. ana Mj. fl. pap. rhœad. p. j. ſem. lin. fœn. græc. ana ℨij fœnicul. ℨiij fic. ping. n. x. dactyl. n. x. f. mitiſſim. decoctum ex ▽ vel vino.

In acido dyſenterico expugnando ℞. Rad. alth. ℨß rhab. toſt. ℨß ℅ nuc. moſch. expr. gr. iij. Laud. op. gr. j. f. pulv.

In Nephritide & in calculo, qui oritur à particulis alcalinis cum acidis. ℞. Rad. aperient ℥ß. alth ℥j glycyrrh. ℨij flor. Melil ſambuc. ana Mj fol. alth. Mj f. decoct. ex ſerolactis Colatur. ℔vj add. ſyr. dialth. ℥vj ⏜ ⊖ dulc. ad grat.

Vidi calculum excretum à puero ſex annorum bene-

neficio cataplaſmatis ſcroto impoſiti ex althæa, lacte & 🝆 lini expreſſo. Ingreditur etiam omnis generis Clyſteres.

LILIUM ALBUM.

Eſt lilium album flore erecto & vulgare C. B. P. radix bulboſa, ſquamoſa, intus alba, extus parum fuſca, ſaporis mucilaginoſi & lenti, odoris nullius.

Planta eſt caule multis foliis cincto, floribus ipſis erectis.

Lentorem habet blandiorem quam althæa, in clavis pedum, & induratis puerperarum pudendis ſingulare eſt. Anodynum eſt, ut & 🝆 ejuſque flores. Additur quoque Clyſteribus anticolicis.

℞. Rad. lil. alb. ℥j fol. Malv. Mj. flor. verbaſc. ſambuc. ana Mß. coq. in ▽ add. Colat. Elect. lenitiv. ℥ß ⊖ gemm. ʒj M. f. Clyſter in fœcibus duriſſimis.

SYMPHITUM.

Eſt ſymphitum ſeu conſolida major C. B. P. radix oblonga, ſurculoſa, pollicem craſſa, nigra, intus alba, ſaporis ſubdulcis mucilaginoſi & viſcidi, odoris nullius.

Planta eſt aſperifolia flore monopetalo cylindraceo, quinque laciniis diſſecto, ſeminibus quaternis.

Craſſiore mucagine pollet quam althæa, cum qua convenit, in ſpecie Vulnerarium eſt, antidyſentericum, & antihæmoptoicum & in exulceratione renum & veſicæ ex cantharidibus orta.

Datur ut althæa, ſed ob craſſiorem mucaginem in minore doſi.

In Clyſteres ſi recipitur, præſtat ut ſindoni involvatur, ne craſſior mucago decocto permiſceatur, &

inteſtinis juſto diutius adhæreat ac aceſcendo tormina producat; corrigitur & alteratur mucago Camphora.

℞. Rad. ſymph. ℥ß urtic. cum toto Mij affund. iij pint. ▽æ, coq. ad conſumtionem ij pint. bis vel ter ſolo hoc decocto D. Hermannus curavit hæmoptoën. Laud. opiatum quodvis extemplo levat ægrum ſi recidivam patiatur.

℞. Conſolid. maj. bol. armen. tragac. Maſtich. ana ℈j conſerv. roſar. r. antiq. ſyr. Corallor. q. ſ. f. Electuarium in hæmoptyſi.

Extra radix applicata confert in hæmorrhagiis, fracturis, luxationibus, herniis, in ſanguine coagulato ſingulare quid præſtare dicitur.

℞. Rad. torment. ſymph. ana ʒj veron. alchymill. plantag. ana Mj. ſummit. Hyperic. Mß coq. in ▽. ℞. Colatur ℔j. add. ſyr. d. ſymph. Fern. ℥j cape pro vulnerario interno. & in quovis hauſtu balſ. 🜂is gt. vij. vel x.

In phthiſi ℞. rad. Chin. ℥ij paſſul. min. ℥iß Conſol. maj. ℥ß. coq. in ſ. q. ▽æ.

Singulare hoc eſt contra herniam. ℞. ſem. naſturt. hort. ℥ß. gumm. carann. Mucag. ex ſymphit. extract. ana contra quamvis herniam recentem ex laxato peritonæo.

In fracturis oſſium intus exhibeo. ℞. oſteocoll. ʒß. cum raſur. conſol. maj. ℈j. cito certe conſolidantur. Externe unguentum ex ipſa mucagine addita oſteocolla mire prodeſt.

Contra fluxum alvi dyſentericum ℞. Conſolid. maj. ℥j torm. biſtort. ℥ß. herb. pyrol. alchymill. ſanicul. Mj. coque in ▽ chalybeata f. decoct. add. Balſ. 🜂 hyperic. addito ☍ roſar. f. Clyſter.

RADIX

***

RADIX CONSTANS MUCAGINE TENUIORE PAUCIS FIXIS SALINIS.

TUSSILAGO.

Eſt Tuſſilago vulgaris C.B.P. radix tenuis, longa, candida, ſaporis ſubacris, cum levi lentore, odoris nullius. Planta eadem quæ Petaſitis.

In blando ſuo lentore nonnihil ⊖ fovet, unde tam incidendo, quam obtundendo acria, ſpecifice prodeſt in phthiſi, tuſſi, tabe & pleuritide.

Datur in decocto ad ℥ij ex vino, lacte, cereviſia, Lindanus magno cum ſucceſſu decoctum ſolum exhibuit.

℞. Rad. tuſſilag. ℥ij flor. papav. err. Mj. bellid. min. Mj f. decoct. add. Mand. luc. piſc. ʒß ⊖ volatil. C.C. ℈ß. M.

℞. Rad. tuſſilag. ℥iiij alth. ℥ß. helen. ʒiij ſarſæpar. ʒv. herb. ſcabioſ. hed. terreſtr. veronic. pulmonar. capill. ♀is ana Mß. ſem. aniſ. fœnic. ana ʒij coq. in ▽ vel vino vaſe probe clauſo, colatur. add. ſyr. violar. vel ſcabioſ. q. ſ. f. apozem.

***

RADICES PARTICULIS TERREIS CONSTANTES.

Terrea concipiuntur corpuſcula irregularia, poroſa, gravia, fixa, ad motum inepta, quandoque ſecum vehentia quippiam ſubtilis vel acido auſteri, vel alcalico aromatici.

Cognoſcuntur quaſi a priori guſtu adſtringente, qui producitur dum ex poris & meatibus Linguæ ab-

ſorbetur & exhauritur fluidum & ſerum, unde pori concidunt, exiccantur. Vehunt in ſe ut plurimum ſubtile acido-auſterum, à quo ſeri fluiditas ſtringitur, quaſique figitur.

Hinc eorum effectus eſt ſanguinem cæteroſque humores fluidiores & tenuiores incraſſare & figere, ut & fibras partium coarctare conſtringere & roborare. Propterea conveniunt in nimio ſanguinis motu & æſtu, in omnibus fluxibus, Vulneribus, internis viſceribus roborandis. Sunt & nonnulla terrea ſimul alexipharmaca ſeu bezoardica, dum nimiam ſanguinis reſolutionem, quam venena maligna minantur, ſua adſtringentia impediunt. Potentius & felicius operantur, ſi ipſis quippiam acidi jungitur.

ALCANNA.

Eſt anchuſæ Monſpeliacæ radix oblonga, craſſa, foris rubicunda, intus alba, rubro colore inficiens, lignoſa, ſaporis dum recens paulum adſtringentis & inſipidi, odoris nullius. Secundo ſationis anno menſe Martio dum ſpiritibus ſuis vegetabilibus gaudet colligenda.

Ex radice Indi cum calce viva extrahunt tincturam, cum qua dentes & ungues rubro colore tingunt, etiam crines equorum.

BISTORTA.

Eſt Biſtortæ majoris radice minus intorta C. B. P. radix oblonga, pollicem craſſa, quaſi tuberoſa & geniculata ſeu nodoſa, nonnunquam retorta, foris ex ſpadiceo nigricans, intus ex flavo rufeſcens, ſaporis auſteriuſculi adſtringentis, odoris nullius.

Planta

Planta eſt pentapetala ſpicata, ſeminibus triangulis ſplendentibus.

Validius adſtringit quam alcanna, hinc potens eſt in fluxibus ſedandis & nimia ſanguinis reſolutione cauſata à malignitate, laudatur quoque in præcavendo abortu.

Datur in qualibet forma, in ſubſtantia ad ʒj decoct. ab ℥j ad ij. ſudores etiam movet, cujus ratio à priori dari non poteſt.

℞. Biſtort. ʒß. conſ. roſ. r. ʒj. ſucc. Citr. q. ſ. f. bolus adſtringens.

℞. Biſtort. ʒj exhibe ex ▽ Plantag. cochl. j. ad præcavendum abortum.

℞. Biſtort. tormentill. ana ℈j borrac. calcinat. ℈ß. conſ. roſ. r. q. ſ. in adſtringenda alvo. [it. in fluxibus ſeminalibus.]

TORMENTILLA.

Eſt tormentilla ſylveſtris C. B. P. radix tuberoſa, dura, nodoſa, pollicem craſſa, in obliquum acta, coloris ex fuſco rubeſcentis, ſaporis auſteriuſculi. Subtilior & paulo temperatior eſt quam biſtorta. In Variolis & Morbillis cum alvi fluxu habetur ſpecifica. Dicitur à nonnullis contra Apoplexiam remedium præſentaneum.

Datur in qualibet forma, in ſubſtantia ad ʒj. melius in decocto quam infuſo à ʒij ad ℥j. Extracto c. Spiritu Vini vel ▽ propria facto ℈j.

℞. Extr. torment. ℈ß. caſtor. gr. v. ⊖ vol. ſuccin. gr. vij. f. pil. in Variolis & morbillis.

℞. Torment. alcann. biſtort. ana ℈j M. f. pulv. add. conſ. roſ. r. ʒj f. bolus.

℞. Nuc. Moſch. toſt. ℈j ſacch. ♄ni gr. vij Laud. op. gr. ij. extract. torment. ℈j c. ſyr. roſ. r. q. ſ. f. pil. vel cum conſerva roſar. r. f. bolus in Dyſenteria.

CONTRAYERVA.

Eſt Clematis paſſionalis Peruviana folio bifido. ſeu Coanenepilli: Radix oblonga, tenuis, nodoſa, fibroſa, foris rubra, intus pallida, ſaporis nonnihil adſtringentis, cum ſuavi ſed debili acrimonia, odoris aromatici ſed exigui. Inter alexipharmaca reponitur. Commendatur magis quam cæteræ radices in morbis malignis.

Datur ut tormentilla in ſubſtantia, infuſo, extracto.

℞. Contrayerv. ʒß. camph. gr. iij ſyr. d. cichor. q. ſ. in Variolis & Morbillis.

℞. Contrayerv. ʒß. Corall. r. ʒj f. pulv. adſtringens excip. conſerv. roſ. r.

℞. Contrayerv. ℥iiij ſantal. r. ℥ß. ♒ theriacal. ℥iiij ♒ ⊕l. dulc. ℥ij ♒ vin. Camph. ℥iij. f. Tinctura Bezoardica. addi quoque poſſunt radices angelicæ, valerianæ, petaſitidis.

CARYOPHYLLATA.

Eſt Caryophyllata vulgaris C. B. P. radices tenues, fibroſæ, ex uno capite craſſiore prodeuntes, ſaporis ſubacris, grate amaricantis, odoris modo aromatici caryophyllati, modo nullius, pro diverſis quibus colligitur locis, odoratior quæ in locis creſcit excelſis, quam quæ in hortis colitur.

Planta eſt pentapetala, ſeminibus in capitulum congeſtis, nudis crinitis.

Recens

Recens & in calidioribus locis nata blandumque aroma caryophyllatum ſpirans, poſſidet in ſubſtantia terrea aliquid Salis volatilis, ob quod contra Catarrhos & capitis obſtructiones commendatur. Sicca vero & ⊖e volatili carens, blande adſtringendo fibras relaxatas, præcipue Ventriculi, confortat. In ſpecie commendatur à nonnullis in ictero à Bile acri & viſcoſa. Stimulatur ſerpillo, majorana.

℞. Caryophyllat. ℥j majoran. ſerpill. ana p. ij affund. ♒ Ceraſ. nigr. f. infuſum pro Catarrhis.

℞. Rad. caryophyllat. ʒß. Caryoph. arom. ℈ß conſ. roſ. q. ſ. f. bolus.

PIMPINELLA.

Eſt Pimpinellæ ſanguiſorbæ minoris hirſutæ C.B.P. radix calami craſſitie, ex fuſco ſubrufa, ſaporis ſubamari, adſtringentis, tenuis non ingrati odoris. Culta in hortis fit delicatior longe. Gaudet foliis pinnatis, flore ſtamineo, ſeminibus ut plurimum quadrangulis.

Reſpondet caryophyllatæ. Hujus radicis uſu hydrophobia fuit curata.

℞. Rad. pimpinell. ʒß. C.C. præpar. ℈j M. f. pulv. addita conſerv. roſar. r. f. bolus. addi quoque poteſt L. O. gr. j. item ☍ Nuciſt. expr. vel magma poſt deſtillationem reſiduum. Exhiberi poteſt in quolibet fluxu.

PENTAPHYLLI RADIX.

Eſt quinquefolii majoris repentis flore luteo C.B.P. radix oblonga, calami craſſitie, foris rubicunda intus incarnata, tranſverſim ſecta crucem vel ſtellam

lam referens. Saporis primum adſtringentis, poſtmodum incidentis. Odoris nullius.

Planta eadem eſt quæ tormentilla, conſtat terreis particulis, volatilibus paucis. Subſtituitur cortex hujus radicis aliquando cum effectu cortici ChinæChinæ, ſi ſumatur ad ʒj.

Subſtantia hujus interior eſt lignoſa, & tota ſtyptica. Cortex vero ob ſui acredinem & ⊖is præſentiam in febribus intermittentibus & quartanis laudatur. Movet ſudorem & Urinam, datur ad ʒij. Ingreditur gargariſmata ad detergenda ulcera, & curandas aphthas.

In decoctis datur pro mictu cruento h.m.

℞. Rad. pentaphyll. ℥ij herb. agrimon. ſummit. hyperic. ana Mj coq. in Cerev. ſ. q.

℞. Rad. pentaphyll. ʒß. ⊖ prunell. ℈j M.

℞. Rad. pentaph Cort. Ch. Ch. rad. gentian. ana ℈j ⊖ prunell. gr.x. M. f. pulv. contra febres add. ſacch. ♄n. gr. vij. L. O. gr. j,

NYMPHÆA.

Eſt Nymphææ luteæ vel albæ majoris C.B.P. radix craſſa, foris ex fuſco flaveſcens, intus pallida, fungoſa, levis, ſaporis adſtringentis auſteri, odoris nullius.

Planta eſt pentapetala, capitulo oblongo ſtriato verrucam exprimente, multicapſularis, in quibus ſemina ſplendentia, triticeo grano minora.

Habet in ſubſtantia levi, poroſa & laxa, nonnihil ⊖is volatilis alcalini, nam ſolutiones cum acidis factas præcipitat, unde & acidorum actiones, phlogoſes cauſantes, temperat & ſiſtit, mitigatque ſpiritus 🜁eo-acidos, & propterea in nimiis vigiliis tanquam ſpecifica

ca extollitur. Ob eaſdem cauſas peculiare quid obtinet in ſiſtendo fluxu ſeminali, pollutione nocturna, ♀ea, inflammationibus renum & veſicæ, ut & in febrium æſtu.

Datur decoctum, ▽ ſuccus, ſyrupus & ipſa ſubſtantia ad ʒj decoctum ad ℥ij.

℞. Rad. nymph. ℥iiij flor. nymph. Mj. anagall. flor. phœniceo Mij coq. ex vino auſtero vel ▽ fontan. f. apozema contra maniam.

℞. Camph. gr.v. rad. nymph. gr.xv. ſacch. ♄ni gr.vij. L.O. gr. ij. excip. conſerv. flor. nymph. & ſyr. menth. q.ſ. f. bolus in mania & quovis delirio.

℞. Cort. ſaſſafras ℥ij rad. Nymph. ℥iij flor. Nymph. p. ij, coque ex vino contra fluorem album mulierum.

*
* *

RADICES TERREIS ET SIMUL MUCAGINOSIS VOLATILIBUS PARTICULIS CONSTANTES.

POEONIÆ RADIX.

Eſt pœoniæ maris folio nigricante ſplendido C.B.P. radix craſſa, tuberoſa, foris fuſca, intus pallida, ſaporis dum recens ſubacris, ſubdulcis, cum levi quadam amaritie, odoris nullius.

Planta eſt multiſiliqua polypetala, ſeminibus lucidis nigris oblongo rotundis.

Hujus vis antepileptica à Medicis decantata, conſiſtit forſan in modica adſtringentia, juncta blando volatili lentori, à quibus contra acidum Epilepticum cerebrum corroboratur, ſpirituumque impetuoſus & violentus motus ſedatur. Eandem ob cauſam laudatur

tur etiam in motibus convulſivis, vertigine, tremore, pavore puerorum nocturno, tanquam prodromis Epilepſiæ.

Datur in qualibet forma, additur quoque Clyſteribus antepilepticis. In ſubſtantia à ʒß ad ʒj. Decoctum ab ℥j ad ℥ij.

℞. Rad. pœon. ſem. pœon. ana ʒij viſc. quern. ʒj ⊖ volatil. C. C. vel ⊖ Urinæ vol. Cinnab. ♁ij ana ℈j in robuſtioribus ʒß. M. f. pulv. pro vj. doſibus: cape ex ▽ flor. tunic. ℥iij.

℞. Rad. pœon. ℥j angel. acor. ℥ß. ſummit. origan. ſtœch. rutæ Mj. coq. in ▽.

℞. Colatur. ℥x add. ⊖ vol. C.C. ʒij f. Clyſter. contra vertiginem & Epilepſiam.

℞. Rad. pœon. ℈ß. Cinnab. ♁ij gr. iiij. M. f. pulv. contra pavores nocturnos infantum.

CYNOGLOSSA.

Eſt cynogloſſi majoris vulgaris C. B. P. radix oblonga, craſſitie digitali, foris ex atro rubens, intus albida, ſaporis fatui muſcoſi, aliquando ſubdulcis, odoris graviuſculi, aliquid fœtidi canis redolentis, narcotici, ſi creſcit in locis uliginoſis, in ſiccioribus inodora fere, admota naribus ſomnum conciliat.

Planta eſt aſperifolia pentapetaloides, ſeminibus lappaceis compreſſis. Eſt ſolani ſpecies. Mulcet acres Catarrhos, & in dolore capitis, fluxu albo in gonorrhœa ut & in aſthmate commendatur.

Externe ſcrophulas & ſtrumas colli curat, & ſi recens ac canino odore adhuc fœta abigit pediculos. Imponuntur nempe infantibus folia cynogloſſæ Capiti ad abigendos pediculos.

Suſpecta

Suſpecta à nonnullis virulentia, idcirco corrigitur cinamomo, croco, Caſtoreo. Vis narcotica exaltatur opio, aliiſque narcoticis.

Pilulæ de cynogloſſa dantur à gr. vij ad ix per ſe. vel

℞. Pil. de cynogloſſ. gr. ij. ▽ papav. rhœad. ℥j ſyr. pap. rhœad. ʒij f. hauſtus ad tollendum capitis dolorem, ubi opium eſt ſuſpectum.

MANDRAGORA.

Eſt Mandragora fructu rotundo C.B.P. radix craſſa, oblonga, bifida, foris cinerea ferruginea, intus pallida, odore ingrato, viroſo, caput feriente & narcotico. Cortices cinereo ferruginei ſcabri, intus albidi.

Planta eſt baccifera, floribus monopetalis in quinque ſegmenta diviſis, ſeminibus compreſſis rugoſis.

Mandragoræ radix duplici notatur virtute. 1. purgante tam per ſuperiora quam per inferiora, cum violentia & virium etiam proſtratione & aliquando convulſionibus ſubſequentibus unde per ſe non datur, niſi doſi minima ad ℈ß. ℈j. Extractum ejus cum Spiritu Vini ex corticibus factum datur pro anodyno. 2. Narcotica ſtupefactiva, quæ in mucagine volatili conſiſtere videtur, à qua ſpiritus æque ac ab opio involvuntur. Inde forſan eſt, quod in Epilepſia & motibus convulſivis quibuſdam commendata ſit.

℞. Cort. Mandrag. Gentian. ana ℈j. Cort. Chin. Chin. ⊖ abſynth. ℈ß. f. pulv. propinandus ex vino. vel

℞. Rad. gentian. ſerpent. Chin. Chin ana ℈j. Mandragor. ℈ß. ⊖ abſynth. gr. v. f. pulv. pro j. doſi quo agyrta curavit febres tertianas & quartanas.

℞. Cort Mandragor. ʒij ſabin. Mj, aſar. ʒj coq. ex Cerevіſia pro expellendo fœtu mortuo.

Empyricis laudatur in quartanis pro Emetico.

Uterum valde movet & abortum cauſatur, unde gravidis non danda. Veteres utebantur cortice infuſo cum vino pro narcotico & ſtupefactivo, præcipue ubi membra erant amputanda. Datur tantum robuſtioribus cum circumſpectione.

℞. Extract. opij. Rad. pœon. cort. mandrag. ana ʒij viſc. quern. ℥j ſem. pap. alb. & nigr. ana ʒiij C.C. coral. r. ana ℥ß. M. f. p. cujus doſis ℈j contra Epilepſiam ſeu convulſiones ſeu impetus ſpirituum animalium. Exhibendus ex ... u Ceraſorum nigrorum. Externe in doloribus dentium à putredine commendari poteſt ſequens.

℞. Cort. mandragor. ʒß. opij ℈j. pil. peſt. Ruffi ℈j ℅ d. caryoph. gt. v. f. turund. in momento ſiſtit dolores.

[Joh Jacob. Manget. in Biblioth. Pharm. Medica ſeqq. profert de Mandragora ex Herman. Mat. Med. MSta. Mandragora eſt planta pentapetala, baccifera, ſeminibus compreſſis rugoſis, quæ nota cum omnium ſolanorum ſit, Mandragora ad ſolanorum ſpecies referri debet. Incipit florere primo vere, quo tempore radix utplurimum efficaciſſima colligitur. Ex odore viroſo colligitur ejus vis narcotica, incidens & & attenuans, ſed omnis hæc vis eſt in cortice, & medulla inferior rejicitur, unde etiam in officinis nonnullis cortices tantum in umbra exſiccati proſtant. Specificum uterinum eſt, ſed caute adhiberi debet, ſin in nimia exhibeatur quantitate dira inde ſequuntur ſymptomata ut ſpaſmi &c. in pulvere doſis eſſe poteſt ʒj (forte ℈j) in infuſo ʒj. Circumforanei utuntur contra febres intermittentes ſed minus tuto. Extractum ejus valet contra tertianam & quartanam. Præparata præterea nulla.]

CAPUT

CAPUT SECUNDUM.
DE CORTICIBUS.
CORTICES CONSTANTES SALE VOLATILI OLEOSO.

Cortex Aurantiorum.
Citri.
Cinamomi.
Caſſiæ Ligneæ.
Caryophyllatæ.
Winteranus.
Guajaci.

CORTEX AURANTIORUM.

HOrum exterior flavedo ⊖e volatili oleoſo fœta, in flatibus & torminibus abdominis præcipue colicis, in paſſione hyſterica, ſtranguria, dyſuria, præſidium ſpecificum eſt. Interior autem fungoſa ſubſtantia ob ſui adſtringentiam in dyſenteria & nimio fluxu menſium prodeſt.

Datur flavedo in ſubſtantia ad ʒß it. 🝆 tam deſtillatum quam ex flavedine expreſſum ad gt. ij vel iij in ▽ appropiata deſtillata, addito ſimul ſyrupo ſaporis gratia.

Amſtelodami proſtat 🝆 ex flaved. Cort. aurant rec. expreſſum, quod ex Hiſpania transfertur, & venditur ſub nomine quintæ eſſentiæ aurantiorum, magis

laudatur ac ☍ deſtillatum. Viget quoque eſſentia, extractum, infuſum.

℞. Flaved. Cort. aur. ʒij Mand. luc. piſc. Ocul. ♋. præpar. ana gr. xv. ſuccin. præpar. ʒß. Zedoar. ℈ß ☍ expr. cort. aur. gt. v. vel vj. M. f. pulv. anticolicus cujus ʒß adjungi poſſunt L. O. gr. ij. ut & in colica convulſiva Caſtor. gr. vij, exhibe ex ▽ appropriata, ſive Tinct. cum Spiritu Vini ex corticibus facta.

℞. Elæoſacch. ☍ Cort. aur. ʒß. Zedoar. gr. v. Nuc. Moſch. gr. vj vel vij M. f. pulv. capiendus ex vino ſeu Cerev. ſeu ▽ chæref. contra ſtranguriam & dyſuriam.

℞. Cort. aurant. inter. fungoſ. flavedine rejecta, coq. forti △e cum ▽ ſ. q. ſume decoctum contra nimium alvi & menſium fluxum.

CORTICES CITRI.

Horum flavedo fragrantior & ſpirituoſior eſt, quam corticum aurantiorum, & propterea in ſyncope, lypothymia & capitis debilitate ad confortandam memoriam ſingulari cum ſucceſſu adhibetur. Videtur etiam aliquid volatilis acidi in ſe habere, & propterea antiſcorbutici audiunt. Dantur ut cortices aurantiorum.

℞. Eſſent. ſ. ♌us Citri ʒij ▽ rub. id. ʒiij acet. theriacal. ʒij ſyr. rub. id. ℥ß. M. f. potus in virium languore & ſyncope, add. conf. alk. ʒß.

℞. Cort. Citr. ʒß. ſem. Citr. Card. Bened. ana ʒij ▽ Papav. q. ſ. f. Emulſ. in febribus malignis.

℞. Flaved. Cort. citr. ʒij Cort. Winteran. ʒß. in fund. in vin. Malvatic. f. infuſum contra ſcorbutum.

CINA-

CINAMOMUM.

Eſt Caſſiæ Zeilanicæ C.B.P. cortex tenuis, ab exteriore pellicula purgatus ex flavo rubeſcens ſeu ferrugineus, in cannulas contortus, ſaporis acris ſuaviter pungentis, ſubdulcis & aromatici, odoris fragrantiſſimi.

Arbor eſt baccifera calyculata nucleo quercino.

Ex radice arboris Cinamomi in inſula Zeilan paratur camphora. [conf. act. Hafnienſi. Vol.3. obſ.23. p.37. ubi Paul. Hermannus noſter allegatur, & Camphoram triplicem ex cortice radicis cinamomi inſulam Ceylon ſuppeditaſſe. 1. Crudam. 2. arte per calcem, 3. per ſe ſublimatam. Duplex oleum, alterum è cortice radicis, alterum è foliis cinamomi elicitum.

Sale ſuo volatili ſpirituoſiſſimo ∽bus animalibus valde homogeneo, junctum habet ſubdulcem blandumque lentorem, hinc præter cardiacam & analepticam virtutem, in tuſſi & pulmonum muco laudatur, tam ob vim incidendi, quam ob lenientem & mitigantem. In Menſium ſuppreſſione & lochiorum fluxu, in corrigenda purgantium violentia cæteris omnibus præfertur.

Vis cinamomi concentratior eſt in pellicula interiori corticis, cui ſubjacet lignum. Nimia hujus acredo ubi inconveniens videtur optime temperatur hordeaceis.

Datur in qualibet forma ut cortices aurantiorum. In ſubſtantia à ʒß ad ʒj it. in infuſo, rarius in decocto, niſi pro aſthmaticis.

℞. Cinam. ʒij Zinz. ʒj flaved. Cort. Citr. Aurant. Nuc. Moſch. ana ℈j. M. f. pulv. & cum vino infuſum confortans.

℞. Cinam. ʒß. borrac. ℈j. M. f. pulv. dandus ex ▽ puleg. ad Menſes & lochia pellenda.

℞. Cinam. ʒß. Myrrh. gr. viij. borrac. ℈ß. ☍ cinamom. gt. j. M. f. pulv.

℞. Cinam. ℥ij oxym. ſcyll. ℥j. M. f. hauſtus in aſthmate & tuſſi.

℞. Gumm. ammoniac. ʒß. ſolv. in ▽ Cinam. ℥iij add. ſyr. puleg. ʒiij f. hauſtus in aſthmate extra paroxyſmum.

℞. Cer. alb. q. v. diſſolv. in ☍ been ſ. q. fiat corpus pro balſamo, huic inſtilla ☍ cinam, pro balſamo conficiendo, ſi velis balſamum hunc tingere nativo colore, adde magiſterii iſtius ſ. q.

℞. Sacch. cand. ℥j ☍ d. cinamom. gt. v. vel vj. ambr. gryſ. gr. v. Moſch. gr. j. M. f. pulv. analepticus ad ℈j exhibendus. Ex ▽ Cinam. Meliſſ. rub. id.

℞. ☍ exp. bacc. laur. ℥ß. deſt. ſuccin. gt. xv. ☍ cinam gt. vij M. f. liniment antiparalyticum.

℞. ⊙ Cinam. ex deſtill. reſid. ʒj fungoſ. ſubſtant. Cort. aur. ʒß. M. f. p. contra Dyſenteriam vel diarrhœam add. Nuc. Moſch. ℈ß L. O. gr. j.

CASSIA LIGNEA.

Eſt Caſſiæ ligneæ Malabaricæ ſeu Sumatrenſis bacciferæ cortex, cinamomi in modum convolutus, cinamomo craſſior, odore & ſapore debilior, cum majore aliquantum lentore. Differt à cinamomo non niſi loco natali, utrumque ex arboribus prægrandibus

dibus inſtar quercus noſtratis, dum ſex annorum eſt, deraditur, cortices hi dum recentes ſunt albicant, beneficio caloris rufeſcunt, & in cannas ſua ſponte convolvuntur.

[De differentia Caſſiæ ligneæ & cinamomi conf. Andr. Cleyer. C. N. C. Dec. II. an. IV. obſ. 6. p. 8. it. d. arbore cinamomi, & caſſiæ ligneæ Theodor. Carol. C. N. C. Dec. II. ann. 8. obſ. 59. p. 149.]

Utraque arbor fructus fert olivæ inſtar, baccas nempe rotas ſebaceas, quarum oleo expreſſo utuntur Indi loco ſebi in candelis, nullius eſt odoris. Hi fructus ſunt magnitudine baccarum lauri, membranæ pulpoſæ incluſi, ſapore juniperino præditi. Flores habent odorem liliorum convallium.

Ratione virtutis aromaticæ debilior eſt, aſt ratione viſcoſitatis efficacior eſt quam cinamomum hinc convenit in aſperitate faucium, tuſſi, inflammatione uvulæ.

Datur ut cinamomum, aſt in doſi majori, in thoracis affectibus præſtantior Cinamomo.

℞. Caſſ. lign. ℥ij paſſul. min. ℥ß. M. f. infuſum ex vino, ſeu leve decoctum ex ▽, contra aſperitatem faucium & tuſſim.

Si recens eſt, fundit 🝆 ſat copioſum, quo adulteratur 🝆 Cinamomi.

CASSIA CARYOPHYLLATA.

Eſt Myrti arboreæ Americanæ Caninge cortex cinamomo tenuior, ex rufo ferrugineus, à pelliculis ſuis exterioribus deraſus, & in modum tubuli convolutus, ſapore acri aromatico, odore ad caryophyllos accedente.

Arbor

Arbor eſt baccifera, baccis rotundis umbilicatis acinos geminos includentibus, ſapore & odore caryophyllorum.

Virtute caryophyllos æmulatur, quibus & hodie ſubſtituitur, habet autem in receſſu multum viſcoſitatis, à qua ⊖ia ejus volatilia multum temperantur. Additur tanquam roborans & confortans diureticis & hydragogis contra hydropem, ut & medicamentis Uterinis ad accelerandum partum.

Datur in ſubſtantia à ℈j ad ʒß.

℞. Caſſ. caryophyllat. ℥ß. Caſtor. ℈j Camphor. ℈ß. f. infuſum hyſtericum in partu laborioſo, admotum naribus, vel cochleatim datum.

℞. Caſſ. Caryophillat. ℥ß. Saſſafr. ℥j radic. vincet. Valerian. ana ʒvj bacc. Junip. ʒv. ſem. fœnic. ʒiij rhabarb. ʒiij infund. in vin. rhenan. f. infuſ. diureticum, hydragogum.

CORTEX WINTERANUS.

Ita appellatur ab Admirali claſſico qui ipſum in freto Magellanico invenit. Non eſt Coſtus, ſed arboris Magellanicæ bacciferæ cortex craſſus, cannulatus, foris integumento cinereo fungoſo inæquali, fiſſo & multis rimis hiante, intus ſolidus, denſus, ad piperis & cochleariæ ſaporem accedens, odoris fragrantis.

[Eſt Laurifolia Magellanica cortice acri C.B.P. Magellanica aromatica arbor Cluſ. Exot. Nonnullis appellatur peryclimenum odoratum.]

Habet ⊖ volatile 🝆ſum cum levi amaritie & acido volatili, in Anglia ſub nomine corticis antiſcorbutici præſcribitur. Naſcitur ferme in locis inacceſſibilibus, unde mirum tantam ejus eſſe hodie copiam.

Salibus

Salibus volatilibus penetrantiſſimis & ſpirituoſiſſimis gaudet, potenter incidit & attenuat, quidpiam acidi volatilis videtur admixtum, & propterea in ſcorbuto ſpecificum eſt. Conducit inprimis in ructibus & Vomitibus acidis. it. in Paralyſi quod confirmat Willis ex Experientia. [Pathol. c. 9. commendat Cereviſiam medicatam ex ſalv. beton. ſtœch. lign. ſaſſafr. & Cort. Winterano.]

Datur in ſubſtantia à ʒß ad ʒj in infuſo ad ℥ij.

℞. Cort. Winteran. ʒß ari præpar. ℈ß. Oc. ♋. ℈j M. f. p. pro ij. doſibus add. L. O. gr. 1. contra Vomitiones acidas.

℞. Cort. Winteran ʒij rad. armorac. ʒiß. fol. cochl. Mj. caſſ. Caryoph ʒiij Cort. Citr. ʒij f. infuſum in ▽ cochleariæ. Seu ſi vinum non noceat affunde vinum Malvaticum.

℞. Cort. Wint. ʒj ſem. Citr. ʒiij Cort. aur. ʒj ſeri lactis pint. iß. f. infuſum in ſcorbuto.

In amiſſa loquela. ℞. Pyrethr. utr. ℥ij ſaxifrag. ℥ij Cort. Winter. ℥ß. ſumm. origan. p. ij f. decoctum ex ▽ & vino. adde colaturæ Eſſent. Caſtor. ℥ß.

Suppeditat oleum copioſum, quod ut ☍ cinamomi partim ſupernatat, partim fundum petit.

CORTEX GUAJACI.

Sunt Guajaci officinarum C.B.P. cortices, tenues, duriuſculi, lignoſi ex pluribus lamellis congeſti, ex cinereo rufeſcentes & nigricantes & maculati, intus pallidi, nitidi & quaſi lævigati, ſaporis acris amaricantis, odoris fere nullius.

Arbor eſt hexapetala nucifera, fructu cordis figura, binis vel trinis nucleis membranæ corticoſæ involutis. Sponte ſua naſcitur in India occidentali.

Corticcs Guajaci nobiliores habent virtutes quam lignum, & conſtant tam ⊖ibus volatilibus quam fixis utroque Balſamo & reſinæ permixto. Horum ratione ſudoriferi ſunt & diuretici, ac in Morbis Chronicis, Catarrhis, odontalgia, ſcorbuto, ſcabie maligna, lue ♀ea mire laudantur. Dantur in decocto, in quod Balſamicam aromaticam vim deponunt.

Oleum deſtillatum & rectificatum, ac cum proprio ſpiritu cohobatum, ſingularis Medicina eſt in affectibus hyſtericis, colicis, febribus malignis, & lue ♀ea, & dicitur Q. E. Guajaci.

Reſina vel ex corticibus extracta, vel ſua ſponte emanans [quæ eſt arida, friabilis, ſubnigricans, pellucida, ſaporis acris, odoris grati] medetur lui ♀eæ & gonorrhææ.

℞. Cort. Guajac. ℥ß Lign. Guajac. ℥iiij paſſul. min. ℥j lign. al. ʒj coq. in ▽ Miiij ſub fin. add. raſ. lign. ſaſſafr. ℥ß. ſem. aniſ. ʒiß. Cinam. ʒij M. f. apoz. in omnibus Morbis Chronicis.

N. quod contra luem ♀cam huic decocto poſſit addi ☿ viv. & ♁ij crudi ana ℔ß. petiis except.

℞. Reſin. Guajac. ℈j ☿ virid. gr. v. ſacch. ♄ni. ℈ß. ſuccin. præpar. gr. xv. Thereb. ſ. Balſ. Peruv. q. ſ. M. f. pil. contra gonorrhæam pro j. doſi.

CORTEX PERUVIANUS SPURIUS CHINÆ CHINÆ FALSÆ.

Sunt arboris cujuſdam Peruvianæ Kinkina dictæ, cortices Kinkina vera tenuiores foris cinerei caneſcentes, intus ferruginei, ſaporis amaricantis & ſimul ſubpinguis, odoris debilis. Potentior eſt & nobilior ratione virium quam china vera, habet ⚯ gratioſiſſimum.

Sale

Sale tam fixo quam volatili cum levi quadam reſinoſitate ſcatet, & præter cæteras facultates quam cum aromaticis communes habet, poſſidet virtutem antiſcorbuticam, & antihyſtericam.

℞. Chin.Chin. falſæ & veræ, ⊖ abſynth. ſ. ⊖ prunell. ana ℈j M. f. pulv. divid. in 2. doſ.

℞. Chin. Chin. falſ. ℥ij Cort. Winter. ℥j Caryoph. arom. ℈j rad. ari. ℥j M. f. cum vino infuſum ſtomachicum & antiſcorbuticam.

℞. Chin. Chin. falſ. cinam. ac. ana ℈j Croc. or. gr. vij. borrac. ℈ß. M. f. pulv. antihyſtericus pro ij doſ.

℞. Elæoſacch. Chin. Chin. falſ. exbibe contra paſſionem hyſtericam.

***

CORTEX SALE FIXO AMARISSIMO CONSTANS.

CHINÆ CHINÆ.

Sunt arboris cujuſdam Peruvianæ Gannanaperide dictæ cortices ex craſſitie mediocri, foris ſcabri, paululum caneſcentes ſæpe muſco obducti, intus læves colore ferrugineo ſeu cinamomeo.

N. Kinkina cum China vera nullam habet convenientiam, ſed quod male pronunciatur à nobis China Chinæ audit.

Arbor eſt procera, craſſa, tiliæ inſtar baccifera.

[De utroque hoc cortice conf. Dale Pharmacol. part. I. p.m. 372. ubi inquit: ſunt qui duas corticis ſpecies eſſe affirmant, videlicet veram, & ſpuriam ſeu falſam, utramque autem ex India occidentali adferri, prima ſeu vera ita deſcribitur cortex craſſitie mediocri foras ſcaber, paululum caneſcens, aliquan-

do muſco obductus, intus lævis coloris ferruginei, ſaporis acris intenſe amari.

Altera ſeu falſa tenera eſſe dicitur, foras coloris cinerei & caneſcentis, intus ex fuſco aut obſcuro ferruginei, ſaporis acris amaricantis ſubpinguis, odoris fragrantis. Quæ deſcriptiones cum Autoris noſtri conveniunt.]

Ob ⊖ia fixa rigidiſſima & amariſſima ſubſtantiæ terreæ & adſtringenti incluſa inſignem obtinet antifebrilem vim, & in ſpecie contra quartanam intermittentem. Quidam volunt hunc corticem eſſe etiam Uterinum, cum aliis præſcriptum.

Datur tam in ſubſtantia ad ʒß quam in infuſo ad ʒij. Præmiſſis purgantibus convenit, vel his tempeſtive interpoſitis, plurimis enim relinquit alvum adſtrictam, unde ſemper purgantia adjunguntur.

Jungitur Gentianæ, centaur. min. abſynth. ſerp. virgin. Stimulatur ⊖e Card. B. abſ. Cent. min. ⊖ prunell. adduntur quoque huic cortici terrea, ſcil. Corall. præpar. ſuccin. &c.

℞. Cort. Ch. Chin. ʒj. rad. ſerp. virg. ℈j ⊖ prunell. gr. xv. M. f. p. antifebr. capiendus ex ▽ taraxac. bihorio ante paroxyſm. vel ex Tinctura bellidis.

℞. Ch. Chin. ℥ß. gentian. ʒiß. flor. cent. min. p. I. pip. long. ⊖☿ri ana ʒij M. f. pulv. cujus ʒj exhibeatur ex ▽ Cichor. vel taraxac. vel acetoſ. bihorio ante paroxyſmum.

℞. Cort. Chin. Chin. ʒij Gentian. ſerp. virg. ana ʒj helleb. n. ʒiß. affund. vin. generoſ. ad pint. iß. f. infuſum, de quo ſingulis diebus bis vel ter ſumendum.

℞. Lap. bez. ℈j vel Bezoard. min. ℈j Chin. Chin. ʒß. lap. prunell. gr. xv. 🝆 d. caryoph. gt. I. f. p. antifebrilis.

CORTICES

⁂

CORTICES CONSTANTES SALE FIXO DIURETICO.

CORTEX FRAXINI.

Sunt Fraxini excelſioris C.B.P. cortices ex craſſitie mediocres, ex cinereo viridantes, ſaporis acris amari, cum adſtrictione.

Lignum ejus ipſo Johannis Baptiſtæ die cæſum à multis habetur vulnerarium, [it. ad hæmorrhagias] Parantur & ex hoc ligno cyathi contra omnis generis venena.

[De ligno hoc vulnerario ſeu ſympathetico dicto prolixe agit, ejuſque vires recenſet Hohberg Kunſt-Buch num. 37. p.30. annex. Georgic.Irioſ. Tom. II. ſed multa hyperbolica & ſuperſtitioſa enarrantur.]

Ob ⊖ acre diureticum & diaphoreticum convenit in Morbis Chronicis, paſſione hypochondriaca, ſcorbuto, nephritide, hydrope, aſcite, Catarrhis, Lue ♀ea & aliis.

Datur decoctum ab ℥iij ad iiij quod aliquando ex vini & aquæ anatica portione paratur. Viget quoque ⊖ per incinerationem factum doſ. à ℈j ad ʒß in Variolis & Morbillis.

℞. Cort. Fraxin. tamariſc. ana ℥j rad. helen. armorac. polypod. ana ℥ß. helleb. n. ʒij ſummit. abſynth. Cent. min. fol. agrim. ana Mj. bacc. Junip. ℥iß. affund. vin. Rhenan. ſ. Malvat. pone in digeſt. f. infuſ. antihydropicum & antiſcorbuticum.

℞. Cort. fraxin. ℥iiij coq. in vin. ℔iiij f. apoz. ex quo cape ⊖ fraxin. gr. vij. l. ℈ß contra hydropem & Urinæ ſuppreſſionem.

CORTEX TAMARISCI.

Eſt tamarix fruticoſa folio craſſiore ſeu Germanica C.B.P. Conſpirant cum corticibus fraxini & cum his ſimul ſolent præſcribi, utrique cortici aſſignatur ſpecifica vis contra vitia Lienis. Unde ex ligno tamariſcino conficiuntur pocula, ex quibus Lienoſi omnia medicamenta & potulenta capiunt. Externe cataplaſma ex corticibus Lienoſis applicatur, admodum conducit, præcipue ſi addatur cortex capparis.

CORTEX CAPPARIS.

Eſt Capparis ſpinoſa fructu minore, folio rotundo C.B.P. Sunt radicum cortices tubuloſi, rugoſi, craſſi cinerei, ſaporis acris auſteri.

Cum tamariſci & fraxini corticibus convenit, niſi quod auſteritatis ratione in receſſu plus adſtringat, datur in ſubſtantia ad ʒj in ▽ Mulſa, oxymell. ad viſcerum obſtructionem ex ſanguine acido aduſto. Decoctum datur ad ℥iiij.

℞. Cort. Cappar. ℥iiij flor. cappar. muria cond. coq. in ▽ Mulſa, vel oxymell. ſeu vino f. apoz. ad induratum lienem.

℞. Cort. cappar. fraxin. tamariſc. ana ℥j herb. cuſcut. epith. plantag. capill. ♀is ana Mj f. decoctum ex ▽ ad quemvis morbum ex obſtructione meſenterii dependentem.

Proſtat & ☍ coct. ex capparibus. ℞. ☍ coct. cappar. diſſolv. gum. ammoniac. ad emolliendos tumores viſcerum.

CORTI-

*
* *

CORTICES PARTICULIS TERREIS CONSTANTES.

CORTICES CASTANEARUM.

Sunt Caſtaneæ ſativæ & ſylveſtris C.B.P. cortices rubicundi, caſtaneam fructum ſeu nucleum immediate cingentes. Eſt arbor nucifera, pluribus in communi integumento echinaceo incluſis.

In ſpecie conducit in ſiſtendo nimio menſium fluxu & metuendo abortu, ut & in gonorrhæa & fluxu albo. Ingreditur quoque gargariſmata refrigerantia & adſtringentia.

Ex pulvere horum corticum cum pinguedine quadam vel oleo appropriato, paratur unguentum, quod in nimio menſium fluxu pudendis illinitur.

℞. Cort. Caſtan. ℥ij auric. jud. ℨj hord. excort. ℥ß. flor. malv. arbor. p. j. coq. ex ▽ plantag. add. colatur. ſyr. violar. q. ſ. ⊖ prunell. q. ſ. f. garg. in inflammatione faucium.

CORTICES GRANATORUM SEU MALICORIUM.

Sunt fructuum mali Punici C.B.P. cortices coriacei duri, rugoſi, foris ex ſpadiceo luteo punicei, intus vero lutei, ſaporis amaricantis adſtringentis auſteri. Eſt arbor pomifera, fructu rotundo, in ſummitate umbilicato, floribus fructui inſidentibus pentapetalis.

Acido-auſterum ⊖licum, particulis terreis admixtum, potenter exaltat horum corticum virtutem adſtrictoriam, propterea conducit in fluxu albo, hæmorrhagiis & profluvio ſeminis. Convenit quoque

in

in Vomitu nimio tam ſimplici quam cruento, ut & gravidis pica laborantibus.

Ingreditur quoque gargariſmata & clyſteres adſtringentes. Externe ad ſuffuſiones Oculorum & leucomata, ut & ad pudenda exulceratata laudatur. Præcavent quoque ne Oculi à Variolis & morbillis lædantur.

℞. Cort. granat. ʒj N. M. toſt. ℈j. Laud. op. gr. j. M. f. pulv. contra omnem fluxum alvi, exceptus conſerv. roſar. ⊕lat. f. bolus.

℞. Cort. granator. ʒß. ſuccin. præpar. oſteocoll. anà ℈j f. pulv. exhibe ex ſucco granator. tam ad fluorem album, quam ad nimium menſium fluxum.

℞. Cortic. granator. ʒß. exhibe ex ſucco granator. ſeu cydonior. contra picam. f. add. miv. cydonior. qt. v. addi poſſunt Zedoariæ grana aliquot.

CORTICES QUERCUS.

Sunt cortices craſſi, foris ex cinereo fuſci, intus pallide rubentes, ſaporis adſtringentis.

Subſtantia terrea craſſiore conſtant quam cortices granatorum, cui tamen non nihil ⊖is alcalini videtur admixtum, decoctum cui aliquid ⊕li adjicitur nigreſcit. Conveniunt in fluxu alvi ut & hæmorrhoidum, teneſmo, item in ſoda, & acida efferveſcentia temperanda. Horum decoctum ex ▽ chalybeata factum datur pro Vulnerario, de cætero conveniunt cum corticibus granatorum.

℞. Cortic. querc. ℥iiij calic. gland. ℥j. coq. ex ▽ chalybeata; add. ſyr. Cortic. granat. q. ſ. f. apozem. contra ardorem Ventriculi.

Paulo ſubtilioris Naturæ ſunt calices glandium, quam cortices quercus, cum quibus in omnibus conveniunt. Et cum his ſimul & ſemel præſcribuntur.

CAPUT

CAPUT TERTIUM.

DE LIGNIS.

LIGNA CONSTANTIA SALIBUS OLEOSIS AROMATICIS.

LIGNUM SASSAFRAS,

Est arboris Virginianæ folio ficulneo Pavame dictæ. Radix deberet proſtare, nam melior eſt, & ſic deſcribitur, ſub cortice tenui, foris cinereo intus ſanguineo, includens materiam ex albo rufeſcentem, lignoſam, aliquantulum ponderoſam, ſaporis ſubacris, ſubdulcis, tenuis aromatici, odoris fragrantis & fœniculati.

Transfertur ex Virginia & Florida. Dicitur ſaſſafras quaſi ſaxifraga. Ob virtutem lithontripticam. Duplex ejus ſpecies, altera folio ficulneo, floribus pentapetalis, habentibus odorem aniſi, quos flores ſequuntur baccæ, & hæc uſualis eſt. Altera non, cum fructibus corni, exceptis foliis valde convenit.

Hujus ⊖ia volatilia temperata ſunt fœniculacea dulcedine, ut tam per diureſin quam diaphoreſin diminuat etiam in remotiſſimis locis viſcoſa, craſſa & vitioſa. Hinc tanquam Catholicum in Morbis Chronicis uſurpatur. In ſpecie corrigit crudam acidamve lympham, [Brunnero in conſiliis audit verum alexipharmacum Catarrhorum] medetur tuſſi, tabi, phthiſi, lui ♀eæ, & in ruptis abſceſſibus internis ſingulare eſt mundificans, ſpecificum eſt in fluore albo.

℞. Rad. helen. ℥ß. lign. ſaſſafr. ℥ij f. decoct. ex vino, quotidie exhib. ad ℥ij add. ʒij Cort. Winteran. ſi ſcorbutus adſit. In hoc exhibe ʒß ſuccin. præpar. in fluore albo. Hoc modo Hermannus puellam curavit.

Datur infuſum, decoctum, eſſentia & oleum deſtillatum.

℞. Raſ. lign. ſaſſafr. ℥ij flor. pap. rhœad. p. ij coq. leniter, add. colatur. ſyr. papav. q. ſ. f. apozema contra Catarrhos.

℞. Cort. ſaſſafr. ℥iß Spir. Vinoſ. ſerpilli ℔ij f. infuſum, quod præſervat à Catarrhis eoſque curat.

℞. Saſſafr. lign. ſanct. rad. ſcorzon. liquirit. ana ℥ij caric. ping. n. ix. f. decoctum mundificans ſanguinem à rupto abſceſſu vel à pleuritide, vel alia cauſa orto.

LIGNUM GUAJACUM ET SANCTUM.

Guajacum eſt arboris Indiæ Occidentalis Guajacan dictæ, lignum ſolidum, denſum, reſinoſum, buxei coloris, medulla intus nigricante duriſſima & denſiſſima, ponderoſa & reſinoſa, [ſaporis acris, amaricantis & aromatici, odoris fragrantis.]

Sanctum eſt arboris Indiæ Occidentalis Hoaxacan dictæ, lignum ſolidum denſum Guajaco albidius, medulla intus ſolida duriſſima &c.

Differt à Guajaco ſpecie non genere, utraque enim eſt ejuſdem figuræ & faciei, utraque prægrandis eſt & nucifera, deſtillat ex utraque etiam reſina. [Reſina arida eſt friabilis, pellucida, nigricans, ſaporis acris, odoris grati.]

Utra-

Utraque hæc ligna quoad vires corticibus reſpondent, & cum his ſimul præſcribuntur in decocto, eſſentia &c. proſtant & eorum oleum & ſpiritus.

℞. Lign. ſanct. cort. mal. granat. ana ℥j thur. maſtich. ana ʒij flor. balauſt. roſar. r. ana Mj. bulliant. in ▽ ſ. q. ad medietatem, f. apozema contra fluorem album, add. tantillum vini auſteri. Pro abluendis pudendis & injiciendo.

Decoctum in Lue ♀ea. vid. ſupra in capite de radice chinæ. p. 42.

✠ Oleum deſtillatum externe applicatum ſingulare eſt in nephritide. Commendatur etiam ℴℴ deſtillatum exterius cum ℴℴ bacc. laur. ſpic. lavend. in arthritide.

LIGNUM ALOES.

Sub nomine Calambac tranſmittitur ex Tartariæ intimis oris & in regnum Sinæ transfertur. vid. du Vall. Reiß-Beſchreibung part. 1. p. 507. Sunt arboris Chinenſis Calambac interioris medullæ fragmenta lignoſa, magnitudine varia, ſolida, denſa, ſpadicea, ſtriis aliquando nigricantibus & reſinoſis conſpicua. Saporis ſubacris, amari, aromatici & pinguis, odoris dum accenditur vel incaleſcit fragrantis. Præſtat ſi annoſum, grave, reſinoſum, æquale eſt. Si odoris fragrantia adeſt à principio accenſionis ad omnimodam conſumtionem uſque. Ex integra ſæpe arbore, quæ quercu magnitudine non inferior eſt, vix ℔ß ligni laudabilioris eruitur, unde tam charum. Chinenſes maxime ejus odoratu delectantur, & quo gratior amicus eo meliorem portionem ipſi offerunt.

Lignum aloës grave, totum reſinoſum & prunis injectum, ſuaviſſimum ſpirat odorem, nullo vel pauco relicto carbone.

Salia volatilia reſinæ ſociata adeo analoga ſunt ſpiritibus noſtris, ut inter analeptica & confortantia catholica habeatur, inprimis Cephalica, ſtomachica, cardiaca & Uterina. Hinc in Lypothymia, Syncope, ſingularem habet energiam, & tanquam ſpecificum confortans jungitur appropriatis ſtomachicis & Uterinis. Ad fluorem album compeſcendum, & abortum præcavendum: ſiſtit quoque Vomitum & ſingultum: curat incontinentiam Urinæ à ſpirituum defectu ortam: laudatur in Epilepſia, & necandis vermibus. Datur in ſubſtantia ℈j in decocto ʒj, ſed decocta reddit nimis amaricantia. Extractum & eſſentia facta cum -^-u vini, vel formicar. vel roriſmar. vel Ceraſ. nigror. habetur optima.

✠ Unà cum ambra hoc ligno aloes ad memoriam confortandam Profeſſor quidam Leidenſis uſus eſt cum ſucceſſu.

✠ ℞. Extr. lign. aloës ℈j margar. præpar. ℈j ambr. gryſ. in ☍cinam. ſolut. gr. vij ſacch. cand. ℥j M. f. pulv. cui add. ☍ cardamom. gt. vij ad ℈j ex ▽ Cephalica uſurpandus, in memoria confortanda, in virium lapſu, ſyncope cardiaca.

℞. Lign. aloës ℈ß theriac ℈j ☍ Mac. gt. ij Laud. op. gr. j. M. f. bolus ad compeſcendum & ſiſtendum vomitum.

℞. Lign. aloës gr. vij Corall. rubr. ſuccin. præpar. Maſtich. ana ℈ß. f pulv. adſtringens & confortans uterum, in fluoro albo, adde ad ſingulas doſes ambræ gr. j. valet quoque in incontinentia Urinæ.

AGAL-

AGALLOCHUM.

Sunt arboris Calambac affinis, fragmenta lignoſa, ſolida, ex fuſco rufeſcentia, ſtriis interdum albicantibus notata, bituminoſa, ligno aloës laxiora & craſſiora, ſaporis amaricantis pinguis, odoris debilioris, earundem cum ligno aloës virtutum, ſed gradu longe debiliori.

[De Agallochi generibus & arbore excœcante conf. Georg. Everhard. Rumphius C. N. C. Dec. II. ann. 3. obſ. 22. p. 74.]

SANTALUM CITRINUM ET ALBUM.

Citrinum eſt arboris Timorenſis bacciferæ Sercanda dictæ cor ſeu medulla, denſa, ſolida, colore citrino ab integumentis ſuis cortice & ligno ſeparato, ſaporis aromatici amari, odoris fragrantis & grati.

Album eſt ejuſdem arboris, [à multis affirmatur ex una eademque arbore produci, ſcribit tamen Garcias ab Horto arom. hiſt. l. 1. c. 17. p. m. 88. tantam eſſe inter utriuſque ſantali arbores affinitatem, ut pallidum à candido diſcerni non poſſit, niſi forte ab ipſis incolis, qui eas cæſas mercatoribus vendunt] lignum ſolidum & ponderoſum pallidum, non proſtat amplius apud nos [apud eund. Garc. l. c. p. 90. in notis: dubitandum an legitimum ſantalum pallidum habeamus, quandoquidem ejus pretium majus ſit apud Indos quam Luſitanos. Fieri igitur poterit ut aliquod aliud odoratum lignum nobis pro legitimo ſantalo obtrudatur. Certe candidum noſtrum plane inodorum]. Indi ſcrinia ex eo conficiunt, eſt enim ſolidius.

Indi unguento ſantalino ſeſe tingunt præcipue puellæ, ſantalum citrinum contuſum inque ſubtiliſſimum pulverem redactum, in oleo recenti inſolant, leniterque inſpiſſant, non debet nimis incaleſcere aliàs ranceſcit, & ex curcumæ radice colorem addunt, gratiſſimi hoc eſt odoris. [Ad æſtus corporis tollendos & odoris conciliandi cauſa. Eſt enim regio admodum calida, & incolæ plurimum odoribus delectantur Garcias l. c.]

Citrinum potentius eſt & odoratius albo, ϴia ſua volatilia tenuia & blanda multis terreis & paucæ reſinæ habet incluſa, hinc præter vim incidentem & Urinam moventem obtinet quoque virtutem adſtringentem & roborantem. Et propterea tam inter deobſtruentia & antihydropica refertur, quam etiam inter adſtringentia, quæ conveniunt in nimio ſudore, præcipue febrium ſudatoriarum, compeſcendo. Hujus æquivalens eſt lignum rhodium, ſtimulatur corticibus Citri.

Datur ejus decoctum, rariſſime ſubſtantia, valet quoque eſſentia cum ♒u vini, melius cum ♒u Cortic. Citri facta.

℞. Rad. helen. ℥j Cichor. ℥ij polypod. ℥ß. ſantal. citr. ʒvj herb. capill. ling. cerv. ana Mj bacc. Junip. ℥j coq. ex ▽ & vino f. apoz. contra obſtructiones viſcerum.

℞. Lign. ſantal. ℥ij Cort. Citr. intern. ℥j lign. aloës ʒß. M. f. decoct. ad nimios ſudores nocturnos & febres ſudatorias.

LIGNUM RHODIUM.

Eſt Aſpalathi albicantis torulo citreo C. B. P. ſeu Cytiſi potius, Cytiſi incani ſiliquis falcatis C.B.P. ab inventore

ventore Bartholomæo Marantha, Cytiſus Maranthæ dicti radix lignoſa, ex flavo pallida, denſa, ſolida, venulis pinguibus ſubinde interdiſtincta, ſaporis amari, odoris fragrantis gravis & roſei.

Fructus eſt ſiliquoſus, papilionaceus, naſcitur in inſulis Canarienſibus.

In radicibus ſunt conſpicui nodi pingues, in quibus maxima vis, ⊖ia ſunt temperatiora quam in ſantalo, in pellenda Urina paulo potentius eſt ſantalo. In oleo ejus deſtillato blanda vis cardiaca eſt, & nonnihil anodyna. Externe convenit cum roſis in mitigando dolore capitis.

Lignum datur in infuſo, ꝏ deſtillatum [odoramentis ſacrum] additur pulveribus dentifriciis & ſternutatoriis & Balſamis cardiacis.

℞. Pumic. calcin. ʒj oſ. ſæpiæ ʒij Corall. alb. ʒß. ꝏ lign. rhod. gt. iij ⊖✳ci ℈j ○ uſt. ℈ß. pro pulvere dentifricio. vel conſerv. cochl. q.ſ. pro antiſcorbutico.

℞. Lign. rhod. ſantal. Citr. ana ℥j bacc. Junip. ℥ij liquirit. ʒiij f. infuſum vel leve decoctum ex ▽ gramin. ad promovendas Urinas.

℞. Lign. rhod. ℈iiij ▽ roſar. ℥j beton. ℥ij + roſac. ℥ß. f. Epithem. ad mitigandos dolores capitis præcipue febriles, addi poteſt ſacch. ♄ni ʒß. it. ſantal. rubr.

*
* *

LIGNA CONSTANTIA SALE BLANDO DIURETICO.

LIGNUM NEPHRITICUM.

Lignum peregrinum aquam cœruleam reddens C.B.P. Eſt arboris Mexicanæ Coarthi vel Coalti dictæ, lignum

lignum ex flavo pallidum, durum, cœruleo tingens colore, includens ſub cortice nigricante medullam fuſcam, ſolidam, duram, gravem, ſaporis ſubacris, amari, aromatici, vim incidentem in lingua relinquens.

Naſcitur in regno Mexicano pyri vel pruni magnitudine, foliis rutæ alatim diſpoſitis, floribus papilionaceis, quos ſequuntur ſiliquæ retiformes includentes ſemen. Arbor refertur ad Cytiſi genera, unde & vocatur Cytiſus Mexicanus.

Robert Boyle tract. d. coloribus. [Part. 3. exp. x. p.m. 78. ad 82. it. pag. 124.] multa cum ipſo fecit experimenta.

▽ hujus ligni 10 vel 12 ♂ poſt infuſionem colorem diverſiſſimum demum acquirit, imo uni unciæ ter vel quater nova poteſt affundi ▽ & tingitur. Color vero debet ad radios ſolares ſpectari, ubi erit elegantiſſimus. Infuſum ex vino Urinam & alvum movet, ex ▽ vero ſola Urinam.

Agyrtæ adſcribunt ei vim alexipharmacam, tincturam enim ejus præbent gallinæ, cui prius dederant arſenicum, & ſervantur.

Ob ſal blandum alcalicum, quod ligni tinctura cœrulea à quovis acido deſtructibilis probat, ſed reſtituitur affuſo ℴℴ ♀ri per deliquium, diureticis, antinephriticis, & antihydropicis annumeratur.

Datur decoctum, infuſum ſeu tinctura cum vehiculo conveniente, ſcil. liquore betulæ, ▽ graminis &c.

℞. Raſ. lign. Nephrit. ℥j affund. ▽ gramin. vel al. deſtillat. ℔ß. f. infuſum ad nephritidem.

℞. Raſur. lign. nephrit. ℥iiij liquor. betulæ ℔ijß Elæoſacchar. citr. q. ſ. ad gratiam, contra dolorem nephriticum.

℞. Raſ.

℞. Raſ. lign. nephr. cort. fraxin. ana ℥j ſantal. Citr. ℥ß. infund. ▽ gramin. prius impragnatæ ⊖e geniſt. f. infuſ. diureticum contra hydropem & obſtructiones viſcerum.

[Nic. Monardes l. ſimpl. ex india allator. ad renum vitia, Urinæ difficultates ac arenulas pellendas uſurpari in nova Hiſpania ait: quod confirmat Boile l. c. h. v. infuſione ipſius in limpida facta contra renum calculos uti ſolent: & ſane inſignis quidam Medicus mihi familiariter notus, qui explorando iſti morbo enixius incubuit, mihi aſſeverat, ſe infuſionem ejuſmodi comperiſſe unum ex efficaciſſimis remediis, quæ unquam adverſus morbum iſtum terribilem propinavit.]

***

LIGNA AMARISSIMA ET SALIBUS FIXIS CONSTANTIA.

LIGNUM COLUBRINUM.

Quod proſtat in officinis potius eſt radix quam truncus. Eſt arboris Timorenſis radix lignoſa, brachium craſſa, ſub cortice ex fuſco ferrugineo ſubinde maculis cinereis notato, includens materiam ſolidam, denſam ponderoſam, ſaporis amariſſimi, odoris nullius.

Ob inſignem ſui amaritiem, ſymbolizat cum gentiana, Centaurio minore, aliiſque amaris.

Fructus hujus arboris ſunt nuces Vomicæ majores, aliæ enim Timorenſes, aliæ Zeilonenſes minores, ſunt venenum, pulpæ luridæ ingratiſſimæ ſunt incluſæ.

Arbor eſt pomiferis affinis, fructu rotundo in cortice tenui duriuſculo includente mucaginem, cui nu-

clei compreſſi, orbiculati involuti ſunt 3 vel 4 ſibi invicem incumbentes. Hujus ligni uſus non eſt, niſi annoſioris, recens enim primo anno maniam, tormina, hinc Vomitum & tandem convulſiones excitat. Laudatur quidem in febribus intermittentibus, quartanis, nec non vermibus necandis, ſed deleteria gaudet facultate.

Datur in infuſo ex vino abſynth. ſ. ex ▽ Card. Bened. impraegnata ⊖e Card. Ben. corrigitur amarities pipere longo.

✠ ℞. Lign. Colubrin. ℥ß. infunde in vin. abſ. per noct. f. hauſtus contra quartanam.

✠ ℞. Lign. colubrin. ℥ß. ſumm. abſynth. p. j. piper. long. ʒj ▽æ Cent. min. impraegnatæ ⊖e abſynth. ℔j f. infuſ. contra febres.

℞. Lign. colubrin. ſem. cinæ ana ℥ß ſumm. abſynth. p. j. ⊖ abſ. ℈j. f. infuſ. ex ▽ gramin. cum qua ▽ ſingul. doſ. cap. ☿ dulc. contra vermes.

[Hîc non abs re judicium Caſp. Hofmanni de Med. offic. l. 2. c 162. p. m. 402. ſubjungendum videtur, ubi de Nuce Vomica agit: In hac rerum incertitudine, cum certo utentis periculo, quid faciendum eſt ei, qui facultates rimatur? quam incerta ſunt iſta; tam certum eſt, quod jam dicam: Utrumque eſt venenum ſi enim ſemel bene ceſſit, non ob id tutum eſt experimentum. Ut in Politicis qui ſemel malus ſemper praeſumitur talis, ita in Medicina non utar incertis, cum certa prae manu habeo. & ibid. lib. 2. c. 4. p. m. 89. ubi de Doronico diſquirit an venenatam qualitatem habeat, concludit tandem, qua de re dubitas, ne feceris.]

LIGNA

LIGNA TERREIS CONSTANTIA.

SANTALUM RUBRUM.

Eſt arboris ſiliquoſæ Indiæ orientalis cor ſeu medulla, ab integumentis exterioribus cortice & ligno ſeparata, ſolida, denſa, ponderoſa, coloris ex fuſco rubicundi, ſaporis & odoris nullius evidentis.

Arbor eſt ſiliquoſa, floribus papilionaceis affinis.

Ob vim terream & adſtrictoriam, commendatur ad nimium ſudorem compeſcendum & ſanguinis æſtum. Foreſtus laudat in phthiſi tanquam corrigens & temperans ſalſam lympham. Externe recipitur ad Epithemata pro capitis dolore & ardore, ubi ſudor in capite ſimul erumpit.

Datur frequentius decoctum, rarius in ſubſtantia.

℞. Crem. 🜿ri ʒß. 🜿r. ⊕lat. ℈j ſantal. rubr. gr. vj. 🝐u ⊕li irroretur f. pulvis, ad roborandas Ventriculi fibras, dicitur bibitorius.

[Coincidit cum pulvere Hepatico rubro Dreſd. quem deſcribit Frid. Hoffmann. Th. Pharm. p.m. 98.]

℞. Decoct. ſantal. rubr. ℔ij ſucc. Citr. rec. ℥ß. ſyr. roſ. rubr. ℥ij ▽ papav. ℔ß. Cinam. ℥j f. Julapium refrigerans in febrium æſtu, maxime in ephemera.

℞. Santal. rubr. ℥ß. Citr. ℥ij ſaſſafr. ℥iß coq. ex ▽ hordei, f. apoz. addi poſſunt paſſul. min. ℥ij, ad temperandam acriorem lympham in phthiſi.

℞. Santal. rubr. ʒj flor. roſar. r. Mj +. roſac. q. ſ. M. f. Epith. ad capitis dolores febriles, add. in nimio æſtu ſacch. ♄ni. ʒj & ubi febris maligna camphor. ℈j.

VISCUS QUERNUS.

Sunt arbuſculorum quercubus innatorum ramuli lignoſi, digiti craſſitie, ſub cortice tenui includentes materiam lignoſam pallide rubentem, odoris & ſaporis nullus.

Viſcus hic eſt planta paraſitica, nullum ſuccum ex terra haurit, ſed ex aliis corporibus. Viſcus corylinus eſt optimus.

Ratione terrearum particularum acida abſorbentium, & nimium ſanguinis motum ſiſtentium convenit in hæmorrhoidum, alvi & menſium fluxu, ut & inflammationibus internis. Forſan ob eandem rationem commendatur in Epilepſia & Convulſionibus. Hujus vis ab autoribus contra incantationes decantata prorſus imaginaria videtur.

N. quoque poſt uſum viſci contra Epilepſiam utendum eſſe convenientibus purgantibus, ne acida in ſua quaſi minera à viſco nimis fixata, febrem excitent.

℞. Viſc. Quern. ungul. alc. ana ʒß. Cran. human. ℈j ⊖ volat. C. C. gr. xv. M. f. p. add. Caſtor. ℈ß. balſ. Copaiv. q. ſ. f. pil. contra Epilepſiam, & ſuperbibe cochl. j. ▽æ Epilepticæ, vel flor. til. vel lil. convall.

℞. Viſc. quern. ʒj cape ex ovo ſorbili contra nimium menſium fluxum.

℞. Viſc. quern. ʒj borrac. calc. ℈ß. f. pulv. exhibendus cum vino ſaſſafr. in fluore albo.

℞. Viſc. quern. ʒj. ſucc. expr. bugl. q. ſ. f. bol. in Dyſenteria ex mente Williſii.

LIGNA

***

LIGNA SALE PURGANTE TAM FIXO QUAM VOLATILI CONSTANTIA.

PANAVA SEU LIGNUM MOLUCCENSE.

Panava [ſeu Pavana, utrumque legitur apud ſcriptores] eſt Ricini arboris indicæ lignum ſpongioſum, leve, rarum, pallidum, tenui cortice cinereo obductum, ſaporis acris mordacis & cauſtici, odoris nauſeoſi.

Arbor eſt tricoccos, floribus muſcoſis, prunorum magnitudine.

Recens & adhuc vegetum adeo potens eſt, ut ſerum & aquas tam per Vomitum, quam per alvum expellat, relinquens ob intenſam ſui acredinem in ano inflammationem. Siccum vero alvum blandius ſubducit, & ſi in parciori doſi exhibeatur ſudorem elicit, in hydrope, leucophlegmatia & pluribus Morbis chronicis ſpecificum eſt.

Pro ſcopo purgandi datur recens in ſubſtantia ℈j ad ʒß. annoſum vero datur ad ʒj. In infuſo & decocto ad ℥ß. Jungitur variis purgantibus, ſtimulatur optime granis Tilli, corrigitur Zinzibere. Temperatur vero mellitis, ſaccharatis vel paſſulis. Pro ſudoribus movendis datur à ℈ß ad ℈j. In infuſo vero & decocto ad ʒij. ʒiij. Additur egregie decoctis ſudoriferis pro ſtimulo.

℞. Panav. rec. ℈j gr. till. gr. j. Zin. gr. vij. ſacch. cand. ʒß. M. f. p. loco ſacchari cape mellis vel conſerv. roſ. q. ſ. f. bolus contra hydropem.

℞. Raſ. lign. guajac. ℥iiij ſaſſafr. ℥j lign. Panav ℥ß. paſſul. min. ℥ij coq. in ſ. q. ▽. f. decoctum in obſtructione chronica.

In paſſione hyſterica ſumunt Indi radic. Curcumæ ʒj terunt ſuper lapide add. Panavæ ʒß. f. pulv. pro iij doſibus ex infuſo Thee vel lavendulæ.

Grana Tilli ſunt fructus hujus arboris à Garcia ab Horto eſt deſcripta, ſed non ſcivit hanc arborem ferre grana Tilli.

CAPUT QUARTUM.

DE FUNGIS.

OMnes fungi ſeu ſpongiæ venenatæ cenſentur, quoniam in earum ſubſtantiam denſam & quaſi coriaceam lympha ſtomachalis digeſtrix agere nequit.

BOLETUS CERVI.

Eſt ſubſtantia fungoſa, ſub terra naſcens, tuberoſa, rotunda, inæqualis, in magnitudinem Nucis Moſchatæ excreſcens, includens inter corticem coriaceum calloſum craſſiuſculum ; ex cinereo nigritantem, in ſuperficie granulatum, medullam raram, fungoſam, ex nigro fuliginoſam, ſubinde purpuraſcentem, & per maturitatem tandem abeuntem in pulverem, ſaporis & odoris dum recens gravis muriatici ſpermatici, dum ſicci nullius evidentis.

Recentiſſimus nondum ſiccus cujus ſubſtantia interna adhuc mollis & mucaginoſa eſt, ſcatet ⊖e volatili, ob odorem ſpermaticum ſpecificum eſt aphro-

diſiacum augens genituram & orgaſmum in ea excitans. Datur ſubſtantia interior cortice rejecto ʒj. Ex vin. Malv. ▽ Magnanimitatis ſ. ⏜u formicar. in infuſo ad ℥j.

℞. Bolet. cerv. rec. ʒj. ſuccolad. q. v. diſſolve in lacte calido ad languorem ♀eum.

℞. Bolet. cervin. ʒj. Elect. diaſatyr. ʒij ambr. gryſ. gr. vij ſyr. gran. cherm. q. ſ. f. Elect. aphrodiſiacum.

Pulvis in boleto contentus Balſamo Apoplectico nigrum ſuum conciliat colorem.

⁂

FUNGUS TERREIS SUBAUSTERIS CONSTANS.

AURICULA JUDÆ SEU FUNGUS SAMBUCI.

Eſt ſubſtantia fungoſa, naſcens ex imo trunco ſambuci veteris, ſpongioſa, levis, coriacea, membranacea, auriculæ inſtar, ſubtus incana & cinerea, ſupra nigricans, ſaporis terrei & aquei inſipidi, odoris nullius.

Ingreditur gargariſmata & epithemata tanquam potens adſtringens & refrigerans, contra inflammationes faucium, anginam, Aphthas, ut & Eryſipelas.

℞. Auric. Jud. q. v. infunde ▽æ deſt. ſambuc. f. infuſum ad inflammationes oculorum.

℞. Auricul. jud. ʒij hord. mund. glycyrrh. ana ʒß. flor. malv. arbor. p. ij. coq. ex aqua plantag. ℞. Colatur. ℔j add. lap. prunell. ʒj ſyr. violar. ℥j. f. gargariſm. ad inflammationes faucium.

FUNGUS

FUNGUS PURGANS.

AGARICUS.

Therebinthina eſt ſuccus collectus Laricum & abietum juniorum. Agaricus naſcitur in laricibus, & ſi creſcit in his arboribus, tunc therebinthinam non amplius reddunt. Naſcitur ex trunco laricis, rarius in ramis invenitur. Eſt excreſcentia fungoſa ex rotundo anguloſa, inæqualis, pugni magnitudine, candidiſſima, leviſſima, friabilis, paucis fibris intertexta, ex cortice calloſo cinereo ſubrufo, cui incluſa exempta. Saporis primum ſubdulcis, mox amari acris & nauſeoſi, cum levi quadam adſtrictione.

Reſinoſus adhuc melior eſt albiſſimo & leviſſimo, ⊖ purgans magis volatile quam fixum reſinæ, per -n- vini extrahendæ ſociatum in levi farinacea ſua terra fovet. Immerſus in acido efferveſcit, & in terram Cretaceam abit, uſtulatus ⊖e purgante orbatur, & in inertem terram mutatur.

Bilem & ſerum viſcoſius per alvum, interdum & Vomitum educit, & in ſpecie tuſſi & aſthmate laborantibus conducit.

Ejus vis purgans tardior ſtimulatur diagrydio, additur etiam ☿ dulc. Ob levem ſui ſubſtantiam ubique ſupernatantem & è Ventriculo facile ad fauces regurgitantem & nauſeam cauſantem, ſub forma pulveris vel hauſtus non exhibetur, ſed potius ſub forma trochiſcorum, vel pilulari. Ejus uſus in ſubſtantia ſuſpectus, tutior in infuſo vel decocto ʒj.

Datur trochiſcatus ad ʒß. in Extr. ℈j. ut & reſina ad gr. xv. Corrigitur vis tormentoſa Zinzibere ºoo caryoph.

caryoph. & macis. Specificatur variis Morbo appropriatis.

℞. Agar. troch. ℈j diagr. gr. v. al. ſuccotr. ℈ß ☍ cinam. gt. j. Elix. propr. q. ſ. f. pil. purgantes.

℞. Agaric. troch. ʒß. Reſin. Jal. gr. v. ☿ dulc. gr. x. aloës ſuccotr. gr. xv. ☍ d. ſuccin. gt. ij. majoran. gt. j. Extr. Caſtor. q. ſ. f. pil. Catarrhales.

℞. Reſin. agaric ℈ß. ☿ dulc. gr. vij diagryd. gr. iij M. f. pil. purgantes.

FUNGUS ORBICULARIS SEU BOVISTA.

Eſt ſubſtantia fungoſa, innaſcens ſuperficiei terræ, levis, aliquando pugnum magnitudine excedens, ſub tenui coriaceo ex cinereo livido inveſtimento includens medullam levem, raram, ſpongioſam, fuliginoſam, lividam, quæ compreſſa pulverem profundit ſubtiliſſimum: odoris & ſaporis nullius evidentis niſi terreſtris & paululum adſtringentis.

Principia activa ſunt particulæ terreſtres. Interne ob metum veneni non exhibetur. Externe adſtringit, incraſſat, abſorbet humida, unde in hæmorrhagiis, intertrigine, puſtulis &c. uſurpatur.

℞. Terr. ⊕l. dulc. phlegm. ⊕li boviſt. ana. q. ſ. f. maſſa in venis apertis, extemplo ſiſtens fluxum ſanguinis.

CAPUT QUINTUM.
DE FRUCTIBUS ET SEMINIBUS.
SALE VOLATILI OLEOSO AROMATICO CONSTANTIA.

Caryophylli aromatici.

ANTHOPHYLLI.

SUnt arboris caryophylloferæ Tſinka dictæ fructus immaturi oblongi, inſtar clavi, anguloſi, depreſſi, rugoſi, ex rubigineo nigricantes, quatuor ſtellatim decuſſatis apiculis notati, cum medio capitello convexo & cavo, futuri floris rudimento.

Sunt fructus immutari, Anthophylli vero maturi, caryophyllis craſſiores & tumidiores, includentes ſub cortice duriuſculo ſemina oblonga, fuſca, ſaporis & odoris ejuſdem cum Caryophyllis. Solent in arbore relinqui, ut ex illis deciduis nova arbuſcula iterum enaſcantur.

Fructus ſunt primo viridis coloris, poſt rubeſcunt, ſiccantur, & ad nos deferuntur.

Caryophylli præſtantiores Anthophyllis, gaudent ſale volatili admodum penetrante, ſubtili & oleoſo, ſpiritibus noſtris adeo analogo, ut vel hos torpentes, vel craſſis viſcidiſve humoribus involutos exſuſcicitent, vel plane proſtratos regenerent. Unde in

Morbis

Morbis ſoporoſis, Apoplexia, Paralyſi, Catarrhis præ cæteris palmam merentur: eandem ob cauſam in primarum viarum muco incidendo, à quo vomitus aliaque mala oriuntur, nulli facile cedunt. Purgantibus tanquam corrigens junguntur, & ſingulare quid obtinent in frigore febrium intermittentium.

Dantur in ſubſtantia ad gr. vij. in infuſ. ad ʒß 🝆 deſt. ad gt. j. ingreditur linimenta & unguenta ad Paralyſin, & Balſamum Apoplecticum.

℞. Caryoph. ℈ß. Cinam. ℈j Nuc. Moſch. gr. xv. M. f. p. pro ij doſibus.

℞. 🝆 Caryoph. origan. ana gt. iij vel iiij camphor. gr. v. excip. goſſyp. & intrudatur cavo dentis ſ. excipiatur opio & f. turunda cavo dentis intrudenda.

Caryophyllis in Electuarium vel cum melle vel ſyrupo redactis, ſtrictiſſime obſervata diæta Luem Veneream curavit D. Hermannus, ſudorem valdopere movet.

AMOMUM.

Sunt arboris Americanæ Caninga dictæ, baccæ rotundæ ex fuſco nigricantes, piperis magnitudine, in ſummo umbilicatæ, includentes ſub tenui pellicula & ſubſtantia fungoſa, binos acinos nigros parvos, tenui membrana veſtitos, ſaporis & odoris ad caryophyllos accedentis.

Sunt fructus caſſiæ caryophyllatæ.

Caryophyllis aromaticis accedunt & reſpondent, ſed virtutibus ſunt debiliores, in ſpecie conferunt in vertigine.

℞. Cort. Winteran. ʒj. caryoph. arom. ʒij amom. ʒiij ſaſſafr. ʒj bacc. Junip. ʒij ſumm. abſynth. Mj. f. infuſ. antihydropicum.

℞. Amom. ℥ß. galang. ʒj rad. doronic. ʒiß. M. f. p. vel addita conſerva florum tunicæ f. bolus contra vertiginem, vel infundatur vino & fiat infuſum.

MACIS.

Eſt Nucis Moſchatæ cuticula membranacea, craſſiuſcula, ramoſa, quaſique reticulata, ex aureo flaveſcens, incumbens putamini corticoſo, cui medulla Nucis moſchatæ incluſa eſt.

Macis ⊖ aromaticum oleo ſociatum paulo temperatius eſt quam caryophyllorum, & ſuas operationes quoque blanda vi anodyna abſolvit.

In colica, Vomitu & pellendis Menſibus ſpecifice commendatur. Datur in Subſtantia ad gr. x. in infuſ. ad ʒij. In India pro pellendis menſibus adhibent macin ex infuſo ▽æ ferventis vel cum floribus lavendulæ. Vis anodyna intenditur Nuce Moſchata, ℴ℘que ejus expreſſo.

℞. Macis. ʒiß infund. vin. ad tormina abdominis, add. Nuc. Moſch. ʒß. contra colicam.

℞. Mac. flaved. Cort. aurant. ana ℈ß. Ocul 69. ℈j f. pulv. ad colicam add. Laud. op. gr. j.

℞. Theriac. ʒß. ℴ℘ d. mac. gt. ij. ſ. macis ℈ß L. O. g. j. ſucc. cydon. q. ſ. f. bolus ad Vomitum ſiſtendum.

℞. ℴ℘ mac. expr. ʒß. Zibeth. gr. viij. f. linimentum in colica à debilitate ventriculi, regioni umbilicali applicandum, addi poteſt ℴ℘ deſt. macis.

NUX MOSCHATA.

Eſt arboris nuciferæ aromaticæ Pala dictæ, nucleus rotundus aut oblongus, ſolidus exterius quodammodo rugoſus & cinereus, intus pallide flaveſcens, venis albicantibus hinc inde interpolatis, ſaporis acriuſculi aroma-

aromatici, ſimulque pinguis, odoris fragrantis ſatiſque grati.

In Moluccis creſcunt non ſua ſponte ſed cultura, arbores magnitudine accedunt ad mala vel pruna, flores habent pentapetalos cum nullis ſtaminibus.

Maci reſpondet, ſed temperatior eſt, & ob terreas adſtringentes partes & 🝆 craſſum habet vim potentiorem anodynam, ad indolem opii quodammodo accedentem, hinc anodyna eſt & hypnotica, ſiſtens nimium ſpirituum & ſanguinis motum, & in vomitu, colica, dyſpnoea, ſtranguria, elegans ſpecificum eſt. In cardialgia, Paralyſi, & ab hac læſa deglutitione ſingularem fert opem.

Stomacho applicata externe eum mire roborat. Datur ut macis, ſed cave ne à nimia doſi ob olei copiam Ventriculi fermentum obtundatur, neve torpor & ſtupor ſpiritibus inducatur. Toſta & 🜔ibus aromaticis orbata, tota terrea, adſtrictoria, anodyna & hypnotica eſt. Et in quolibet fluxu compeſcendo auxilium fert.

Oleum ſuppeditat triplex. 1. 🝆 deſtillatum tenue & ſubtile 2. expreſſum, craſſum & ſebaceum, in quo vis aromatica adhuc integra hæret. 3. coctum, itidem craſſum & ſebaceum, aſt virtute aromatica plurima ex parte orbatum. Hoc colligitur ex reſiduo deſtillationis Nucis Moſchatæ, cujus ſuperficiei inſtar ſebi ſupernatat.

℞. Nuc. Moſch. toſt. gr. xv. Corall. r. præpar. C.C. uſt. ana ℈j Laud. op. gr. ij. M. f. pulv. vel cum conſerv. roſ. ⊕lat. f. bolus ad quemvis fluxum alvi.

℞. Vitell. ovor. q. v. Nuc. Moſch. ℈j + roſac. par. coque ſimul ad duritiem, ad quoſvis alvi fluxus.

℞. ⚬⚬⚬ Nuc. expr. gr. x. L. O. gr. ij. conſ. roſ. ⊕lat ʒß. f. bolus in alvi fluxu, adde ſi velis, Corall. præpar. ſ. C.C. præpar. ut & ſacch. ♄ni gr. vij.

℞. ⚬⚬⚬ N. M. ex deſtillat. reſid. ℈j Corall. r. præpar. C.C. præpar. ana gr. xv. L.O. gr. ij Conſ. roſ. ⊕l. & ſyr. nonnihil vel q. ſ. f. bolus præſtantiſſimus in omnibus fluxibus.

℞. ⚬⚬⚬ Nuciſt. expr. & multiplici affuſione ♒us vini odore aromatico orbati ʒj ⚬⚬⚬ ſuccin. gt. v. caryoph. gt. iij ⚬⚬⚬ expreſſ. ex flaved. cort. aurant. gt. viij cinam. gt. ij. M. f. Balſamus apoplecticus vel analepticus in quovis caſu ex torpore ſpirituum.

℞. ⚬⚬⚬ Nuciſt. expr. ʒj ⚬⚬⚬ ſuccin. gt. x. ſpicæ gt. xv. camph. gr. vij f. linim. antiparalyticum. Hujus grana iij vel iiij poſſunt exhiberi ad convulſiones.

℞. ⚬⚬⚬ Nuc. Moſch. expr. ʒij rad. ir. flor. ʒiß. Croc. or. ℈ß. L.O. gr. xv. ſtyrac. benz. ana ʒj Balſ. Peruv. q. ſ. M. f. pomum hypnoticum, naribus admovendum, ad conciliandum ſomnum.

CUBEBÆ.

Sunt fruticis convolvulacei Javanenſis Curanc dicti, rotunda piperi ſimilia, paulo majora, fuſca, rugoſa ſcabra, cum appenſo tenui pediculo caudæ inſtar, includentia ſub cortice tenui nucleum parvum rotundum, foris nigricantem, intus albicantem, ſaporis aromatici potenter acris, odoris ſuavis.

Adeo penetrante acri & aromatico ⊖e pollent, ut in ore volutatæ faucium & linguæ poros aperire & quaſi diſſolvere videantur. Hinc quoque in Catarrhis, glandularum obſtructionibus, raucedine, Paralyſi, Apoplexia maximam habent virtutem, confe-

runt

runt quoque ad confortandam memoriam, & ſtimulandam ♀em, in ſiccitate oris à ſpirituum animalium denegato influxu, ſingulare quid promittunt.

Dantur in ſubſtantia à gr. v. ad viij in infuſo ad ʒij ☍ deſt. ad gt. ij ad memoriam confortandam.

℞. Cubeb. ʒij ſacch. penid. ℥j infund. ▽ puleg. q. ſ. f. infuſ. ſ. coque ſaccharum ad conſiſtentiam & fac rotulas contra raucedinem.

℞. Cubeb. gr. vij. caric. ping. ʒj cape ſingulis diebus ad præcavendam Apoplexiam.

℞. Cubeb. ℈ß. Caſtor. gr. v. ſacch. par. M. f. pulv. aut fac trochiſcos ſeu rotulas ad nimiam ſiccitatem oris, ortam à ⁓uum animalium denegato influxu.

CARDAMOMUM.

Cardamomum majus non eſt in uſu, nam non fert ætatem ut minus. Minus ſunt plantæ indicæ capſulæ membranaceæ triquetræ, palleſcentes, tribus per longitudinem diſtinctæ cellulis, quibus includuntur acini anguloſi ex fuſco nigricantes, involuti tenui & albicante pellicula, ſaporis acris aromatici, ejuſdemque odoris.

Majus ſunt plantæ indicæ Benzal dictæ, capſulæ membraneæ triquetræ, cardamomo minori triplo longiores, tribus per longitudinem diſtinctæ cellulis, ſaporis & odoris minori fragrantioris.

Minus majori præſtantius. Vi aromatica gaudet remiſſiore, & minus deſiccativa, quam cubebæ, attenuat ſatis & penetrat, ut in vertigine, confortanda memoria cubebas ſuperet.

Datur in ſubſtantia ℈ß in infuſo ut cubebæ, ſtimulatur camphora.

℞. Car-

℞. Cardamom. ʒß. Coriandr. ʒij ſinap. ℥ß. M. f. p. pro Apoplexiæ præſervatione.

℞. Ambr. gryſ. gr. xx. ſolv. in 🜅 cinam. cubeb. cardamom. ſ. q. ſacch. albiſſ. ℥iiij Moſch. or. gr. iij M. f. p. ad confortandam memoriam.

PIPER ALBUM ET NIGRUM.

Piper album & longum non differunt ratione plantæ, ſed ratione fructus. Datur etiam factitium. [Quod ita deſcribit Sam. Dale Pharmacol. p. m. 446. Javanenſes & Amſtelodamenſes immittunt piper nigrum in ▽ ut decorticetur, & cum involvunt in Creta, ut fiat piper album factitium.]

Nigrum ſunt fruticis ſarmentoſi piperiferi grana rotunda, piſi magnitudine, globoſa, nigricantia, rugoſa, includentia ſub cortice intus albicante nucleum nullum vel exiguum ſaltem. Saporis acris & fervidi aromatici, odoris fragrantis.

Album ſunt ejuſdem grana, cortice cinereo tecta, pellicula minus rugoſa.

Album ſi naturale ſit, non factitium, gratius eſt & laudibilius nigro, utrumque gaudet ⊖e aromatico acerrimo pungente, fixo potius quam volatili, cum pauco oleo ſubtili, acredine piperata fere privato, quod mirandum.

Ob ſalis penetrantiam, viſcida & acida potenter penetrantis ſtomachium eſt & anticolicum, leniens ſimul horripilationes febrium intermittentium, ipſarumque febrium paroxyſmos, præcipue ſi fomes earum ſit in Ventriculo.

Externe reſolvit nervos obſtructos, uvulam elongatam & à Pituita obſeſſam curat. Integrum ſeu groſſo

groſſo modo contuſum felicius datur, quam minutius tritum, hoc enim membranas Ventriculi & inteſtinorum aliquando inflammat, illud vero aromaticam ſuam virtutem tranſmittit in ſanguinem [vid. hac de re Caſp. Hofmann. de Medic. officinal. l. 2. cap. 175. p.m. 424.]

Decocta piperum, quia ⊖e fixo gaudent, acerrima fiunt.

In ſubſtantia dantur gr. iij. in infuſo datur ad ℈ß. nimia acredo temperatur oleoſis, præcipue baccis lauri, ſaccharo & fœniculo.

℞. Piper. gr. iij integra cape ad cruditates Ventriculi ex paſſione colica, vel contund. & add. conſerv. menthæ.

℞. Piper integr. gr. iij cap. ex cochl. j. ♒us vini in febribus ante paroxyſmum.

℞. Piper. gr. iij bacc. laur. ℈ß. L.O. gr. j. M. f. p. ad febres.

℞. Piper. gr. vij O gr. v. f. pulv. uvulæ applicandus.

℞. ☍ piper. deſt. gt. v. Extr. gentian. abſynth. ana ℈j f. bolus ad febres intermittentes præcipue quartanam; ut & illarum horripilationem.

℞. ☍ laurin. expr. ʒij ☍ piper. deſt. ℈j ☍ nardin. ʒj f. linimentum dorſo applicandum, contra horrores febriles.

N. quod fructus piperis Hiſpanici recentioris fumum ſuppetant, qui attractus tuſſim excitat vehementem, adeo ut omnes qui adſunt in conclavi tuſſire cogantur. Ope hujus tuſſis poteſt rejici, ſi quid incidit in aſperam arteriam.

PIPER LONGUM.

Piper longum ſunt ſarmentoſi fruticis fructus oblongi, teretes, cinericei, [inſtar nucamentorum ſeu Julorum Coryli,] ex numeroſis vermiculato plexu diſpoſitis granis minutis compacti, ſaporis pipere nigro acrioris [& amaricantis.]

Piper hoc rotundo quidem ingratius, ſed virtute potentius, & proinde ad excitandos ſoporoſos, emendandas Ventriculi cruditates uſus elegantioris eſt, quam piper rotundum.

℞. Piper. Caſtor. rad. pyrethr. ari, fol. origan. ana q. v. f. infuſ. ex vino vel ♒u vini adde ſi vis Nuc. Moſchat. [In eſſentia ſtomachica Thomſoni, in cruditate acida & viſcida commendatiſſima piper baſin conſtituit.]

℞. Piper. caſtor. helleb. alb. ana ℈j ꝏ rut. major. ſalv. ana gt. ij. M. f. pulv. ſternutatorius ad excitandos ſoporoſos.

GRANA PARADISI.

Sunt ſpecies Cardamomi, arilli anguloſi, nitidi, fuſci, fragmentis lapillorum granatorum ſimiles: utuntur Indi loco piperis [ſaporis acris & fervidi medii inter cardamomum & piper.]

Virtute cum pipere coincidunt, ejuſque defectum ſupplent, tam condimenti quam medicamenti loco. Uſurpatur ut piper, idcirco huic admiſcetur, & iis adulteratur. Præbent ꝏ acerrimum.

℞. ꝏ laurin. ʒj ꝏ gran. parad. gt. viij in Ventriculi debilitate.

Capiuntur ad Epithemata contra Paralyſin.

Infuſa

Infuſum ex vino in Ventriculi debilitate ejuſdem eſt facultatis cum infuſo acori.

SEMEN AMMIOS.

Sunt Ammios majoris C. B. P. ſemina parva, petroſelino minora, ex cinereo ſpadicea, ſaporis amari, acris, odoris fragrantis.

Minoris ſemine apii, odore origani. J. B. ſunt ſemina parva, ſtriata, petroſelino minora, ex rufo ſpadicea, ſaporis acris aromatici, quaſi ex pipere & cumino compoſiti, odoris fragrantis ad origanum accedentis.

Minus majori præſtantius & præter virtutes aromaticis affines vim habet ſpecificam anticolicam, ex plurimorum ſententia vim ad promovendam fœminarum fœcunditatem. [vid. hac de re Matthiol. lib. 3. in Dioſc. c. 61. p. m. 557.] ob ſalis aromatici & ~uoſi penetrantiam, quod ſe ad Uterum extendit, etiam fluori albo conducit, unde quoque ingreditur Clyſteres Uterinos.

Majus datur ad ʒß minus ad ℈j.

℞. Sem. ammios ʒß fol. horm. p. j. lign. ſaſſafr. ʒj affund. ▽ cinam. & fac infuſum, quod cape cum toto. In fluore albo.

℞. Sem ammi. ʒß. ſacch. cand. par. f. pulv. ad ij vel iij doſ. exhibeatur infantibus contra tormina.

℞. Ariſtol. rot. ℥ß rad. Zed. ʒj Cent. m. roriſmar. ana Mj. chamom. artemiſ. ana Mß. coq. in ▽ f. Clyſter Uterinus.

Indi exhibent à ʒj ad ʒij contra ſterilitatem mulierum, ſæpe cum fructu, ubi ſcilicet adeſt, obſtructio vaſorum.

R 2 ℞. Ci-

℞. Cinam. ℥j caryoph. ʒij Nuc. Moſch. piper. galang. ana ℥ß ſem. ammios ʒiij affund. 🜄 regin. Hung. datur infuſum cochleatim contra ſterilitatem.

SEMEN ANISI.

Sunt Aniſi herbariis dicti C.B.P. ſemina oblonga, rotunda, in obtuſum acumen deſinentia, parva, ſtriata, colore ex rufo ſpadiceo.

Duplex eſt ſemine majore & minore, ſed ſolent confundi.

Semina conſtant 🜔e volatili 🝆ſo dulcibus incluſis, ſive ſalibus aromaticis dulcedine temperatis gaudent, unde in raucedine aſperitate faucium, tuſſi, colico dolore, & promovenda Urina conducunt, ſiſtunt ſingultum, & ſimul laxant alvum tardiorem puerorum & duriorem; impediunt quoque lactis coagulationem, corrigendo acidum & amandando illud ad Urinam. Datur à ʒß ad ℈ij.

℞. Sem. aniſ. ʒiß coq. in 🜄 ſ. vino ad paſſionem colicam.

℞. Sem. aniſ. 🜔 geniſt. ana ℈j f. pulv. diureticus.

℞. Rhabarb. ʒj ſem. aniſ. ℈j f. pulv. ad alvum laxandam.

℞. Flor. benz. ʒj 🜔 volatil. C.C. 🝆 d. aniſ. gt. ij. ſacchar. q. ſ. in tuſſi.

℞. Sem. aniſ. ℈j 🝆 d. aniſ. gt. ij rad. alth. ʒj ſacch. q. ſ. f. pulv. ad raucedinem aſperitatem & tuſſim.

Oleum aniſi deſtillatum blandum anodynum eſt & alvum ſimul laxat, in colica & fluxu albo ſpecifice laudatur. Externe ſuffuſionibus, coryzæ, cardialgiæ, inflationibus abdominis & hypochondriorum; ſingultibus medetur.

SEMEN

SEMEN ANISI CHINENSIS.

Aniſum peregrinum C. B. P. aniſum Philippinarum inſularum. Aniſum ſtellatum, fœniculum Sinenſe. Semen Badianum. Zingi etiam vocatur.

In Moſcoviam transfertur ex Tartaria. Semine ſunt abſque omni odore & ſapore. In involucris conſiſtit omnis vis aniſata, in ſemine nihil.

Oleum hoc nunquam concreſcit, ut noſtrum ☍ aniſi.

Convenit virtute cum aniſo vulgari, ſed ob dulcedinem gratiorem & fere penetrantiorem vim aromaticam, quæ in ſeminum involucro conſiſtit habetur præſtantius. Creditur vulgo confortans & roborans catholicum, Magnates volvunt ſub lingua ad conciliandam gratiam ori, ad arcendam nauſeam & faſtidium, ut & tollendum aëris fœtidioris incommodum. Chinenſes infundunt cum radice Ninzin, ſeu maſticant in ore & ſuperbibunt potum Thee, horum exemplo noſtrates utuntur hoc ſemine cum potu Caffee.

SEMEN FOENICULI.

Sunt fœniculi vulgaris Germanici C.B.P. ſemina fœniculo dulci ſimilia, ſed longiora, odore & ſapore debiliori. Dulcis ſeu Italici ſunt ſemina excinereo ſpadicea, oblonga, ſtriata, vulgari fœniculo rotundiora, ſaporis dulcis, acris & ſuavis, odoris fragrantis.

Fœniculum dulce, quod optimum, convenit cum ſemine aniſi & propria radice, qua tamen potentius eſt, in Variolis & Morbillis, ſtranguria, calculo, augendo lacte in frequentiori uſu eſt. Externe medetur oculorum aciei.

Siccum præfertur viridi quod ob excrementitiam humiditatem torminoſum eſt, datur ut ſemen aniſi

℞. Sem. fœnic. ℈j aniſ. ℈ß 🜕 deſt. fœnic. gt. j. Sacchar. par. f. pulv. contra tormina.

℞. Rad. helen. ℥ij ſem. fœnicul. ℥ß. paſſul. min. ℥iß. M. f. decoct. in aſthmate & tuſſi. vid. plura d. radice fœnic. p. 33.

℞. Olei fœnicul. ⏜ ⊖✳ci M. invicem, inungantur hypochondria contra inflationes. Infantibus obeſis & robuſtis hæc mixtura valde convenit, contra qui non ſunt ſat robuſti ferre non poſſunt ob ⏜um ⊖✳ci.

℞. ▽ fœnic. Cent. min. ana ℥ij ſyr. C. B. ℥j ⊖ C. C. gr. v. Tinct. Caſtor. ʒß ut & ſi libuerit. Tinct. Myrrh. ℈ß. f. mixtura contra Variolas & Morbillos.

℞. ▽ fœnicul. ℥iiij ⊖ ♀ri gr. vij ad ℈ß fol. ſenn. ʒvj f. infuſ. in loco calido, ſi lubet add. ſacch. cand. par. ſi fortius expetatur add. rhabarb. ʒij.

℞. ▽ fœnicul. Euphraſ. ana q. v. add. fol. ſenn. vitr. ♁ij vel croc. metallor. pone in loco calido utere ad Inflammationem, aliaque vitia Oculorum.

SEMINA LEVISTICI.

Sunt Liguſtici vulgaris C. B. P. ſemina oblonga, depreſſa, membranacea, ſtriata, ſaporis ſubdulcis, aromatici nauſeoſi.

Conf. radix Leviſtici p. 15. cum qua ſemina conveniunt.

℞. Sem.

℞. Sem. Leviſtic. ʒß. Caſtor. gr. x. Croc. or. gr. vij L. O. gr. j. M. f. pulv. pro ij doſ. contra dolores & tormina colica.

***

SEMINA SALE VOLATILI AROMATICO CUM BLANDA VISCOSITATE CONSTANTIA.

SEMEN ANETHI.

Sunt anethi hortenſis C. B. P. ſemina foliacea & membranacea, orbicularia, compreſſa, in parte exteriore tribus ſtriis vel ſulcis notata ex atro virentia.

In ⊖e volatili temperato quaſique incluſo, fovet quippiam anodyni & narcotici, & propterea in colica, ſingultu, cardialgia & ileo ſpecifice commendatur. Semina recentia contuſa & temporibus applicata puerorum, placidum ſomnum accerſunt.

℞. Sem. aneth. ℈ß. Caſtor. gr. iij L. O. gr. j. M. f. pulv. ad ſingultum.

℞. Summit. aneth. Mj. coq. ex ▽ vel Cereviſ. ſit potus ordinarius in ſingultu.

℞. Summit. aneth. q. v. coq. ex ▽ papav. ſeu adde capita paveris albi pro potu, ad ſomnum conciliandum.

℞. Olei d. aneth. gt. v. ☍ amygd. dulc. rec. expr. ℥ß. f. hauſtus ad ſingultum.

℞. Olei d. aneth. gt. iij C. C. præpar. ℈j L. O. gr. j. Cort. aurant. condit. cum proprio ſyrupo q. ſ. f. conditum contra tormina.

℞. Olei

℞. Olei Papav. expr. ℥ß 🜄 aneth. gt. xv. ung. popul. ℥ij M. f. liniment. add. L. O. ℈ß. ad ſomnum conciliandum.

SEMINA CUMINI ET CARVI.

Sunt Cumini ſemine longiore C. B. P. ſemina ſtriatata, oblonga, ex cinereo ſpadicea, fœniculi ſeminibus longiora & craſſiora. Sapore acri aromatico ingrato. Carvi ſunt Cumini pratenſis ſeu Carvi officinarum C. B. P. ſemina ſtriata, parva, oblonga, tenuia, paululum incurva ſeu falcata, fuſca. Saporis acris aromatici, odoris fragrantis. Hoc tetrum illud redolens cumini non habet,

Sale aromatico oleoſo temperato admodum anodyno & carminante gaudent, & in tympanitide, flatulentia ventriculi, colica flatulenta tam interne quam externe laudantur.

Semen Carvi gratius eſt, quam Cumini. Hinc ſemen carvi felicius interne, cumini vero externe præſcribitur.

℞. Sem. carv. ſaccharo incruſt. q. v. datur ad flatulentiam ventriculi, add. Oleum proprium deſtillatum pro ſtimulo.

℞. Emplaſtr. d. cumin. malax. in ſ. q. Olei cumin. & carvi. in Doloribus iſchiaticis, in inflationibus ventriculi & abdominis.

℞. Semin. cumin. & carv. q. v. contunde, ſacculo involve, & applica abdomini contra tormina.

℞. Olei carv. deſt. gt. iiij ♒ Menth. ℥iij vel iiij L. O. gr. i. f. mixtura in cardialgia.

BACCÆ

BACCÆ JUNIPERI.

Sunt Juniperi vulgaris fruticoſæ C.B.P. baccæ rotundæ per maturitatem nigricantes, includentes ſub tenui pellicula pulpam, acinos tres oblongos triangulos, ſaporis ſubacris ſubdulcis & reſinoſi, odoris fragrantis. Vis conſiſtit in pulpa interiore.

Harum ⊖ aromaticum oleoſum & quodammodo Therebinthinatum, cœteris excellit in promovenda Urina, ſudore, Ventriculi flatulentia, in menſium obſtructione, præcavendo calculo, & ad reſolvendas Catarrhoſas obſtructiones. Hydropicis ſingulariter conveniunt ob virtutem ſtomachicam & diureticam. Externe laudantur in ſuffitu ad Catarrhos & membra Paralytica. Dantur in ſubſtantia ad ʒj.

℞. Bacc. Junip. ℈j ꝏ d. junip. gt. iiij rad. rub. tinct. gr. xv. f. p. capiendus ex convenienti liquore ex ▽ gramin. vel ▽ al. deſt. propria, pro diureſi excitanda.

℞. Rad. Angelic. Leviſtic. ana ℥ß. roriſmar. Matricar. ſabin. ana Mj. bacc. Junip. ℥iß. Cinam. ʒiij Croc. or. gr. xv. vel ℈j. ⊖ 🜿ri ℈j affund. Vin. q. ſ. ad menſium obſtructionem chronicam, adde ſi menſes inſtant Rhabarb. ʒij hell. n. ʒj f. infuſum.

℞. Roob. Junip. ℥iiij rhabarb. ʒiij Cinam ʒj f. Elect. diureticum, hydropicis convenientiſſimum add. roob. ſambuci.

℞. Spiritus Junip. ℥iiij balſ. Peruv. ʒj f. lavamentum ad dolores partium à Catarrhis ortos.

Olei deſtillati guttulæ aliquot pulveribus antifebrilibus adduntur.

℞. Olei Junip. gt.iij. Ocul. 69.℈j. ⊖ prunell. gr.xv. ſacch. ♄n. ℈ß. L. O. gr. ij f. p. contra febres intermittentes.

℞. Vin. Junip. ℥iiij. Tinct. Nephritic. ℥j ♒. Carmi. Sylvii ℥ß. f. mixtura diuretica.

Vinum Juniperinum paratur ſi muſto baccæ ejus contuſæ injiciantur & fermentent ſimul.

BACCÆ LAURI.

Sunt Lauri vulgaris C.B.P. baccæ oblongæ, rotundæ, nigricantes, ſub pellicula tenui includentes nucleum geminum nigricantem, ſaporis acris amaricantis aromatici oleoſi, odoris non ingrati.

Blandum ſuum oleoſum aromaticum ⊖ fovent in ☍ craſſo admodum, anodyno & hypnotico, virtute diuretica & Uterina conſpirant cum baccis Juniperi, his tamen virtute carminativa & anodyna longe ſuperiores ſunt.

Hinc ſpecifice prædicantur in Colica, Paralyſi, doloribus & Uteri vitiis, in febribus malignis ſingulare & blandum ſudoriferum conſtituunt, præcipue ſi camphora addatur. Externe applicatæ conveniunt in inflatione ventriculi, paralyſi, convulſione, ileo, iſchuria, doloribus poſt partum, colicis doloribus, item in enecandis pediculis, nam odorem non ferunt, in ſananda ſcabie & maculis faciei.

Cave ne nimia quantitate exhibeantur, quia ob ſui oleoſitatem facile Vomitum cauſant.

℞. Raſ. lign. ſaſſafr. ℥ij. roriſmar. Mj. bacc. laur. ℥j coq. in ▽ & vino ad Paralyſin.

℞. Bez. min. gr.xv. ⊖ volatil. C. C. gr. v. vel vij bacc. laur. ℈ß. f. p. quietem concilians in doloribus.

℞. Flor.

℞. Flor. chamom. Mij. bacc. laur. ℥ij f. decoct. vaſe clauſo leni △e, adde colaturæ ☍ expr. bacc. laur. ʒj f. Clyſter in colica optimus.

℞. Bacc. laur. cort. aurant ana q.v. infunde vin. q. ſ. exprime fortiter, expreſſum exhibe in colica paſſione.

℞. Flor. ♃is, ♁ diaphoret. bacc.laur. ana ʒß. M.f.p. diaphoreticus de die uſurpandus in febribus malignis, addi poſſunt Camphoræ gr. vij.

℞. Bacc. laur. ℈j. borrac. gr.vj. cinam.ac. gr.xv. ☍ Mac. gt. ij M. f. p. in menſium remora cum dolore.

℞. Flor. Cham. Mvj. fol. Malv. Mij. flor. ſambuc. Mj. bacc. laur. ℥j coq c. ▽ cui inſideat iliacus.

℞. Olei bacc. laur. expr. ℈ß, (ſi pluſcula doſis exhibetur Vomitus ſequitur ob oleoſitatem, non tamen moleſtus vel noxius) ☍ ſpicæ. thereb. ana gt.v. ☍ ſuccin. gt. j. camph. g. j. ☍ N. M. expr. gr. v. M.f. Balſamum pro ij vel iij doſibus, ex Cereviſia calida exhibent, idem balſamum applica partibus convulſis, præcipue infantum externe.

℞. Olei laurin. expr. ℥ß. Caſtor. ʒj. ☍ deſt. chamom. ℈ß f. liniment. ad dolorem iliacum, colicum alioſve.

℞. Olei expr. bacc. laur. ℥j. ung. Mart. ℥ij & f. ung. quo inungatur regio pubis infantum & ſubito fluet Urina ſuppreſſa.

℞. Olei expr. bacc. laur. ℥ij flor. ♃is. ʒij ſacch. ♄ni. ʒj. Camph. ℈ij f. liniment. ad ſcabiem: addi poſſunt ☿ dulc. ʒij.

Elect. ex bacc. laur. ad ʒj datur cum camph. gt. v. & ⊖is vol. ſuccin. gr. xv. pro ij vicibus, conſtituit bolum ſudoriferum, anodynum & carminativum.

℞. Olei d. bacc. laur. gt. iij. L. O. g. j. cran. human. ʒß. M. f. p. anodynus in colica ab acido.

CARPOBALSAMUM.

Sunt Balſami Ægyptiaci Proſp. Alpini fructus oblongo rotundi, rugoſi, piſo minores, ſpadiceo fuſci in parvum aculeum deſinentes, includentes aliquando dum matureſcunt exiguam medullam balſamicam oleoſam albicantem, odoris grati, aliquando nullam medullam includunt, ſed putamen tenerum eſt incaſſum.

[Conf. Proſp. Alpin. dialog. d. Balſamo cap. 6. p. m. 75.]

Naſcitur nullibi ſua ſponte, colitur in hortis magnatum, habet folia inſtar liguſtri. Conveniunt cum balſamo Ægyptiaco, quod vid. ſuo loco. Vis conſiſtit in reſina liquida ſeu ſucco balſamico fructuum cavitati incluſo.

℞. Spec. diambr. ℥ß. lign. aloës, carpobalſami ana ʒij affund. -∽- roſar. ardent. q. ſ. f. eſſentia, ejus gt. xx. ex ſyr. chermes pro cardiaco exhiberi poſſunt.

Commendatur in India ad gonorrhæam ſiſtendam. ℞. Carpobalſ. ℈j oſſ. ſep. rhab. toſt. ana ℈j M.

℞. Carpobalſ. ℥j affund. ꝏ Hyperic. ſuccin. Junip. vel Therebinth. f. balſamum Vulnerarium internum & externum.

SEMINA DAUCI.

Sunt vel Cretici vel vulgaris. Cretici ſunt Dauci foliis fœniculi tenuiſſimis C. B. P. ſemina oblonga, incana, ſuperiori in parte acutiora, villis reſtricta, ſaporis acris aromatici grati, odoris debilis.

Vulga-

Vulgaris ſunt paſtinacæ tenui foliæ ſylv. Dioſc. ſeu Dauci officinarum C.B.P. ſemina compreſſa, cinerea, Dauco Cretico breviora, latiora, circa marginem utrinque villo obſita, ſaporis dauco cretico debilioris, odoris fragrantis.

Utriuſque dauci ſemina ob ⊖ blandum ☍ſum carminativum conveniunt in calculo, nephritide, menſium ſuppreſſione, doloribus poſt partum ut & in ſtranguria.

℞. Sem. dauci ʒij infunde in vino albo & exhibe hyſtericæ.

℞. Bacc. Junip. laur. ſem. dauc. ana ʒiij coq. in Cereviſia ad nephritidem.

℞. Fol. ribes. nigr. Mj. ſem. dauc. ℥ß coq. ex vino rubro contra ſtranguriam.

[Singulariter contra calculum commendat ſemen dauci Helmont. de Lithiaſi c. 1. §. 19. p.m. 14. it. cap. 7. §. 14. p.m. 39. it. cap. 8. §. 23. p. 48.]

SEMEN SESELIOS ET SILERIS.

Sunt Liguſtici, quod Seſeli officinarum C.B.P. ſemina oblonga, ſtriata, ex viridi ſpadicea, notata per marginem fimbria exigua folioſa, ſaporis acris amaricantis aromatici, quaſi ex corticibus aurantiorum & cumino mixti.

Vulgaris ſunt ſeſelios pratenſis C. B. P. ſemina oblonga ſtriata, fœniculo proxima.

Utraque cum Dauci ſemine conſpirant, ſed virtutis ſunt remiſſioris, interim in diſcutiendis flatibus & menſibus commendantur.

℞. Sem. ſeſel. dauci ſiler. ana ʒiij. flav. Cort. aurant. Citr. ana ℥j infund. vino ſeu ▽ gramin. ⊖e geniſtæ impregnatæ f. infuſum diureticum.

SEMEN PETROSELINI.

Sunt apii hortenſis Petroſelini vulgo C.B.P. Semina parva, oblonga, ſtriata, atrovirentia, ſaporis acris. Symbolizant cum radice propria, ſed ſunt præſtantiora.

℞. Sem. petroſel. ℥ß. cape ex decocto propriæ radicis vel affunde ▽ gramin. ⊖e geniſt. imprægn. & pone in digeſt.

SEMEN APII.

Sunt Apii paluſtris & Apii officinarum C.B.P. ſemina parva ſtriata ex fuſco ſpadicea.

Convenit cum ſemine petroſelini & propria radice, qua tamen potentius eſt, præcipue in educendo calculo & Urina, commendatur una cum radice in Uteri affectibus.

℞. Sem. apii ℥ß. ⊖ vol. C.C. gr. v. cape ex ▽ chelid. maj. vel chærefol. contra calculum vel icterum.

SEMEN RUTÆ.

Sunt rutæ hortenſis latifoliæ C.B.P. ſemina parva, recurva, ſemilunata, coloris modo glauci, modo nigricantis, ſaporis ingrati, odoris gravis. In capſulis ſeminalibus omnis vis, ex illis enim plus olei extrahitur quam ex integris ſeminibus & caulibus rutæ.

De rutæ inſigni virtute quæ in ſeminali capſula concentrata v. cap. d. herb. rutæ. Semen commendatur inprimis ad lumbricos, gonorrhæam & extinguendum ſemen ob nimiam ſui ſiccitatem.

℞. Sem. rut. lin. tanacet. ana ℈j f. p. ad lumbricos add. ☿ dulc. ℈ß.

SEMEN

SEMEN TANACETI.

Sunt Tanaceti vulgaris lutei C. B. P. cum ſemine rutæ & ſantonici conſpirant, ſunt vero quoad amaritiem remiſſiora, quoad acredinem aromaticam vero intenſiora. Cum ſantonico ex ▽ graminis datur verminantibus pueris & cum ▽ artemiſiæ contra Uteri obſtructiones.

℞. Sem. tanacet. ſanton. ana ℈j. ☿ dulc. ℈ß. f. p. contra lumbricos, vel cum conſerv. fl. perſic. ad ʒj f. bolus.

℞. Sem. tanacet. ʒß. caſtor. gr. xv. f. p. exhibendus ex ſyr. artemiſ. ʒiij & ▽ artemiſ. ℥iij contra Uteri obſtructiones.

***

SALE AROMATICO BLANDO CONFORTANTE CONSTANTIA.

GRANA KERMES.

Sunt ilicis aculeatæ cocciglandiferæ C.B.P. Grana excrementitia rotunda, piſi magnitudine, rubicunda, ejuſdem foliis adnata, dum recentia plena vermiculis & ſucco ſanguineo, fatiſcente in annoſis in ſubſtantiam raram fungoſam concolorem, ſaporis ſubacris ſubamaricantis, odoris non ingrati. Arbor fert glandes in calicibus, floret ut quercus noſtrates, ſunt igitur hæc grana ut gallæ quercuum excrementa, in trunco & ramis ejus majoribus excreſcentia. Adſperguntur aceto, hinc intereunt vermiculi, ſolent vero illico hæc grana una cum vermibus in ſyrupum coqui & ad nos deferri ex agro Monſpelienſi.

Sale

Sale aromatico blando Spiritus augente & confortante conſtant, unde decantatiſſima ſunt in ſyncope, lypothymia, palpitatione cordis, Melancholia, præcavendo abortu, confortanda memoria, in ſingultu. Horum vis conſiſtit in interiore pulpa ſeu in vermiculis incluſis.

℞. Gran. Kerm. ʒß. cap. ex ▽ borrag. c. ſyr. borrag. ad præcavendum abortum.

℞. Ambr. gryſ. ℈j. gran. Kerm. ʒij affund ℘ Meliſſ. q. ſ. ad diſſolvendam ambram. ſacch. cand. q. ſ. f. p. doſ. à ℈j ad ʒß, ad præcavendum abortum ab Uteri debilitate.

℞. Gran. Kerm. ℥ß. flaved. Cort. Citr. ℥ß. ℘us Cortic. Citr. q. ſ. f. infuſ. vel Eſſent doſ. cochl. ß. pro doſi.

℞. Eſſent.' gran. Kerm. cum ℘u Meliſſ. fact. ℥iiij extr. lign. aloës. ʒß. ſyr. flor. tunic. vel cinam. ʒv. M. f. mixtura ad confortandam memoriam.

℞. Conf. alkerm. ʒß. L. O. g. j. conſ. flor. borrag. ʒj M. f. bolus confortans & hypnoticus.

℞. Gran. Kerm. ℈j. Corall. r. præpar. ʒß. conf. alk. ſin. moſch. ʒß. cape ex ovo ſorbili ad præcavendum abortum.

VAINIGLIA.

Planta eſt convolvulacea ſcandens. Fructus ſunt inſtar vaginæ, habent ſubdulcem, ſubacrem aromaticam ſubſtantiam, in qua grana minutiſſima, odoris dum recentes gratiſſimi. vid. Franc. Hernandez.

Salem blandum aromaticum & confortantem in ſua mucaginoſa ſubſtantia fovet, virtute granis Kermes reſpondet, ſed potentius eſt & ſimul menſes & Urinam pellit, confortat memoriam & ſpiritus auget.

Datur

Datur ejus ſubſtantia, ingreditur chocolatam tanquam roborans.

℞. Vainigl. ʒj infund. 🜄 gramin. ferventiſſ. f. infuſ. ad ciendas Urinas.

℞. Vainigl. ʒß. conf. alkerm. ℈j ambr. gryſ. gr. iij in 🝕 cardam. ſolut. add. ſyr. cinam. q. ſ. ad confortandam memoriam, addi poteſt conſerva florum tunicæ.

SEMEN CORIANDRI.

Sunt Coriandri majoris C.B.P. ſemina rotunda, ſtriata, ex cinereo fuſca, includentia ſub cortice tenui medullam ſubfuſcam duriuſculam, in medio quaſi diſſepimentis diſtincta, ſaporis ſubdulcis ſubacris, odoris aromatici.

Ratione 🜔is blandi aromatici, & nonnihil oleoſi temperati, laudatur tanquam carminativum & ſtomachicum ſingulare, tollit quoque cruditates acidas Ventriculi, & inde enatos ructus, flatus, unde in præcavenda vertigine & apoplexia ex vitio ventriculi orta laudatur. Externe in flatulentis herniis laudatur.

Vulgo commendatur coriandrum, quod maceratum in aceto, ſed præſtat leviter exſiccatum, non maceratum. [Dum recens eſt habet odorem gravem, viroſum, caput tentantem, qui tamen per exſiccationem & progreſſu temporis evaneſcit,]

℞. Sem. coriandr. aniſ. fœnicul. ana ℥ß. Cinam. ac. ℥j Mac. ʒij flav. Cort. Citr ʒij pan. biſcoct. ℥iiij ſacchar. q. v. f. p. apoplecticus, poſt paſtum capiendus, addi poteſt aliquid ambræ & moſchi.

℞. Sem. Coriandr. ℥ß. ſem. cumin. aniſ. ana ʒij. ☉ crud. ʒß. Vin. & aq. ℔j coq. & applica decoct. hern. flatulentæ, addi poteſt ſem. naſturt. q. ſ.

SEMEN NIGELLÆ.

Sunt nigellæ flore ſimplici minore candido C.B.P. ſemina anguloſa, nigra, ſaporis ſubacris.

Ob blandum ⊖ aromaticum valde debile 🝆 craſſo incluſum, commendatur in Catarrhis & coryza tum interne, tum externe, ſi naribus admoveatur, ſeu oleo ejus expreſſo nares inungantur.

Foreſtus commendat ſemen ad menſes provocandos, præſcribitur tam crudum, quam toſtum, ut & 🝆 expreſſum.

℞. Sem. nigell. ʒij. 🝆 majoran. gt. iij vel iiij f. p. naribus admovendus in coryza.

℞. Sem. nigell. ʒij fol. majoran. p. iij. 🝆 majoran. gt. ij ſuccin. aniſ. ana gt. j. f. nodulus contra coryzam, addi poteſt aliquid ambræ & moſchi ut & ſtyrac. calamit.

***

SEMINA SALIBUS ACRIBUS FIXIS CONSTANTIA.

SEMEN SOPHIÆ CHIRURGORUM.

Sunt Naſturtii Sylveſtris tenuiſſime diviſi C.B.P. ſemina minuta ſubrubentia, ſaporis acris fervidi ad naſturtium accedentis.

Ratione ⊖is acris & fixi acidum dyſentericum infringunt, Urinam & calculum pellunt, externe ulcera ſordida mundant, eorumque acidum corroſivum infringunt.

℞. Sem. ſoph. ʒj conſerv. roſ. r. ʒiß. L.O. gr. j. 🝆 N. M. expr. gr. xv. f. bolus in Dyſenteria.

℞. Sem.

℞. Sem. ſoph. ℥ß cap. ex 🜄 Petroſel. vel gram. vel chæref. ad pellendum calculum & Urinam.

℞. Sem. ſoph. perfoliat. ana ℈j M. f. p. exhibendus ex decocto ſymphyti in herniis à nimia humiditate ortis.

SEMEN THLASPI.

Sunt Thlaſpi arvenſis Vaccariæ incano folio C.B.P. ſemina parva, oblonga, rufa, ſaporis acris & fervidi, reſpondent ſeminibus ſophiæ ſed potentiora & acriora ſunt. Propterea in rumpendis abſceſſibus internis, in pellenda Urina & menſibus uſurpantur. Cavendum vero ne facile gravidis exhibeantur quia Uterum nimis commovent & abortum cauſant. Externe ulceribus inſperſa, ea mundificant & abſtergunt.

℞. Sem. Thlaſp. ℥ß vel ℈ij Theriac. q. ſ. f. bolus ad rumpendos abſceſſus.

℞. Sem. Thlaſp. ℥ß. cap. ex 🜄 Puleg. vel Cinam. vel artemiſ. ad pellendos menſes & Urinam.

SEMEN NASTURTII.

Sunt Naſturtii hortenſis C.B.P. ſemina parva, oblonga, ſublutea, aliquantum compreſſa, in cacumen pallidum deſinentia, ſaporis acris amaricantis.

Horum 🜔: acre penetrantius ſubtilius & ſpirituoſius quam in ſemine Thlaſpi, reſpondent cochleariæ virtute, qua tamen paulo debiliora ſunt. Singulari quadam proprietate conveniunt in herniis tam interne, quam externe.

℞. Sem. Naſturt. ʒj cape ex vino rubro adſtringente ſingulis diebus. ℞. Sem. naſturt. album. ovi q. ſ. f. maſſa applicanda parti hernioſæ. Vel ℞. Sem.

Naſturt. gumm. Carann. ſagapen. M. c. album. ovi applica parti.

SEMEN SAXIFRAGIÆ.

Sunt pimpinellæ ſaxifragæ majoris C.B.P. conveniunt cum radice, quo recentiora eo præſtantiora ſunt.

***

SEMINA SALE VOLATILI ACRI CUM OLEO CRASSO CONSTANTIA.

SEMEN SINAPI.

Sunt ſinapi rapi folio C.B.P. ſemina rufa, rotunda, parva, ſaporis acris fervidi.

Ratione ſui ſalis acris, acida, fixa, terrea craſſa ſea mucaginoſa incidunt, attenuant, exterminant, hinc cibis craſſis, fumo induratis, ſub ſpecie embammatis tanquam appropriatum condimentum adduntur. In malo Hypochondriaco, inflationibus ventriculi, obſtructionibus lienis aliiſque morbis ab acido pendentibus tam interne quam externe conveniunt. Huc ſpectant cachexia, ſcorbutus, affectus ſoporoſi. Venerem quoque ſtimulant, diureſin promovent. Externe ſub forma ſinapiſmi applicantur cuti, ad rubefaciendum, ad excitandas puſtulas.

Oleum expreſſum externe applicatur in Paralyſi & affectibus frigidis.

℞. Sem. ſinap. maſticetur leviter & volvatur ſub lingua, ad præcavendum ſcorbutum.

℞. Sem. ſinap. ʒvj cardamom. ʒj caric. ping. n. vj vel vij. ∽ Cochlear, q. ſ. f. Elect. contra Apoplexiam.

℞. Sem.

℞. Sem. ſinap. ʒij. ſucci paſtinac. ʒij. 🜍 ovor. formicar. ʒj f. ung. quo glans inungitur ad tollendam impotentiam. Interne ℞. Elect. diaſatyr. ʒij ſem. ſinap. ʒß. f. bolus.

℞. Sem. ſinap. p. j. bacc. junip. p. ij infund. 🜄 vin. camphorat. vel 🜄 C. C. ſuccinat. q. ſ. applica membro paralytico.

SEMEN NAPI.

Sunt Napi ſativæ C. B. P. ſemina parva, rotunda, ſubrufa, ſaporis acris amaricantis.

Remiſſiore ſunt acredine, quam ſemen ſinapi, interne in Variolis & Morbillis pellendis commendantur, de cætero cum ſinapi conveniunt.

℞. Sem. nap. aquileg. Card. B. ana ʒß. ▽ fœnic. ſ. chærefol. q. ſ. f. Emulſ. ad Variolas & Morbillos, ſtimuli loco add. ⊖ volatil. C. C. gr. vij. ad ℈ß.

℞. Sem. nap. ſinap. eruc. raphan. ana q. v. affund. 🜄 vin. Camph. f. epithema antiparalyticum. 🜍 expreſſum convenit cum 🜍 ſinapi.

Hoc 🜍 in Paralyſi poſt dolores colicos & iſchiaticos feliciter ſæpe uſus eſt Hermannus. ℞. 🜍 Napi, eruc. ſinap. oleris atri. raphan. ana ℥ß. camph. ʒiij f. liniment.

SEMEN OLERIS ATRI.

Sunt braſſicæ fimbriatæ C. B. P. ſemina rotunda, glabra, nigricantia, ſaporis acris aromatici, odoris grati ſed debilis.

Convenit cum ſeminibus napi.

SEMEN RAPHANI.

Sunt Raphani majoris orbicularis vel rotundi C.B.P. ſemina rotunda, rutila, parva, ſaporis acris.

Serum craſſum incidunt, & ad Urinam deponunt, adeoque cum Napo conveniunt.

℞. Sem. raphan. ʒj. ▽ chærefol. q. ſ. f. Emulſ. diuretica, quæ ſimul incidit mucum primarum viarum.

℞. Sem. raphan. cape ex proprio ſucco radicis, addito tantillo ⊖is communis, pro diuretico & calculo pellendo.

℞. Sem. raphan. ℥ij coq. ex ▽ f. apozema diureticum.

***

SEMINA SALE FIXO AMARO CONSTANTIA.

SEMEN FUMARIÆ.

Sunt ſemina rotunda, parva, ex nigro viridantia, ſaporis amari ingrati. Ob ⊖ fixum cum paucis volatilibus laudantur in depurando ſanguine ubi acido-viſcoſa, ſcorbutica, icterica & ſcabioſa diſpoſitione laborat, potentior tamen eſt herba integra, quam vide.

℞. Sem. fumar. ʒß exhibe ex ▽ chelid maj. contra acidas Cachexias.

℞. Sem. fumar. Naſturt. ſoph. ana ℈j f. p. diaphoretic. contra ſcabiem & malum hypochondriacum.

℞. Sem. fumar. ʒß. croc. or. gr. vij Camphor. gr. iij cape ex conſerv. borrag. ſ. fumar. ſ. hyperic. contra malum hypochondriacum.

SEMEN

SEMEN CINÆ.

Semen Cinæ vel Zinæ, Santonici. Zedoariæ, ſanctum, it. ſem. lumbricorum.

Sunt Abſynthii ſantonici Alexandrini C.B.P. ſemina parva, oblonga, quaſi ex ſquamoſis funiculis compacta, flaveſcentia, ſaporis acris amari, odoris graviuſculi.

Vermibus unice ſunt dicata, tum quod vermes horum amaritiem ferre neſciunt, tum quod horum ⊖ia fixa putridum acidumve vermium nidum deſtruant.

℞. Sem. Cinæ ʒij Mannæ ℥ij ſolv. in ▽ gramin. cape contra vermes add. ☿ dulc. gr. vij pro qualibet doſi, addito ſaccharo f. rotulæ ſ. morſuli.

℞. Sem. cin. tanacet. ana ℈j. ☿ dulc. gr. vij. M. f. p. contra vermes, exhibendus ex decocto radicis filicis, polypod.

℞. Summit abſynth. ſem. tanacet. Cinæ. ana q. v. f. c. ♒u vini extractum, cujus inſpiſſati cape ℈j add. ☿ dulc. gr. vij. fac pilulas ☉o foliato involvendas contra lumbricos & vermes.

SEMEN CARDUI BENEDICTI.

Sunt Cnici Sylveſtris hirſutioris ſive Cardui Benedicti C.B.P. ſemina oblonga craſſiuſcula, ſtriata, ex fuſco ſublutea, ſaporis inſigniter amari.

Salia fixa includunt in mucagine tenui, hinc blandos movent ſudores & Urinas, relinquuntque in receſſu vim anodynam, & propterea in pleuritide, peripnevmonia, variolis & morbillis commendantur.

℞. Sem.

℞. Sem. Card. B. Card. Mar. bardan. pap. alb. ana ℥ij. ▽ flor. pap. rhœad. q. ſ. f. Emulſ. cum qua cape mandib. luc. piſc. l. dent. apri. l. ♄ diaph. ℈j contra pleuritidem.

℞. Sem. Card. B. ℥ij. ▽ Cent. min. q. ſ. f. Emulſ. add. ⊖ prunell. ℈j exhibe contra febres intermittentes hora ante paroxyſmum. add. tinct. Caſtor. ſ. Myrrh. ſ. ⊖ volat. C. C. exhibe contra Variolas & Morbillos.

SEMEN CARDUI MARIÆ.

Sunt Cardui albis maculis notati vulgaris C. B. P. cum Cardui Benedicti ſeminibus omnino conveniunt, verum ad fluorem album commendantur ſpecifice.

℞. Sem. Card. Mar. ℥ij exhibe ex vino ſaſſafr. ſ. ex ▽ fœnicul. contra fluorem album.

SEMEN BARDANÆ.

Sunt Lappæ majoris. Arcii Dioſcoridis C. B. P. ſemina oblonga, ſtriata, fuſca, craſſiuſcula, ſaporis ſubamari, ſubacris, odoris nullius. Radici reſpondent in omnibus.

℞. Sem. bardan. ℥ij violar. ℥j Xanthii ℥ß f. Emulſ. ex ▽ gramin. ad calculum.

℞. Sem. bardan. ℥ij Card. B. ℥j f. Emulſ. ex ▽ chærefol. cum qua exhibe ♌ ⊖ ✳ci ℈j contra hecticam & phthiſin.

SEMEN URTICÆ.

Sunt urticæ vulgaris urentis maximæ C.B.P. ſemina parva, compreſſa, lubrica, ſpadicea, lævia, ſaporis ſubacris & tenuis.

Romanæ

Romanæ ſunt urticæ urentis pilulas ferentis I. Dioſc. ſemine lini C. B. P. ſemina pene orbicularia, compreſſa, lubrica, lævia, ex ſpadiceo nigricantia, ſaporis ſubacris, pinguis, cum quadam tenuitate.

Majoris, minoris & Romanæ gaudet ⊖e diuretico ℅ſa quaſi mucagine temperato, unde non tantum in affectibus renum & veſicæ, ſed etiam in phthiſi, tuſſi, dum materiam peccantem ad Urinaria vaſa deducunt proſunt. Laudantur ſpecifice in qualibet interna hæmorrhagia, nempe hæmoptyſi, vomitu ſanguinis, mictu cruento, fluxu hæmorrhoidum.

℞. Sem. urtic. teſt. ovor. N. M. toſt. f. p. ad dyſuriam, omiſſa N. M. pellit quoque Urinam, capiendus ex ſucco urticæ.

℞. Sem. urtic. ʒij Carthami. ʒj. ▽ Hyſſop. q. ſ. f. Emulſ. contra tuſſim, addi quoque poteſt tantillum ſyrupi papav. l. violar.

℞. Sem. urtic. ʒj cape ex ſucco proprio ſeu decocto ad quamvis hæmorrhagiam.

℞. Veron. Mij heder. terr. Mj. rad. helen. ℥ß. ir. florent. ʒiij coq. in ▽ addito hydromell. ℞. Hujus decoct. q. v. fac cum ſemine Carthami & urticæ Emulſ. contra tuſſim.

*
* *

SEMINA CONSTANTIA SALE BLANDO DIURETICO.

SEMEN ALKEKENGI SEU HALICACABI.

Sunt Solani veſicarii C. B. P. baccæ rotundæ, ceraſi magnitudine, incluſæ folliculo folioſo, per maturita-

turitatem rubente, continentes ſub pellicula tenui pulpam flaveſcentem, in qua ſemina compreſſa, orbicularia, rugoſa, ſaporis ſubacris, ſubamari, odoris nullius evidentis. [Fructus hi dulcacidi ſolo contactu digitorum leviſſimo amareſcunt. conf. G. H. Welſch. obſ. Med. Phyſ. Hecatoſt. 1. O. 9. p. 10.]

Fructuum horum virtus contra icterum, ſanguinem congrumatum, arthritidem, ſtranguriam, Urinæque remoram, conſiſtit in ⊖e diuretico, qui materiam morbificam per vaſa Urinaria excludit. Fumus ſeminum ore exceptus mire prolicit vermiculos ex cavo dentis.

Datur eorum ſubſtantia, trochiſci, ▽ deſtillata cum corticibus Citri laudatur ab Ettmüllero. [Cum cort. Citri præbet ▽am Nephriticam ab Hœfero in Hercule Medico l. 3. p. m. 176. deſcriptam & laudatam.]

℞. Sem. alkek. ʒj exhib. in ⌒u vel vino Junip.

℞. Sem. alk. ʒß cape ex ▽ chel. maj. contra icterum.

℞. Ocul. ♋. ſem. alkek. ana ℈j ſperm. Cet. ℈ß. ⊖ vol. C. C. gr. v. f. p. cap. ex ▽ chæref. ad diſſolvendum ſanguinem congrumatum.

℞. Fruct. alk. q. v. infund. ⌒u Junip. facto per ferment. f. eſſentia diuretica, quæ ſi exhibeatur cum Tinct. Caſtorei valet, ſi Urina ſuppreſſa à cauſa quadam Paralytica.

[De virtute diuretica baccarum alkekengi ſeq. hiſtoriam habet Arnold. de Villanova in libro de vinis, his verbis: Fuit tempore meo Cardinalis quidam, qui jam per quatriduum Urinæ abnegationem paſſus eſt, turgidus, & ut dicitur, deſperatus, nullis aliis proficien-

ficientibus Medicinis, qui de conſilio parvi Medici per potum vini alkekengi liberatus eſt, & ſoluta eſt Urina in quantitate, ut impleretur quaſi concha una. Ex eo ſolo experimento Medicus ille cum parva ſcientia & pauper, factus eſt magnus & dives & multis quidem Deus dat gratiam & affert utilitates & contra, & totum eſt, quia judicia Dei abyſſus multa.]

SEMEN GENISTÆ.

Sunt geniſtæ anguloſæ & ſcopariæ C.B.P. ſemina reniformia, compreſſa, fuſca, ſaporis ex dulci amaricantis nauſeoſi & ingrati, dum annoſa farinacei.

In ſubſtantia farinacea latet ⊖ diureticum, quod per incinerationem copioſum elicitur, unde in movenda Urina uſurpatur.

Semina recentia quippiam ſapiunt nauſeoſi, unde quoque horum infuſum ex ▽ fervente factum blandos Vomitus movet, ob quos in arthridite prædicantur. Cæterum quoque alvum nonnunquam movent, ſi cum hydromel. dantur.

℞. ⊖ geniſt. gr. vij ſem. geniſt. ʒß. f. p. diureticus, exhibendus ex ▽ chærefol. vel ſi alvus ſit adſtricta ex Emulſione carthami.

℞. Sem. geniſt. ʒj cap. c. hydrom. ad movendam alvum & Urinam, in ſpecie ſi adſit hydrops, icterus & tumor lienis.

℞. Sem. geniſt. cont. ℥ß vel ʒvj infund. ▽ fervent. q. ſ. cape infuſum tepidum ad provocandum Vomitum.

[Aliàs flores geniſtæ nondum expanſi eadem ratione ut cappares condiuntur, & à Nephriticis condimenti loco cum fructu uſurpantur.]

SEMEN MILII SOLIS.

Sunt Lithoſpermi erecti majoris C. B. P. ſemina rotunda, glabra, lævia, ſplendentia, cinerea, duriſſima, ſaporis farinacei, odoris nullius.

Subſtantia horum in omnibus fere cum oculis ♋ convenit, hinc in urina & calculo expellendo exhibentur. Freitag. [in auror. Medic.] ſuo, experimento laudat in movendo partu, it. in gonorrhæa ſiſtenda.

℞. Sem. mil. ſol ʒj. ⊖ geniſt. ℈j f. p. diureticus, vel adde lap. prunell. ʒß. Datur cum Rhabarbaro & moſcho in ſanguine congrumato.

℞. Sem. mil. ſoliis ʒij exhibe cum lacte ad promovendum partum.

℞. Sem. mil. ſol ʒiß. pulv. ling. cervin. ſ. phillitidis. ʒß. ſuccin. præpar. ℈ij M. f. p. exhib. ʒj ſingulis diebus contra gonorrhœam.

SEMEN RUSCI.

Sunt Ruſci C.B.P. baccæ rotundæ, aſparagi baccis majores, rubentes, includentes ſub pellicula tenui modo unum, modo binos acinos duriſſimos, fulvos, ſaporis ſubdulcis.

Conveniunt cum radice, quam vide.

Ad excreſcentiam in ductu Urinario tollendam.

℞. Rad. ruſc. ℥ß. ſem. ruſc. ℥ij. f. decoctum.

SEMEN ASPARAGI.

Sunt Aſparagi ſativæ C.B.P. baccæ rotundæ, rubræ, piſi magnitudine, includentes tres acinos nigros corneæ duritiei, ſaporis ſubacris, & ſubdulcis, odoris nullius. Convenit cum radice.

℞. Sem.

℞. Sem. aſpar. ruſc. alkek. mil. ſol. ana ℈j. ⊖ geniſt. ℈ß l. lap. prunell. gr. xv. f. p. diureticus, exhib. ex ▽ chærefolii.

SEMEN COFFEE.

Eſt Euonymo ſimilis Ægyptiaca, fructu baccis lauri ſimili C.B.P. ſponte naſcuntur arbores in Perſia, Arabia felici, ad magnitudinem tiliæ. Sunt igitur hujus nuclei pallidi, faba paulo minores, altera parte convexi, altera plani, inſigni ſulco notati, bini, ſibi invicem cohærentes, tecti folliculo membranaceo tenui, ex fuſco nigricante.

Coffee crudum more leguminum in farinacea ſubſtantia fovet ⊖ fixum diureticum, quod in uſtulato paululum extricatur, ut à ſuperfuſa ▽ fervente reſorberi vel imbibi poſſit, hinc ſolum uſtulatum Coffee uſurpatur in infuſo ▽æ ferventiſſimæ.

Sale quidem conſtat diuretico, ſed admodum blando & nonnihil volatili & temperato, unde viſcida diſſolvit, & ad Urinam diſponit, ſuppreſſos torpidoſque ſpiritus ſuſcitat, & alacres ad obeunda munera reddit. Conducere videtur potius obeſis & craſſis, pituitoſis vel phlegmaticis, quam bilioſis & macilentis, & quorum ſpiritus inquieti & nimis agiles.

In Perſia uſtulantur cortices ſeu tegumenta granorum. Singulis diebus grana Coffee cremant pro uſu, procraſtinare non ſinunt, optimum eſt infuſum, quod ex recenter uſtulatis granis eſt confectum. Cortices raro ad nos transferuntur, aliàs eorum infuſum multo præſtantius.

***

SEMINA SALE PURGANTE ACERRIMO CONSTANTIA.

SEMEN MEZEREI SEU LAUREOLÆ.

Sunt Laureolæ folio deciduo ſeu flore purpureo C. B. P. baccæ rotundæ pipere majores, rubentes; per maturitatem nigricantes, includentes ſub tenui pellicula nucleum rotundum ſaporis fervidi & nauſeoſi. Vocatur piper Germanicum.

Tam potenter purgante & cauſtico ⊖e conſtant, ut earum uſus non niſi robuſtiſſimis concedatur, licet plebeji tribus integris granis deglutitis ad hydropem utantur. Extractum cum 🜅u vini ſeu aceto factum, mitius eſt & ad ℈ß cum caryophillis aromaticis tanquam correctorio hydropicis propinari poteſt. Interim nonnulli hoc Mezereo eorumque extracto ſtimulant & quaſi reſtituunt purgantia effœta.

℞. Gran. mezer. gr. iij integra, cape ex ▽ roſar. pallid.

℞. Conf. Hamech. ʒß. gran. mezer. gr. iij. f. bolus.

℞. Mezer. q. v. infund. vin. cydonior. exime grana & exſicca & exhibe pro doſi gr. iij integra vel in pulverem redacta.

℞. Mezer. q. v. macera in lixivio, fac turundam & inſere auriculæ perforatæ ad oculorum inflammationem tollendam, exemplo Riverij. [Conf. G. H. Welſch. Mictomim. Cent. 2. obſ. 63.]

SEMEN

SEMEN STAPHISAGRIÆ.

Sunt Staphiſagriæ C. B. P. ſemina rugoſa, ſcabra, nigra, anguloſa, triquetra, ſaporis fervidi.

Semina ſunt plantæ quæ refertur ad angioſpermas multiſiliquas ſiliculis corniculatis oblongis, floribus cucullatis quæ vel purgantes vel venena ſunt, ut aconita, napelli, helleborus. Eſt enim ſtaphiſagria ſpecies aconiti.

Sale purgant potius volatili quam fixo, verum ἄνω καὶ κάτω purgante conſtant, ſed cum ob violentam & moleſtam acredinem fauces exurant, & metum ſuffocationis & ſtrangulationis inferant, uſus internus ideo rariſſimus, interim maſticata & ſub lingua volutata ſalivam adeo copioſe proliciunt, ut inter apophlegmatizantia referantur. Propterea Sylvius [Meth. Med. l. 2. c. 13. ſ. 59. & 60. p. m. 115.] pro ptyaliſmo excitando ad L. ♀ commendat, ſed præterquam, quod ob prædictam moleſtiam eorum uſum ægri ferre recuſent, inſufficientia & plane incongrua videntur ad Luem ♀eam. Licet enim ptyaliſmum inducant, non talem tamen, quæ ita ſerum fundit, veluti ☿ialia faciunt. Salivatio enim tamdiu adeſt, quamdiu ſapor acris in lingua. Pediculis ſingulariter exitioſa ſunt, & ob id unguentum pediculorum ingrediuntur.

Vis purgans juvatur oxymelle. Saporis moleſtia corrigitur cubebis & nuce moſchata.

℞. Staphiſagr. ℈j oxymel. ℥j M. pro ſurſum & deorſum purgante.

℞. Sem. ſtaphiſagr. ſinap. ana ʒj rad. pyrethr. ʒiß. Cubeb. ʒj piper. alb. ℈j N. M. ʒj f. p. apophlegmatizans contra Catarrhos.

℞. Sta-

℞. Staphiſagr. q. v. coq. cum ▽ tene decoctum in ore ad ptyaliſmum excitandum.

℞. Mucag. helen. ℥ij ſtaphiſagr. ℥ß. ☍ laur. q. ſ. f. ung. add. ♀is ʒj & ſi videtur ☿ii vivi ʒß. ad enecandum pediculos.

SEMEN PSYLLII.

Sunt Pſyllii majoris erecti C. B. P. ſemina parva oblonga, puliciformia, lubrica, lucida, ex fuſco punicea, ſaporis primum mucilaginoſi, poſtmodum acris & nauſeoſi.

In horum mucagine acria quævis temperante, occultatur ⊖ blandum, bilem expurgans.

Sal purgans volatile probant ſemina cocta & uſtulata, quæ amiſſa virtute purgante adſtringunt. Bilis expurgationem evincit uſus, nam acres bilioſos humores manifeſte evacuant, eoſque ob mulcentem mucaginem obtundunt & temperant. Cæterum ob eandem mucaginem proſunt in raucedine, angina, ardore & ſiccitate faucium & tuſſi. Exterius conveniunt ad ambuſta.

Rarius uſurpatur ob ſui mucaginem, quæ ob tenacem adhæſionem Ventriculo & Inteſtinis nonnunquam torminoſa eſt. Datur ſubſtantia à ʒj ad ʒij. Datur & Emulſ. Elect. & ſyr. Vis purgans juvatur manna, hydromel ſtimulatur diagrydio. Vis temperans & refrigerans intenditur æquepollentibus nempe ſemine cydoniorum. Mucago ſeminum toſtorum additur clyſteribus ad dyſenteriam & acredinem temperandam.

℞. Sem. Pſyllii ʒiß. f. Emulſ. c. hydromel. ad alvum laxandam.

℞. Sem.

℞. Sem. pſyllii ʒij violar. ʒij ▽ violar. ℥ij contunde & exprime, expreſſo add. Mann. ℥j exhibe cochleatim infantibus per vices ad laxandam alvum, ſi robuſtis exhibendum add. gr. j. vel ij diagrydii.

℞. Sem. pſyll. cydon. lini ana ʒj ebulliant leviter in ▽ roſar. add. integrum ovum, ℴℴ roſar. q. ſ. ſacch. ♄n. ad fuſcum colorem inducendum ☿ viv. ʒß quo Hermannus cum ſucceſſu uſus eſt in hæmorrhoidibus.

℞. Sem. pſyll. cydon. gum. tragac. ana ʒj f. mucilago c. ▽ roſar. in inflammatione hæmorrhoidum.

℞. Sem. pſyll. ℥ß. ſem. cydon. papav. ana ʒj. ▽ roſar. calid. ℥ij f. mucago add. ſacchar. penid. ℥j ad raucedinem, ardorem faucium & tuſſim.

℞. Sem. pſyll. ℥ß. ▽ Nymph. plantag. ana ℥ij ſyr. violar. q. ſ. f. mixtura refrigerans vel temperans ſi adſit aridura linguæ.

℞. Mucag. pſyll. cydon. ſacch. ♄. ℴℴ lini q. v. f. linimentum contra ambuſta.

*
* *

PURGANTIA CONSTANTIA SALE ACRI PURGANTE TAM VOLATILI QUAM FIXO.

COLOCYNTHIS SEU ALHANDAL.

Sunt Colocynthidis fructu rotundo minoris C. B. P. fructus rotundi, pugni magnitudine, ex flavo virides, includentes ſub cortice tenui coriaceo medullam candidam, fungoſam, levem & ſemina parva compreſſa, oblonga, cucumeracea, quaſi ſordida, albicantia, ſaporis acris amari & nauſeoſi.

Inter purgantia vegetabilia validiſſima eſt, humores quoſcunque viſcido pituitoſos & tenaces, qui cæterorum purgantium vim omnem eludunt, ex ultimis quaſi articulis provocat, & per ſuperiora, frequentius tamen per inferiora educit, hinc tanquam catholicum in morbis pertinaciſſimis, Iſchiatico dolore, Lue ♀ea commendatur. Vermibus quoque tam interne quam externe contrariatur.

Ingreditur clyſteres pro ſtimulo in ſpecie contra morbos ſoporoſos, ſed cave ne pulpam injicias ſed ſindoni includas. Additur quoque pro ſtimulo fotibus vaporoſis ad menſes & fœtum mortuum expellendum, jungitur & ſuppoſitoriis.

Vis conſiſtit in ſubſtantia fungoſa & reſinoſa. Semina ex uſu proſcribuntur.

In ſubſtantia pulpa per os raro datur, quia ob nimiam levitatem Ventriculo adhæret, & tormina & hypercatharſin cauſat, ſed prius cum tragacanth. bdell. vel gummi Arabico formatur in trochiſcos, qui à gr. vj ad ℈ß ad ſummum exhiberi ſolent.

Externe applicata alvum ſolvit. Ita ☍ colocynthidis coctum externe applicatur.

Datur reſin. ad gr. x Extract, ad gr. xij.

Virulentia corrigitur ⊖e ♀ri, aceto, vel Urina in qua pulpa maceratur. [De hac re ita Laz. River. Prax. Med. l. I. c. 15. p.m. 191. Colocynthis ad cerebrum expurgandum efficaciſſima eſt, ſed ſummo amarore infenſa eſt, vehemenſque illius eſt operatio. Corrigitur utrumque infuſione in Urina, ſic enim amaritudinem deponit, & fere inſipidum remedium evadit, tum vehementia illius infringitur ut ad ʒj tuto exhiberi valeat, fitque excellens ad omnes capitis morbos à

cauſa

cauſa frigida oriundos remedium.] Aſt optima corcectio fit ⟜u vini, cujus quoque beneficio reſina & extractum fit, quæ optima & tutiſſima colocynthidis præparata ſunt.

Vis tormentoſa emendatur Mace, Nuce Moſchata, cinam, &c. ſpecificatur variis morbo appropriatis.

℞. Troch. alhand. ℈j infund. vin. coq. par. f. cholagog.

℞. Pulp. Coloc. ſ. potius troch. alhand. gr. x. Mac. gr. v. conſ. roſ. pallid. ſ. fl. per ſic. ʒj M. f. bolus.

℞. Fol. ſalv. flor. chamom. ſumm. origan. ana Mj. rad. pyrethr. ℥j pulp. Coloc. in pet. ligat. ʒj coq. in ſ. q. ▽. ℞. Colatur. ℥x add. El. hier. picr. ℥j Mell. anthoſ. ʒvj Extr. Caſtor. ʒj ⊖ gemm. ʒj f. Clyſter contra morbos ſoporoſos.

℞. Oleum Colocynth. M. c. ung. d. arthanita q. ſ. ad formam linimenti, hoc inunge totum abdomen in alvi adſtrictione.

℞. Helleb. n. ʒß. calam. ar. ℈ij pulp. colocynth. ℈j. bacc. laur. ʒj ſem. cumin. ℈ij ⟜us vini ℥iiij f. infuſum, quod guttatim inſtilla auri ad ſurditatem tollendam.

℞. Extr. coloc. ℈ß. extr. hell. n. gr. iij. ℅ d. caryoph. gt. j. Maſtich. gr. vij. M. f. pil. purgantes.

℞. Pil. aloëph. gr. xv. Extr. coloc. gr. iij. ℅ Mac. gt. j. M. f. pil. cephalicæ, convenientes in Catarrhis.

℞. Pil. Maſtichin. gr. xv. Extr. Coloc. gr. iij. ⊖ ſuccin. volat. gr. v. f. pil. cephalicæ anticatarrhales.

℞. MP. fœtid. gr. xv. Extr. Coloc. gr. iij Caſtor. gr. v. f. pil. Uterinæ.

Reſina colligitur ex pulpa Colocynth. cum ⟜u vini, poſtea præcipitatur cum ▽ communi.

℞. Reſin. Coloc. gr. v. Diagr. gr. iij. ⊖ ♀ri gr. iij. ☍ cinam. gt. j. conſerv. roſ. pallid. ʒß. M. f. bolus.

Olei deſtill. coloc. per 6am ſi abdomen eo inungatur juvat in fœcibus duriſſimis vel etiam ☍ ejus coctum.

NUX VOMICA.

Nux Vomica in officinis C.B.P. ſunt arboris Malabaricæ fructuum nuclei orbiculati, plani, compreſſi, cornea duritie, colore argenteo, foris parum lanuginoſi, ſaporis amari. Arbor eſt nucifera, floribus monopetalis oblongis in quinque ſegmenta diviſis, racematim diſpoſitis, fructibus rotundis aurantiæ forma & magnitudine. Hi fructus includunt ſub putamine tenui duriuſculo, fragili, pulpam mucilaginoſam involventem nucleos.

[Conf. ſupr. de ligno colubrino p. 113. ſ. Arbor hæc ſub nomine Caniram deſcribitur in Hort. Malabar. T. 1. p. 67.]

Sale purgante emetico adeo hoſtili conſtant, ut citra convulſionis metum vix interne adhiberi queant. Veteres eas alexipharmacas pronunciarunt, & ad Electuarium de ovo dictum receperunt, aſt conſultius videtur eas ex hoc Electuario & ex alexipharmacorum numero præſcribi. v. Zvvelff. An. Ph. Aug. p. 259. Aſt in tam parva quantitate ingreditur, ut vim ſuam minus exercere poſſit. Electuarii ʒj vix ultra gr. j. ingreditur. [Nonnulli etiam exhibuerunt in febribus tertianis ſeu quartanis ſed in dubiis recte mitiorem & tutiorem ſequimur viam conf. Wedel. Amœn. Mat. Med. l. 2. ſ. 2. c. 4. p. 337.]

D. Hermannus exhibuit integram nucem Vomicam, aliquando cani, qui exinde mortuus eſt ſpumeſcenſque

ſcenſque ſanguis in Pulmonibus fuit repertus, itaque diſſolvit ſanguinem. Canibus aliquot grana data omnino exitioſa ſunt, dum ſpiritus in turbas concitant, ſpaſmos convulſiones & tetanos excitant, ſanguinis nexum ſolvunt, & tandem ſpiritus extinguunt.

Pari modo infeſtæ ſunt corvis & cornicibus; interim quibuſdam avibus indicis noxiæ non ſunt, nam ex horum ventriculis ſæpe integræ nuces eximuntur, & reperiuntur inter alvinas fœces ex iiſdem rejectas. Indi cum aceto tritas imponunt capitis doloribus: quidam ex his infuſum parant ad quartanam v. c. ℞. Raſur. Nuc. Vomic. ʒj infund. vin. abſynth. exhibe infuſum contra quartanam, ſumitur hora ante paroxyſmum.

*
* *

SALE PURGANTE TAM FIXO QUAM VOLATILI OLEO CRASSO INCLUSO CONSTANTIA.

GRANA TILLI.

Pinus indica nucleo purgante C. B. P. ſunt ricini arboreſcentis ſemina, ſeu fructus oblongi, ovales, cataputia media paulo majores, una parte gibbi, altera paulo planiores, quatuor levioribus cingulis notati, includentes ſub pellicula corticoſa cinerea, nucleum albicantem, pinguem oleoſum, ſaporis exurentis.

[Vid. ſupr. Pavana ſ. lign. Moluccenſe p. 117.]

Adeo potenter humores ſeroſos, viſcidos & tenaces incidunt, & per ſuperiora & inferiora educunt, ut hoc nomine ipſam Colocynthidem ſuperent, adeoque in morbis chronicis contumacibus, in ſpecie in hydro-

pe, tanquam catholicum propinantur, pro ſtimulo potius quam pro baſi.

Vis concentrata eſt in foliolis iſtis ſeminalibus, quæ in media granorum ſubſtantia latent, ut & in oleo craſſo quod prelo ex granis elicitur.

Ob excedentem ſui acredinem, fauces, palatum & nonnunquam anum inflammat. Datur in forma pilularum, & corrigitur coctione, vel cum aceto vel in Urina puerorum. Corrigitur Zinzibere, ſucco glycyrrhizæ aliiſve vel aromatibus, vel dulcibus viſcoſis. Alii ſaccharum ♄ni & opium pro correctorio addunt. Cortices blandius purgant, ac ipſa ſubſtantia interior.

Datur integra ſubſtantia nucleorum, rejecto cortice exteriore, licet purgativa virtute haud careat, à gr. iij ad v. unumquodque granum, ſedem molitur unam, ſi ▽ calida vel juſculum ſuperbibatur, ita tamen ut iij grana, communiter v faciant ſeceſſus. Alvus vero in momento ſiſtitur, ſi ſuperbibitur hauſtus largior ▽æ frigidæ vel ſi pedes madefiant ▽, vel ſi manus laventur ▽ frigida & naribus admoveatur.

Dantur duo folia ſeminalia fructus medietati incluſa, vulgo germen dicta vel Cyma, quantitate ad j. vel ad ſummum gr. ij.

Datur 🝆 gran. til. expreſſum gr. j. robuſtiſſimo cum oleo ex ricino vulgari expreſſo. Datur & extractum beneficio ſpiritus vini factum à gr. iij ad v.

℞. Gran. till. gr. iij z. z. gr. vij M. f. pil. iij vel iiij.

℞. Fol. ſeminal. gr. till. gr. ij Zz. gr. v. ſucc. glycyrrh. in ▽ roſar. ſol. q. ſ. f. pil. j. ad purgandum ſerum.

℞. Extr. Cathol. gr. xv. gran. till. gr. j. vel ij ſyr. roſ. q. ſ. f. pil.

℞. Extr.

℞. Extr. gum. gutt. gr. iij reſin. Jalap. gr. vij gr till. gr. ij Zinz. gr. v. M. f. pil. antihydropicæ.

℞. Fol. ſeminal. gran. till. gr. j. fol. ſenn. ʒß Zinz. gr. v. ſacchar. q. ſ. ſcil. ʒß vel j. f. pulv. l. ſacchar. ſ. q. ſolv. in ▽ roſar. & ad conſiſt. coq. f. rotulæ purgantes.

℞. Olei expr. gran. til. gt. j. exhibe c. ☍ expr. ſemin. ricin. vulg. ad cochl. ß vel ☍ amygd. dulc. rec. expreſſ. cochl. j. exhibe pro purgante. [Deſcribit hoc Petr. Borellus ſub titulo facilis purgationis C. 3. obſ. 29. p.m. 221. ſi olei nuclei pinei indici expreſſi guttæ duæ potui infundantur, vel juſculo, & ſic procurabit purgationem facilem. Addit: alii eo umbilicum fricant, & ſic etiam incaute ægrum purgant.]

Olei hujus expreſſi uſus frequentior eſt externus. I. In alvi retentione, ſi eo umbilicus inungatur. Scil. ℞. ung. d. arthanit. ʒj ☍ Coloc. ʒiß. ☍ gran. till. expr. gt. xij f. liniment. quo inungatur abdomen. II. In inflammatis hæmorrhoidibus cœcis & carunculis exedendis. ℞. ☍ expr. gran. till. gt. v. add. ſacch. ♄ni par. f. linim. quo inungantur hæmorrhoides cæcæ inflammatæ, & ani carunculæ, 3 vel 4 de die repetatur. III. In excitandis ſedibus, dum ejus odor naribus excipitur. Hinc ex eo conficitur pomum regium catharcticum ſequenti modo. ℞. ☍ expr. gran. till. q. ſ. huic immitte pomum aurantium ſeu Citreum & relinque per menſem integrum, poſt exime & aſperge pulvere ſantali Citrini, admove naribus ad alvum movendam, vel frica hoc pomo fortiter manus & ad move naribus, quotieſcunque pomum naribus admovetur, ejuſque odor fortiter attrahitur, toties movebitur alvus.

RICINUS

RICINUS MEDIUS.

Sunt Ricini vulgaris C.B.P. ſeu Cataputiæ mediæ fructus oblongi, ovales, faba minores, includentes ſub pellicula corticoſa, livida & maculoſa nucleum bipartitum album, pinguem, ſaporis ſubdulcis, acris nauſeoſi.

Vim purgantem incluſam habet oleo temperatiore, ita ut 🝆 expreſſum ex hoc ricino ad integrum cochleare exhiberi poſſit, pro laxanda alvo, in ſpecie ſerum & aquam evacuat.

Dantur grana in Emulſione, ad ʒß. vel j. ex ▽ fumar. vel Cichor. it. in ſubſtantia dantur à gr. x. ad xv vel xx.

Oleum, vulgo infernale dictum à cochl. ß. ad integrum exhibetur pro purgante, externe vero applicatum partibus paralyticis medetur.

℞. Ricin. gr. x. f. Emulſ. ex ▽ Cichor. q. ſ. add. pro ſtimul. gr. ij. ſcammon.

℞. Olei ricin. expr. cochl. ß. ad alvum laxandam. vel ℞. 🝆 ricin. expr. inunge regioni umbilicali pro alvo laxanda.

℞. Olei ricin. med. expr. ℥ij. 🝆 Nuc. Moſch. expr. ℥ß. M. f. linim. contra pedes paralyſi affectos, quem affectum Indi & Bontius Beriberi vocant. [De affectu hoc præter Bontium agit quoque Tulp. libr. 4. c. 5. p. m. 300. & Ol. Borrich. act. Hafn. Vol. 4. obſ. 43. p. m. 144.]

RICINUS AMERICANUS.

Sunt Ricini Americani majoris ſemine nigro C.B.P. ſeu Cataputiæ majoris fructus oblongi, ovales, fabæ

fabæ magnitudine, una parte convexi, altera depreſſi, includentes ſub pellicula corticoſa tenui duriuſcula nucleum candidum bipartitum, ſaporis cum præcedente.

Validior eſt ricino medio & propterea in minori doſi præſcribendus, cæteroquin utiliſſima Medicina purgans hydropicis.

Vis hujus conſiſtit vel in nucleo, vel oleo, potiſſimum foliis ſeminalibus. Subſtantia datur à gr. v. ad vij. vel ix. datur in emulſione, datur & 🜖 ejus expreſſum.

℞. Ricin. americ. gr. vij ▽ Cichor. q. ſ. f. Emulſ.

Oleum expreſſ. etiam exhibetur ad quartanam vel per ſe, vel cum oleo expreſſo ricini medii.

[De ricino Americano conf. prolixe Sim. Paulli Quadr. Botan. Claſſ. Quart. p.m. 549. & ſeqq.]

CATAPUTIA MINOR.

Secundum quoſdam ſpecies eſt Ricini, ſed revera eſt ſpecies Tithymali. Sunt Lathyridis majoris C.B.P. grana oblonga, rotunda, ovalia, piſo majora, includentia ſub pellicula corticoſa nucleum album pinguem.

Surſum & deorſum ſerum potentiſſime evacuat, hinc velut eſula hydropicis robuſtioribus conducit. Folia ſeminalia etiam habent vim purgantem concentratam. Externe 🜖 expreſſum abdomini applicatum etiam alvum ſolvit. Datur in ſubſtantia à gr. v. ad vij & x.

℞. Sem. Catap. min. gr. vij Zinz. gr. v. contunde. fiant pil. in hydrope exhibendæ. Non opus eſt, ut aliquid addatur ad formandas pilulas, nam ipſa ſubſtantia ſeminum contuſa ſat viſcoſa eſt.

℞. Catap. min. gr. vij exhibe ex ovo ſorbili vel oxymell.

℞. Ung. d. arthanit. ℥ij. ☍ Catap. expr. ʒj. ☍ diacolycynth. ʒij M. f. ung. ad ſolvendum alvum.

SEMEN CNICI SEU CARTHAMI.

Optima transferuntur ex agris Argentoratenſibus. Sunt Cnici ſativi ſeu Carthami officinarum C.B.P. ſemina oblonga, anguloſa, lævia, alba, includentia ſub cortice duriuſculo pulpam albicantem, ſaporis dulcis poſtea acris & nauſeoſi.

Planta eſt capitata pappeſcens, floribus tinctores utuntur; multæ aviculæ veſcuntur his ſeminibus abſque purgatione, hinc quia vis languida ſatis, raro uſurpantur. Ob lentorem ſubdulcem cui vis purgans incluſa eſt conveniens eſt catharcticum ſeriducum tuſſientibus aliiſve vitiis Pulmonum laborantibus.

Datur in ſubſtantia, melius in Emulſione ad ʒij, in qua commode reſinæ purgantes ſolent exhiberi. Extractum datur ad ʒj Elect. diacarthami datur ad ʒvj.

℞. Sem. Cartham. ʒij. ▽ fœnicul. q. ſ. f. Emulſ. contra tuſſim & alvi adſtrictionem.

℞. Sem. Carthami ʒij Citr. ʒj aq. fumar. ℥iiij Cinam. ℥ß f. Emulſ. add. Mann. ℥ij.

℞. Extr. diacartham ʒß. diagr. gr. iij extr. troch. alhand. gr. j. ☍ d. fœnic. gt. ij f. bolus purgans.

BALANUS MIREPSICA SEU NUX BEEN.

[Sunt glandis unguentariæ C. B. P. fructus avellanæ magnitudine, triquetri, continentes ſub putamine corti-

corticoſo cinereo & pellicula candida fungoſa nucleum pinguem & oleoſum veluti amygdalæ, ſaporis ſubacris amari & oleoſi odoris nullius.]

Ob vim purgantem, quæ in 🜖 acri & amaro conſiſtit, referri quidem debet ad purgantia, ſed ob vim purgantem tardam ut plurimum ex eorum cenſu proſcribitur.

Hodie uſurpari ſolet 🜖 expreſſum pro menſtruo ad extrahendos florum odoratorum odores, nam neque facile rancorem contrahit, neque alio peregrino odore florum odores inficit.

℞. Vas vitreum in imo ſtrictius, in ſummo latius, in hoc mediante crate ex ſetis equinis facta, & circulo ligneo formata, ſtratificentur flores & goſſypium tenuiſſime carptum oleo balanino imbutum per quatuor horas, exprime ex goſſipio oleum, & novis floribus prædicto modo reaffunde, donec oleum ſufficienter ſit imprægnatum florum odore. [Conf. ulterius Joh. Zvvelffer. animadv. in Pharm. Auguſtan. p.331.]

℞. Ceræ albæ q. v. 🜖 balan. q. ſ. f. ungt. pro baſi omnium balſamorum, cui inſtilla olea deſtillata pro arbitrio. Cera alba in officinis plerumque adipe eſt adulterata, ideoque rancorem contrahit. Si vero non adulteratam habeamus, non ranceſcit, & præfertur 🜖 Nuc. Moſch. expreſſo pro balſamo conficiendo, nimis enim fragrans hoc exiſtit.

MYROBALANORUM
QUINQUE SPECIES.

Belliricæ. Sunt Myrobalani rotundæ belliricæ C. B. P. arboris pruniferæ indicæ folio laurino ſubcinericeo fructus rotundi, ſæpe anguloſi, magnitudine

majoris gallæ, in pedunculum craſſum deſinentes, ex rubigineo flaveſcentes, includentes oſſiculum durum ſub putamine carnoſo duriuſculo, ſaporis adſtringentis poſt ſubdulcis. Arabibus Belleregi dicitur.

Chebulæ. Sunt Myrobolani maximi oblongi anguloſi pituitam purgantes C.B.P. arboris folia perſicæ ſimilia, fructus oblongi, omnium maximi, rugoſi, coloris obſcuri fuſci. Includentes ſub putamine carnoſo oſſiculum oblongum, craſſum & cavernoſum. Arbor Quebuli dicitur.

Emblicæ. Sunt Myrobalani Emblicæ C.B.P. arboris palmæ magnitudine foliis minutim inciſis, fructus rotundi, ſexangulares, belliricis minores, ex cinereo nigricantes, includentes oſſiculum leve, pallidum, avellanæ minoris magnitudine hexagonum, tribus cellulis inſignitum, in quo nucleus pomaceus ſpadiceus triangularis. Arbor Embelli dicitur.

Citrinæ. Sunt Myrobalani teretes citrini bilem purgantes C.B.P. Arboris ſorbi foliis, fructus oblongi, rotundi quinque angulis rugoſi, chebulis minores, coloris citrini, includentes ſub putamine carnoſo oſſiculum anguloſum. Arabibus Azafar dicitur.

Indæ. Sunt Myrobalani nigræ octangulares C.B.P. arboris folio ſalicis, fructus oblongi reliquis graciliores & tenuiores, ſuperficie octangulari, ſolidi & duri intus & extus, piceæ nigredinis inſtar bituminis judaici. Arabibus Aſuar dicitur.

Conſtant ⊖e purgante blando, cum terrea ſubſtantia adſtringente. Purgativa vis conſiſtit in ſubſtantia pulpoſa, quæ ſimul aliquid reſinæ habet, adſtrictiva vero in ſubſtantia duriore corticoſa, hinc in dyſenteria aliisve alvi fluxibus, ubi ſimul purgatione & adſtrictione opus eſt conducunt. In-

Infuſum laxat alvum abſque adſtrictione, in ſubſtantia & decocto purgat & ſimul adſtringit, toſtorum pulvis potentius adſtringit, & nihil vel parum purgat, præ cæteris in uſum vocantur Chebulæ.

Dantur in infuſo ad ℥ij in decoct. ad ℥iß. in ſubſtantia ad ʒj.

℞. Chebul. ʒß. Rhabarb. ʒiß. f. p. pro ij doſ. in Lienteria, & Dyſenteria.

℞. Chebul. ℥ij paſſul min. ℥j rhabarb. ʒiß. infund. ſer. lactis, add. ſyr. d. Cich. c. rhab.

℞. Myrobal. toſt. rhab. toſt. NM. toſt. ana ℈j. L.O. gr. j. f. pulv. adſtringens in quolibet fluxu alvi.

FRUCTUS CASSIÆ.

Eſt Caſſiæ fiſtulæ Alexandrinæ C.B.P. ſiliqua oblonga, rotunda, cylindracea, nigricans, includens ſub putamine duriuſculo corticoſo in certis & ordine diſpoſitis loculamentis ſemina depreſſa, nitida, fulva & pulpam nigricantem melleam. Ægyptiaca eſt optima, corticem habet è nigro rufeſcentem. Pulpa ⊖ purgans blandum in ſubſtantia viſcoſa nonnihil dulceſcente occultat, purgat bilem & ſerum, cietque ſimul Urinam, unde gonorrhœa, ſtranguria & calculo vexatis, ſi alvi adſtrictione laborant, conducit. Quibuſdam tanquam ſingulare antipleuriticum laudatur. Sectis à calculo hujus uſus eſt interdictus. Cæterum ob viſcoſitatem torminoſa eſt & tardæ operationis, propterea correctorio & ſtimulo eget. Hanc ob rationem hypochondriacis non conceditur.

Interſtitia lignoſa poſſidere vim purgantem experientia ſimiarum conſtat, quæ hæc devorantes fortiter purgantur.

Fallopius & Fabricius vim contra ſtranguriam improbant facto experimento in pluribus.

Datur pulpa recenter ex fructibus extracta ab ℥j ad ij.

℞. Pulp. Caſſ. ℥j ſacch. cand. ℥ß. ſucc. liquir. ʒiß. cape ad vitia renum & expellendum ſabulum & præcavendam generationem calculi.

℞. Pulp. Caſſ. ℥j detur in Pleuritide.

TAMARINDI.

Sunt ſiliquæ Arabicæ Tamarindus dictæ C. B. P. fructuum pulpa ex fuſco nigricans, ſaporis ſubacidi raris per longitudinem lignoſis filamentis pertexta, diſtincta in membranacea loculamenta, in quibus incluſa latent oſſicula dura quadrata, ex fuſco rubicunda & ſplendentia.

Horum pulpa in ſubſtantia acido-vinoſa poſſidet vim laxantem blandam ad morem ♀ri, propterea appropriatiſſima eſt & tutiſſima, in æſtu reprimendo, præcipue febrili, ubi opus eſt ſimul laxante, ejus temperato acido bilis mulcetur & ſanguinis nimia efferveſcentia ſedatur, ſtomachus quoque roboratur, ejuſque fermentum reſtituitur. In nidoroſa cruditate cum rhabarbaro valde commendatur. In nimio hæmorrhoidum fluxu à ſanguine acriore & bilioſo laudatur à Mercato pro laxante: ♀us etiam ex horum pulpa paratur.

Datur ſubſtantia ab ℥j. ad ij vel cruda, vel ſoluta cum ▽ appropriata. Decoctum ad ℥iij. Pulpa ad ℥iß.

℞. Pulp. tamarind. ℥j. Rhabarb. ʒj f. bolus in cruditate nidoroſa.

℞. Pulp.

℞. Pulp. tamarind. ℥ß. Crem. ♁ri. ℈ij ſyr. d. rhabarb. ʒv. ∇ Cichor. q. ſ. pro diluenda pulpa M. f. potio purgans.

℞. Pulp. tamarind. ℥ij coq. in ſ. q. ∇æ. ℞. Colat. ℔j add. Tinct. roſar. violar. ana ʒj ſyr. rib. ſ. berber. ℥j f. julapium ad Linguæ ariditatem.

℞. Pulp. tamarind. ℥ij paſſul. min. ℥j coq. cum ſero lactis f. decoctum ad ſitim fallendam.

℞. Decoct. hord. Corn. Cerv. ana ℔j pulp. tamarind. ℥iiij f. potus in febribus malignis uſitatiſſimus. add. tincturam bezoardicam quamcunque vel ⌒um theriacalem camphoratum, vel aquam theriacalem camphoratam.

SEMEN ATRIPLICIS.

Sunt atriplicis hortenſis albæ, ſeu pallide virentis C. B. P. folliculi membranoſi orbiculati ex binis folliculis conflati, includentes ſemen unicum compreſſum orbiculatum ſpadiceum, ſaporis dum recens nidoroſi.

Sale nidoroſo emolliente & laxante pollent ſed debili, adeo ut vix in uſu ſint. Interim Matthiolus [l. 2. c. 112. p. m. 364.] ex Serapione vim purgantem emeticam ſatis validam his attribuit, quæ vis tamen intelligenda eſt de recentioribus. Quibuſdam commendantur in ictero, dum obtundunt & evacuant bilem.

℞. Sem. atriplic. ʒij propina cum ∇ calida, ad laxandam alvum.

℞. Sem. atripl. cannab. ana ℥j coq. in lacte, datur in ictero.

SEMEN

SEMEN SAMBUCI.

Sunt Sambuci fructu in umbella nigro C. B. P. acini exigui oblongi, plani, ſaporis ſubacris & nauſeoſi. Grana ſeu Semina habent vim blandam laxantem & Urinas moventem ſimulque in receſſu adſtringentem. Vis purgans conſiſtit in oleo, quod ex his exprimitur. Baccarum pulpa diaphoreſin potius quam Catharſin excitat.

℞. Sem. ſambuc. coq. cum juſculo gallinaceo add. rad. petroſel. q. v. f. apoz. contra hydropem.

℞. Pulp. gran. actes, farin. ſecalin. f. paſta, exhibe cum decocto C. C. contra pleuritidem. Quercetanus laudat quoque in dyſenteria.

℞. Olei expr. gran. actes ℥j propina ex Cereviſia butyracea ad movendum Vomitum hydropico, quod ut arcanum laudat Hoffmannus in Clave, dicitque 🝆 illud exprimi ex arillis baccarum poſt ſucci tranſcolationem in filtro relictis. Laudatur & ſequens hoc:

℞. Olei expr. gran. act. q. v. inunge ventriculi regioni pro laxanda alvo in hydrope.

* * *

SEMINA BLANDA VISCOSITATE CONSTANTIA.

SEMINA GOSSYPII SEU BOMBACIS.

Sunt Goſſypii fruteſcentis ſemine albo C. B. P. ſemina oblonga, rotunda, aliquantum compreſſa, magnitudine ciceris, quæ viſcida lanugine candida includuntur,

duntur, pericarpiis corticoſis in tres quaſi carinas dehiſcunt, ſaporis ſubdulcis, odoris nullius.

Planta in qua naſcitur goſſypium eſt flore malvæ vel altheæ.

Duæ ſunt ſpecies annua & perennis, annua adfertur ex India Occidentali, facile etiam hujus ſemina germinant in noſtris terris. Perennis ex oriente, tertia etiam datur ſpecies, quæ lanuginem ſericam profert, quæ ex plantis perennibus præſtantior.

Ratione ſubdulcis & blandi viſcoſi Pulmonibus dicata ſunt & acria quævis temperant, conveniunt in phthiſi tuſſi & raucedine.

℞. Sem. bombac. Card. Ben. Mar. ana q. v. c. ▽ puleg. q. ſ. f. Emulſ. in tuſſi vel ▽ fl. papav. contra Pleuritidem.

SEMEN HORDEI.

Viſcoſitate adeo blanda & temperata conſtat, ut tam pro medicamento quam alimento habeatur, tanquam anodynum & mulcens acria quævis, & ex his natos affectus, hinc in febribus ardentibus, in nimia ſanguinis effervesſcentia, in compeſcenda ſiti, aliiſque uſurpatur. Externe quoque emollit, maturat, & anodynum eſt.

Datur hordei emulſio, cremor, ſemina etiam mundata vel uſtulata, vel ad crepaturam cocta, datur etiam decoctum.

℞. Sem. hord. ad crep. coct ℥ij. ▽ Card. B. q. ſ. f. emulſ. temperans & refrigerans in febribus.

℞. Decoct. hord. ℔j. ▽ Cinam. ℥j ſyr. roſar. & violar. ana ℥ß. lap. prunell. ʒj f. Julap. temperans & refrigerans in febribus ad ſitim fallendam.

℞. Decoct. hord. ℔ij ſyr. limon. ℥ij. -⁀- ⊕l. q. ſ. f. Jul. refrigerans.

℞. Hord. ℥ij liquir. ℥ß. paſſul. min. ℥j M. f. decoct.

℞. Raſ. C.C. hord. ana ℥ij coq. ex ▽ ſ. q. add. ſyr. Card. Ben. f. decoctum temperans in Variolis & Morbillis, in quo & ſpecifica poſſunt exhiberi.

℞. Hord. mund. ℥ij rad. chin. ℥ß. coq. in ▽ f. apoz. in febribus ardentibus.

℞. Decoct. hord. ▽ ſperm. ranar. ana ℥ij ſem. melon. cucum. ana ʒj amygd. d. n. vij f. Emulſ. ad fluxum hæmorrhoidum nimium à ſanguine acriori, in alvo adſtricta poteſt addi Mannæ. ℥ß.

℞. Hord. ⊖ commun. ana Mj inde ſacculo & applica abdomini ad tormina & inflationes: de uſu farinæ hordei v. cap. de lino & fœnu-græco.

SEMEN CANNABIS.

Sunt grana ſubrotunda parva, quæ ſub tenui cortice ex cinereo ſpadiceo glabro includunt medullam albidam, ſaporis ſubdulcis.

Hujus mucago nonnihil olei conjunctum habens, acrem volatilemque bilem à qua icterus oritur mire temperat & compeſcit, obtinet quoque vim ſtupefactivam, quæ ex recentis plantæ odore gravi, idem exſiccatione manifeſte percipitur, orgaſmum genituræ maxime compeſcit, & propterea tanquam antaphrodiſiacum & ſpecificum contra gonorrhæam laudatur.

Turcæ præparant ſuum Maſlach ſeu Banque hoc modo. Recipiunt folia cannabis minoris recentiſſima, in pulverem terunt in olla, in qua butyrum fuit aſſervatum, torrent, madefaciunt, hocque aliquoties repetunt, & utuntur ſingulis diebus, quando ſibi ſtuporem inducere volunt.

In

In India orientali alia eſt planta ex qua fit Banque, Althæa goſſypii folio ſapore acetoſæ, terunt etiam in pulverem, miſcent cum butyro & oryza [deſcripta eſt hæc planta ab Hermanno noſtro Hort. Lugdun. Batav. p. 25. conf. etiam Garz. ab Horto l. 2. c. 25. p. 242 it. de Maſlach. Turcarum Wedel. opiolog. l. 2. ſ. 4. p. 1. p. 168.]

Datur in Emulſione, decocto. it. ☍ expreſſum.

M: ℞. Sem. cannab. ℥ij coque in lacte bubulo ad crepaturam uſque, cape hujus decocti ℥v bis vel ter in die, contra icterum ex acri & volatiliore bile.

℞. Sem. cannab. ℥ij ſem. lactuc. portulac. ana ʒij f. Emulſ. ex ▽ portulac. & Nymph. ſ. q. ad extinguendam ♀em, adde ſi præſtantiorem expetas ⦶ præpar. ℈ß. ſacch. ♄n. gr. vij. ſem. agn. caſt. ℈j.

℞. Sem. cannab ʒvj. 4. frig. maj. ana ʒj f. Emulſ. cum ▽ Menth. add. ſuccin. præpar. oſſ. ſepiæ ana ʒj ſyr. nymph. q. ſ. ad gonorrhæam.

Oleum expreſſum inungitur regioni renum contra gonorrhæam.

***

FRUCTUS ET SEMINA SUBDULCIBUS PARTICULIS ET OLEOSIS TEMPERATIS CONSTANTIA.

AMYGDALÆ DULCES.

Hi nuclei terræ commiſſi facile progerminant, & citiſſime in arbores excreſcunt, aliquando vero ex dulcibus fiunt arbores fructus amaros ferentes & vice verſa ex amaris, in meliorem terram tranſplantatis.

Ob vim alimentoſam conveniunt in phthiſi, hectica, tabe, atrophia, ob vim medicamentoſam conveniunt in omnibus affectibus acrioribus, mulcendo & obtundendo, tormina quoque tollunt.

Duplex parari poteſt 🜕, unum ex amygdalis parum crematis præloque commiſſis, quod optimum, ſi plane non cremarentur adhuc melius eſſet. Alterum fit evaporando emulſiones factas, tum 🜕 remanet, quod nullius momenti.

Ob oleum quo gaudent, fœces alvinas induratas emolliunt & alvum ſolvunt. Urinæ acredinem temperant, temperant quoque acidum, à quo ſæpe inflammationes internæ, pleuritis nempe & nephritis oriuntur.

Datur eorum ſubſtantia, emulſio & 🜕 expreſſum: oleum interne datur, quando recens eſt expreſſum, nam inveteratum rancidum eſt & nauſeoſum.

℞. Amygd. d. excort. ℥ij. aq. rub. id. q. ſ. aq. roſar. par. f. Emulſ. nutriens & confortans conveniens inprimis in phthiſi & tabe, add. Magiſt. perlar. ℈j it. aq. Cinam. tantill.

℞. Amygd. dulc. ℥ß ſem. pap. alb. ʒij 🜄 Nymph. q. ſ. f. Emulſ. add. Ebor. ʒß. ſyr. d. Nymph ℥ß. datur pro refrigerante & ſomnum inducente.

℞. Decoct. hord. aq. ſperm. ranar. ana ℥iij ſem. pap. alb. ʒij gland. immatur. num. ij amygd. d. n. v. f. Emulſ. add. Miv. Cydon. ʒvj Magiſt. Corall. ℈j Unicorn. foſſil. gr. xv. L. O. gr. j. exhibe contra fluxum hæmorrhoidum.

℞. Olei amygd. dulc. ℥iiß add. Mann. ℥iß cape ex juſculo gallinæ contra colicam, docente ita Sennerto.

℞. Olei amygd. dulc. ℥ij Ocul. ♋. ʒß. ſyr. dialth. par. cape ad dyſenteriam.

℞. Olei

℞. Olei amygd. ℥iß. Vin Malv. ℥ß. ſyr. papav. ℥ß. M. f hauſtus in colica paſſione, alii addunt ſaponem Venetum.

℞. Olei amygd. ℥vj. 🝆 laur. expr. ℥ij f. Clyſter in colica flatulenta. [De uſu olei amygdal. dulc. in forma clyſteris injecti conf. Sim. Paulli Quadr. Bot. Claſſ. 2. p.m. 18.]

AMYGDALÆ AMARÆ.

Ob ſui amarorem ad caliditatem inclinant, & minus temperatæ ſunt ac dulces, harum 🝆 expreſſum, in quo potior vis conſiſtit, diureſin movet, calculum & nephritidem curat, eſt quoque antipleuriticum. Convenit externe in tinnitu aurium & ſurditate à ſordibus auri impactis

An amygdalæ amaræ comeſtæ ante potum ob ſui oleoſitatem ebrietatem præcavent? experientia conſulatur.

Dantur per ſe, harum Emulſio, 🝆 expreſſum ut & elixir.

℞. Olei amyd. amar. ℥ij detur contra dolorem nephriticum, calculum & pleuritidem.

℞. Olei amyd. amar. q. v. adde ſacch. q. ſ. ad conſiſtentiam mellis, f. Electuar. applica ulceri vel vulneri faucium, œſophagi, quin & aurium.

℞. Amygd. amar. ℥ij. 🜅 vin. rectific. ℔j cinam. ℥ß. digere, deſtilla ex cineribus: cape ℥ß. vel ʒij ad flatulentiam & pleuritidem.

[Amygdalæ amaræ lethales ſunt vulpibus, gallinis, ſciuris &c. vid. C.N.C. Dec. I. ann. 8. obſ. 99. p. 183. & ſqq. it. Dec. 2. ann. 8. obſ. 154. p. 335.]

PISTACIA.

Sunt Piſtaciæ peregrinæ fructu racemoſo ſeu Therebinthi Indicæ Theophraſti C. B. P. fructus oblongi, anguloſi avellanæ magnitudine, altera parte elatiores, altera depreſſiores, coſta notati, ſub putamine tenui includitur nucleus ex albo vireſcens, ſaporis oleoſi ſubdulcis.

Eſt vel major vel minor, ex Perſia raro transfertur & multo ſapidior eſt quam major, in noſtris creſcens etiam terris.

Nuclei qui ſoli in uſu ſunt conveniunt cum amygdalis dulcibus, ſed potentius augent genituram, vis omnis conſiſtit in oleo recenter expreſſo, datur ut amygdalæ.

℞. Olei piſtac. expreſſ. ℥iß vel ij. vitell. ovor. n. j. ambr. gryſ. ℈ß. ſacch. cand. ʒij redige ad ignem cinerum in formam Electuarii, propina ſeni ad augendam ♀em.

℞. Piſtac. rec. amygd. dulc. nucl. pini f: Emulſ. ex lacte contra phthiſin, tabem & maraſmum, add. ſem. Eruc. ℈ß mac. gr. v. Cubeb. gr. vij f. Elect. aphrodiſiacum.

NUCLEI PINI.

Sunt Pini ſativæ C.B.P. nuclei oblongi, candidi, pingues, incluſi mediante rutila tunica oſſiculis craſſis & duris, in turbinato & ſquamato cono diſpoſitis, ſaporis ſubdulcis.

Virtute piſtaceis accedunt, ſed ſpecifice commendantur in temperanda lympha acriore, ſtillante in

in pectus, ut & in mulcendo acido ſtranguriæ & dyſuriæ. Vis omnis conſiſtit in oleo.

℞. Piſtac. amygd. d. pinearum ana ℥ß. ſem. eruc. urtic. ana ℈j ſpec. diambr. ʒß Mell. q. ſ. f. conditum aphrodiſiacum, add. diamoſch. ℈ij Elect. diaſatyr. ℥ij.

℞. Olei rec. expr. nucl. pini ℥j cape cum vino ad erigendam ♀em vel etiam ☍ Majoran. gt. iij.

NUCLEI CERASORUM.

In horum ſubſtantia mucaginoſa & ☍ſa latet aliquid ſalis diuretici, in oleoſa vero virtus anodyna, unde ſingulariter laudantur in nephritide.

℞. Nucl. Ceraſ. ʒiij ▽ chæref. q. ſ. f. Emulſ. diuretica.

℞. Olei expr. nucl. ceraſ. ʒij exhib. ex ▽ chæref. contr. nephritidem.

NUCLEI PERSICORUM.

Cum nucleis ceraſorum conveniunt, ſed amariores, acriores & potentiores ſunt in pellenda Urina.

Oleum expreſſum & infuſum propriis floribus ſingulare eſt in ſurditate, tinnitu & eliciendis vermibus ex aure. Dantur ut nuclei ceraſorum.

℞. Nucl. Perſicor. ʒij. ▽ gram. q. ſ. f. Emulſ.

℞. Nucl. Perſicor. ʒvj ſem. melon. cucum. ana ʒij papav. alb. ℥ß. ▽ veron. tuſſilag. farfar. ana ℥iij f. Emulſ. in phthiſi.

℞. Nucl.

℞. Nucl. Perſic. ▽ Nymph. verben. acet. q. ſ. f. Epith. ad capitis dolorem.

ANACARDIUM.

Sunt arboris Malabaricæ Oepata dictæ, fructus compreſſus aviculæ cordis figura, nigricans, includens ſub duplici involucro, exteriore tenui rufeſcente, nucleum ſolidum albicantem, ſaporis ſubdulcis & intra illud duplex involucrum quaſi in poris favis apum ſimilibus, liquorem melleum glutinoſum rufum, ſaporis acris & exurentis, qui ab Indis habetur venenatus, à Chirurgis ſcirrhis etiam duriſſimis impoſitus eos miraculi inſtar diſſipat.

Hujus nuclei piſtaciorum vices ſupplere poſſunt. Toſti horum nuclei ſub cineribus ab Indis avide comeduntur.

Veteres integro fructui adſcripſere vim memoriam confortandi, & in hunc uſum proſtat, an vis cerebrum confortandi conſiſtat in nucleis his innocentibus, an in muco oleoſo acerrimæ exedentis & volatilis Naturæ, qui intra corticis duplicaturam reperitur, aliis judicandum relinquo.

Hoc mucoſo acri oleo utuntur Indi ad abſumenda condylomata, ficus, caruncul as, ſcrophulas aliaſque excreſcentias carnoſas.

℞. Confect. anacard ℈j cape pro confortando cerebro juſto humidiore.

CACAO.

CACAO.

Sponte ſua naſcitur in India occidentali non orientali, ut quidam perperam ſcripſere.

Sunt arboris Cacaiferæ Americanæ nuclei, amygdalis magnitudine forma & conſiſtentia ſimiles, incluſi copioſo numero in involucro corticoſo, duro, rugoſo, fuſco oblongo peponem imitante. Saporis ſubpinguis oleoſi & ſubamari, odoris nullius.

Virtutem amygdalarum pinearum & piſtaciarum æmulantur, vim habent nutriendi potentem adeo, ut ex quorundam ſententia carnibus omnibuſque aliis alimentis ſint anteponendi. Vis omnis conſiſtit in mucoſa blanda & oleoſa ſubſtantia, qua totus fructus ſcatet.

Si ex retorta debite deſtilletur ſuppetit phlegma inſtar juſculi carnis, cujus ſpiritus & oleum redolet pinguedinem ex aſſata carne deſtillantem. Hinc emaciatis tabidis & qui pingueſcere cupiunt valde convenit. Datur uſtulatum in emulſione cum ſaccharo, lacte ſeu vino, additis vitellis ovorum, & ſi lubet aromatibus.

Indi & nunc Europæi ex hoc componunt maſſam Chocolatam dictam, quam in lacte, aqua & etiam vino fervente diſſolvunt, eamque ſorbillant.

℞. Cacao. toſti ℔j. farin. tritic. turcic. ℥vj. ſacchar. ℔ß. ſolut. in ▽ roſac. contund. invicem & agitetur ſuper ignem continuo ne aduratur, & redigatur in maſſam.

Modus quomodo in Hiſpania à ditioribus conficitur. ℞. Cacao excortic. mundi & toſti ℔vj Cinam. ℔ß Vainigliæ n. vij in ſyrupo ſolut. caryoph. arom.

n. vj. vel vij. farin. frumenti indici ℔iß piper. hiſpanic. ℥j orleanæ, ſuppetit colorem rubellum in ▽ roſac. ſeu ſyr. ſolut. ℥ij ſacchar. q. v. circiter ℔iij vel iiij contunde & miſce in cacabo ſuper △e leni, agita continuo, ac omnia exacte invicem mixta ſint, redige in maſſam & ſub finem adde eſſent. ambr. & moſch. ſi velis.

***

FRUCTUS PARTICULIS DULCIBUS VISCOSIS CONSTANTES.

DACTYLI.

Sunt Palmæ majoris C. B. P. dactyliferæ fructus oblongo-rotundi, glandium magnitudine & forma ſed paulo majores, includentes ſed pellicula fulva & tenui pulpam melleam ſubdulcem, & ſub hac nucleum oblongum durum, per longitudinem ſulco notatum. Ob vim mulcentem & temperantem acria quævis eorum caro commendatur in tuſſi, aſperitate faucium, raucedine vocis, & catarrhis ab acriore lympha oriundis, ut & in dyſuria & ardore Urinæ tanquam ſpecificum. Nucleus terreis conſtat, & in dyſuria pro acido imbibendo datur.

Præſcribuntur ut ficus & paſſulæ.

℞. Carn. dactylor. ficu. ana n. xx. paſſul. min. ℥ij rad. helen. ʒ. x. f. c. ▽ decoct. add. ſyr. violar. q. ſ.

FICUS ſeu CARICÆ.

[Sunt ficus communis C. B. P. fructus pyriformes, figuræ p. m. rotundæ ac depreſſæ, coloris ex flavo fuſci, intus copioſiſſimis granulis per pulpam infarcti,

farcti, ſaporis dulcis, odoris debilis non ingrati.] Variæ ſunt ſpecies & tot ferme ficus quot pyrorum.

Acria quævis mulcent potius quam dactyli, hinc in tuſſi, aſthmate, raucedine, nephritide, ardore urinæ, mictu cruento & iſchuria, mire commendantur. Urinam adeo potenter pellere dicuntur, ut tam eorum decoctum quam arilli manifeſtè à quibuſdam conſpecti fuerint. In ſpecie temperant acidum Variolarum & morbillorum, temperando ſanguinis ebullitionem & præcavendo ne acidum profundas foveas faciat.

Externe impoſiti maturant, emolliunt, leniunt & tollunt tumores gingivarum, uvulæ & faucium ut & horum omnium inflammationem. Nimium eorum uſum generare pediculos experientia tam veterum, quam recentiorum confirmat. [Conf. Sebiz. d. alim. fac. l.5. probl. 169. p.m. 1538. cujus ſententiam & noſtram facimus.]

Datur in ſubſtantia, decocto, gargariſmate &c.

℞. Hord. mund. paſſul. min. ana ℥j. liquir. ʒij ficu. n.vj. herb. capill. ♀is, hyſſop. ana Mß. ſem. braſſic. ʒij pin. recent. ℥ß. f. c. ∇ dec. ad raucedinem.

℞. Ficus q.v. macera in ∽u vini aniſato, deflagra, cape reſiduum pro tuſſi, raucedine & aſperitate.

℞. Ficus n. xv. fol. millefol. ſummit. hyperic. ana Mj coq. in ∇ contra mictum cruentum.

Cum ſucceſſu adhibetur in bubonibus ♀eis pulpa ficuum cum ☿io maritata, loco emplaſtri de ranis cum ☿io.

℞. Caric. ping. n. iiij vel v. herb. bellid. Mj coq. in ∇ add. tantill. ⊖is prunellæ, f. gargariſma contra inflammationes quævis oris.

JUJUBÆ ſeu ZIZIPHI.

Sunt ſpecies prunorum, deferuntur ex Ægypto, Hiſpania, Mauritania. Sunt jujubæ majoris oblongæ C.B.P. pruna ſeu fructus olivæ magnitudine & forma, puniceí ſeu rutili, involventes ſub pellicula craſſiuſcula oſſiculum oblongum, utrinque acuminatum, incluſum pulpæ molli fungoſæ, ſaporis mellei & vinoſi.

Symbolizant cum dactylis & ficubus, ingrediunturque decocta pectoralia & nephritica.

℞. Flor. Malv. Mj. jujub. ſebeſt. ana n. xv. coq. in ▽ ad ardorem Urinæ.

℞. Syr. jujub. violar. liquir. ana ℥ß. ſyr. pap. ʒij f. linctus contra tuſſim.

SEBESTEN.

Sunt Sebeſtenæ domeſticæ C. B. P. fructus prunis domeſticis ſimiles, nigricantes in plicas ſive rugas contracti, includentes in ſubſtantia carnoſa mellea oſſiculum tribus angulis vel coſtis notatum, in quo nucleus conditur exilis.

Virtute cum jujubis conveniunt.

℞. Elect. diaſebeſten. ℥ß. cape ad mulcendos quoſvis æſtus ſanguinis

PRUNA DAMASCENA.

Sunt pruna magna dulcia atrocœrulea C.B.P. recentia ſiccis longe præferenda, præter virtutes mulcentes & temperantes quas cum ſuperioribus, ſebeſten & ficubus habent communes laxant ſimul alvum, & cum facile ſint fermenteſcibilia, excitant quoque fermenta-

tionem in ſanguine, unde nimis comeſta dyſenteriam & alvi fluxus cauſant.

Viſcoſitas eorum quæ aliquando juſto diutius inteſtinis adhæret & tormina facit, corrigitur cinamomo, aniſo, fœniculo.

Datur horum decoctum, pruna laxativa, electuarium, pulpa.

℞. Prunor. damaſc. paſſul. major. ana ℥j coq. in 🜄 add. fol. ſenn. ℥j cinam. ℥ß. ſem. fœnic. ʒiij ad laxand. alvum pro aliquot doſibus.

℞. Prun. damaſc. ℥iiij coq. in 🜄 ſimpl. colatur. add. fol. ſenn. ʒvj Crem. 🜍ri ʒij cinam. ʒij mane ebulliant leniter, colaturam reaffunde prioribus prunis, ſerva in vaſe ut humiditas paulatim exhalet, dantur pro laxante pruna v. vj. vel etiam plura.

℞. Prun. damaſcen. n.xv. coq. in 🜄 ad ℔j. colatur. infund. fol. ſenn. ℥j ſem. fænicul ℥ß. flor. bugloſſ. violar. ana p. j. add. Mannæ ℥ij ſicca leniter ſuper 🜂 cape pruna vij vel plura pro laxante.

℞. Elect. diaprun. ℥ß. ad alvum laxand. add. diagryd. gr.v. pro ſtimulo & troch. alhand. gr. iij ad purgandum.

PASSULÆ MAJORES ET MINORES ſive CORINTHIACÆ.

Eſt uva paſſa major βότρυς Græcis C. B. P. ſunt baccæ exſiccatæ depreſſæ, rugoſæ ex obſcuro rufeſcentes, continentes ſub pellicula tenui acinos anguloſos, involutos pulpæ melleæ ſubrufæ.

Harum ſubſtantiam temperatam ſubacrem, ſubdulcem lenientem & alvum blande laxantem conſtare in particulis 🜔inovolatilibus & 🜍uoſis docet vi-

num, quod ex fermentatis conficitur ut & earum fermenteſcibilitas.

Ob ſubſtantiam viſcoſam ſubdulcem & temperatam emolliunt, leniunt, temperant, & propterea in tuſſi, raucedine, aſperitate conveniunt & partibus nutriendis alimentum apponunt, quem in finem decoctis infuſis acrioribus, tanquam temperans adjunguntur, ob ſui fermenteſcibilitatem alvum ſolvunt, eamque apertam detinent, ut proinde in alvi retentione & fœcibus induratis mire proſint. Ob ſalia vero volatilia aperiunt & deobſtruunt, adeoque ſcorbuticis, cachecticis & hypochondriacis ſingulari proprietate conducunt.

Datur earum ſubſtantia, pulpa, infuſum decoctum.

Minores ſunt uvæ paſſæ minores ſive paſſulæ Corinthiaæ C.B.P. quæ præferuntur majoribus. Si majores præſcribuntur ſignentur paſſulæ majores exacinatæ, quia in acinis conſiſtit vis adſtrictoria. [Acinos exemptos & toſtos in pulverem redigit ſubtiliſſimum & à ℈j ad ʒj alvi fluxu laborantibus cum ἀφοϱία adhibet propter ſingularem vim ſtegnoticam Joh. Ægidius Euth. C.N.C. Dec. III. ann. 7. & 8. obſ. 5. p.m. 8. imo ex iis beneficio -🜍-us vini rectificati tincturam extrahit ſaturate tinctam quæ mirificam cum marte vel terra Japonica habet analogiam.]

℞. Paſſul. cum pyris moſchatellin. q. v. contunde, ſine ſuccum effervescere ut fiat vinum, quo in cachexia utere, ſeu loco pyrorum cape poma redolentia contra hypochondriacam paſſionem.

℞. Pulp. paſſul. ℔ß rhabarb. ʒij ſacchar. ℥ij M. cape cochl. j ante menſam ad dyſcraſiam ſanguinis, aromatiſa etiam cinamomo.

℞. Pulp.

℞. Pulp. paſſul. tamarind. ana ℥j cape in ſanguinis æſtu cum adſtrictione alvi.

℞. Paſſul. min. ℥ij coq. in ▽ ſimpl. add. colatur. adhuc ferventi fol. ſenn. ℥ß. Crem. ♀ri ʒiß. f. laxans

℞. Pulp. paſſul. min. ſyr. d. cich. c. rh. ana ʒij corall. r. Oc. ♋ ana ℈ß M. datur infantibus ad expurgandum meconium, etiam in febribus in alvo adſtricta.

℞. Pulp. paſſular. cape c. ▭u vini aniſato contra tuſſim & raucedinem.

℞. Paſſular. amygd. dulc. ana q. v. cape ſingulis diebus pro corpore pinguefaciendo.

℞. Paſſul. minor. hord. mund. ana ℥j rad. liquirit. ʒij caric. ping. n. vj. herb. capil. ♀is Mj. pinear. recent. ʒj M. f. decoct. ad tuſſim & raucedinem.

SILIQUA DULCIS ſeu CERATONIA.

Sunt Siliquæ edulis C.B.P. fructus oblongi, compreſſi, paululum intorti, craſſi, digito longiores, pollice latiores, fuſci ſplendentes, dorſo elatiore includentes pulpam carnoſam, rufam, dulcem, ſub qua intra membranaceos loculos latitant nuclei plani, compreſſi, orbiculati, lucidi, fuſci. Habent dulcedinem majorem in pulpa, quam in cortice. Hujus dulcis & viſcoſa ſubſtantia in ſpecie temperat ſodam ardorem Ventriculi & urinæ. Datur integra cum pulpa & cortice.

℞. Siliq. dulcis ʒj. cape cum ▽ Nymph. contra ſodam & ardorem ventriculi.

SEMINA

*
* *

SEMINA PARTICULIS AQUOSIS ET CRASSIS CUM OLEO PAUCO CONSTANTIA.

SEMINA QUATUOR FRIGIDA MAJORA.

Sunt Citrulli, melonum, cucurbitæ, cucumeris. Vocantur majora, quia majorem habent vim refrigerantem.

Citrulli ſemen. Sunt Anguriæ Citrullus dictæ C.B.P. Semina oblonga plana, rhomboidea, ad margines æqualia abſque eminentia, modo nigra, modo nigricantia, ſub cortice duriuſculo includentia medullam albam, ſaporis aquei.

Planta eſt pomifera ſcandens monopetala.

Melonum ſunt ſemina oblonga ex flavo albicantia, ad ſemen cucumeris accedentia.

Cucurbitæ ſunt ſemina thyroidea, oblonga, plana pallida, parte ſuperiore cornuta, ſeu auriculata duabus apophyſibus quaſi anguloſis eminentibus, & duabus lineis per longitudinem excurrentibus notata.

Cucumeris ſunt ſemina oblonga, plana, in medio paululum tumida ſeu convexa, candicantia, ſeminibus melonum anguſtiora, puſillo ſubtus apiculo protenſo.

Semina hæc, inter quæ præſtantiora ſunt ſemina melonum, conſtant particulis viſcoſis aquoſis craſſis, cum blando & pauco ☍, propterea temperant & obtundunt ⊖ia acria quævis, eaque ad renes præcipitant & per urinam expellunt, hoc nomine proſunt etiam

in

in ſupprimendo orgaſmo genituræ conveniunt, etiam in linguæ ariditate, & in mulcendis doloribus inteſtinorum à bile acri. Dantur horum pulpæ excorticatæ vel per ſe, vel in Emulſione.

℞. Sem. 4. frig. maj. ana ʒij amygd. dulc. ℥j decoct. hord. ℥vj ▽ roſar. ℥j. f. Emulſ. temperans & refrigerans, add. add. ſyr. papav. alb. q.v.

℞. Sem. 4. frig. maj. ana ʒij ſem. cannab. ʒij f. Emulſ. ex ▽ lactuc nymph. portulac. q. ſ. add. ⦶tr. præpar. ℈j. ſacch. ♄ni ℈ß. contra nimiam ſalacitatem.

Semina Citrulli ſpecifice in menſibus decoloratis laudantur ſi xij excorticata ſimul deglutiantur & hauſtus lactis ſuperbibatur.

SEMINA QUATUOR FRIGIDA MINORA.

Sunt Lactucæ, Scariolæ, Portulacæ, Plantaginis.

Lactucæ ſativæ C. B. P. ſunt ſemina parva oblonga, compreſſa, pallida, utrinque acuminata, ſaporis aquei.

Scariolæ ſunt Intybi ſativæ anguſtifoliæ C. B. P. ſemina Cichorii ſeminibus ſimillima, colore pallidiora.

Portulacæ latifoliæ ſeu ſativæ C.B.P. ſunt ſemina exigua nigra.

Plantaginis latifoliæ ſinuatæ C. B. P. ſunt ſemina parva oblonga, puliciformia, coloris obſoleti ac ferruginei.

Endiviæ vulgaris C. B. P. ſunt ſemina parva oblonga anguloſa ſpadicea, in ſummo acetabulo inſtar coronæ notata.

Coincidunt cum ſeminibus frigidis majoribus, excellunt vero ſemina lactucæ, ſemen vero plantaginis habet ſimul virtutem adſtringentem, ut proinde forſitan rectius ad terrea quam ad aquoſa amandari poſſit. Dantur in Emulſ. ut frigida majora.

℞. Sem. frig. min. ana ʒij f. Emulſ. ex decoct. hord. add. ſyr. pap. alb. q. ſ.

℞. Sem. plantag. portulac. corall. præpar. borrac. calcinat. ana ℈j. ſacch. ♄n. gr. v. L. O. gr. ij f. pulv. ad ij doſes, exhibendus in nimiis alvi & menſium fluxibus.

*
* *

SEMINA ET FRUCTUS PARTICULIS CRASSIS AQUOSIS SEU MUCAGINOSIS ABSQUE OLEO SE TAM SUBITO MANIFESTANTE CONSTANTIA.

SEMEN ALTHEÆ.

Sunt Altheæ Dioſcoridis & Plinii C.B.P. ſeu Malvæ paluſtris ſemina ſemicircularia ſemilunam exprimentia, ex dorſo craſſiori in acutiorem aciem deſinentia, ſaporis viſcoſi. Sua mucaginoſitate conſpirant cum radice, niſi quod mucago ſeminum ſit paulo tenuior.

℞. Sem. alth. ʒß. ſacch. ℈j f. pulv. in dyſuria, ſi potentiorem velis, add. Oc.♋. ℈j ⊖ ſtip. fabar. ℈ß.

℞. Sem. alth. ʒij. ▽ roſar. q. v. f. Emulſ. ſ. mucago ad dolores ex ambuſtione ortos.

SEMEN MALVÆ.

Sunt Malvæ folio ſinuato & rotundo C.B.P. ſemina in rotulæ aut caſeoli formam circumacta &c. inferiora ſunt & debiliora ſemine althææ,

℞. Sem.

℞. Sem. Malv. Alth. ana ℥ij. ▽ chærefol. f. Emulſ. vel mucilag. diuretica & ſimul anodyna.

SEMEN AQUILEGIÆ.

Sunt aquilegiæ ſylveſtris & hortenſis ſimpl. C.B-P. ſemina nigra lucida. In mulcente ſua mucagine habent aliquid ſalis, quod ſimul movet blandos ſudores, hinc in acido Variolarum & Morbillorum expellendo, & in nimia ſanguinis efferveſcentia tollenda laudantur.

℞. Sem. aquileg. ℨj naſturt. ℨß. ▽ Card. Bened. f. Emulſ. temperans fervorem ſanguinis in variolis & morbillis.

SEMEN BARDANÆ.

Sunt Lappæ majoris C.B. P. ſemina oblonga, ſtriata, fuſca craſſiuſcula, ſaporis ſubamari, habent partes mucilaginoſas includentes ſalem alcalinum.

℞. Sem. bardan. violar. ana ℥ij f. ex ▽ gramin. petroſel. Emulſ. diuretica.

FOENUM GRÆCUM.

Sunt Fœnugræci ſativi C.B.P. ſemina oblonga flaveſcentia ſeſamacea, compreſſa, anguloſa, ſaporis mucilaginoſi odoris jucundi, cum quadam gravedine.

Horum mucago in farinacea ſubſtantia conſiſtens, includit quidpiam ⊖is volatilis, quod odor indicat, unde non tantum emolliunt & leniunt ſed etiam diſcutiunt incidunt. Horum uſus potior externus eſt in cataplaſmatibus emollientibus & maturantibus, ut & in clyſteribus anodynis, dantur etiam in fotibus, in

 ſeſſibus.

ſeſſibus. Uſurpatur farina, mucago, ipſa ſubſtantia, rarius interne, frequentius externe datur.

℞. Fœn. græc. ʒiij ferment. ℥j ſterc. columb. ʒij. ☍ chamom. ℥iß. Mell. ℥j contund. miſc. f. catapl. in maturandis abſceſſibus.

℞. Fœn. græc. ſem. lin. herb. chamom. chæref. coq. c. ▽ ad form. Catapl. ad tumorem lienis.

℞. Mucag. ſem. lin. fœn. græc. alth. ☍ amygd. d. lilior. alb. ana q.v. applica tumoribus emolliendis.

℞. Sem. fœn. græc. pſyll. ana ʒij flor. verbaſc. Mj. roſar. pallid. Mß. lumbric. terr. ℥ß. coq. omnia c. lacte f. clyſter. in teneſmo add. ſebi cervin. ℥j.

SEMEN OROBI.

Sunt Orobi ſiliquis articulatis ſemine majore C. B. P. ſemina ſeu grana anguloſa, ſubrotunda, fuſca, ſpadicea, ſaporis amaricantis & ingrati paululum.

Ob farinaceam mucaginem reſpondet fœnu græco in emolliendis & maturandis abſceſſibus, ſed quoniam more omnium leguminum gaudet ⊖e diuretico ideo medetur Urinæ & calculo pellendo. Datur ut fœnum græcum.

℞. Sem. Orob. Cicer. ana ℥ij rad. gramin. ℥j. f. decoct. cui add. ſyr. dialth. Fern. q. ſ. contra calculum.

SEMEN CICERIS.

Eſt vel rubrum vel album, rubrum officinale, album culinarium. Cicer rubrum ſunt Ciceris ſativi C. B. P. ſemina rotunda anguloſa, piſo majora, unico angulo acuto prædita, coloris ex fuſco rubentis, saporis viſcoſi & leguminoſi.

Symbo-

Symbolizant cum ſemine orobi, aſt temperatiora ſunt & gratiora, provocant urinam, menſes lochia & fœtum. Dantur ut ſemina fœnugræci.

℞. Sem. Cicer. q. v. coq. cum brodio vel juſculo carnis ad menſes & urinam provocandam ut & ad lochia expellenda.

℞. Sem. Cicer. ℥ß. ſtipit. fabar. cremat. Mj rad. gramin. petroſel. ana ℥ij f. decoct. diuretic. cui add. ſyr. dialth. Fern.

℞. Farin. Cicer. lupin. fabar. ana ℥ij f. Catapl. in tumoribus duris leniendis.

℞. Cicer. rubr. ℥j flor. verbaſc. Mij coq. in ▽ colatur. add. ſevi hircin. ℥ß. f. Clyſter. in teneſmo.

SEMEN LUPINI.

Sunt Lupini ſativi flore albo C.B.P. ſemina orbicularia plana, compreſſa, ſaporis leguminoſi ingrati.

Coincidunt cum orobo & ob ſui amaritiem lumbricos pellunt, tum interne in decocto data, tum externe. Datur decoctum & farina.

℞. Farin lupinor. 🜕 amygd. amar. applica umbilico contra vermes.

℞. Sem. lupinor. ℥j f. dec. contra vermes.

SEMEN FABARUM.

Sunt Fabæ vulgaris C.B.P. ſemina oblonga, elata, ovalia, coloris varii, cum ciceribus conveniunt, tum ratione virtutis diureticæ, tum emollientis, uſurpantur ut cicera, proſtant etiam ſtipites fabarum uſtulati, ut & ⊖ è ſtipitibus paratum.

Farina eſt abſterſoria, unde Coſmeticum. Exterior cortex fabarum habet vim adſtrictoriam.

℞. Farin. Fabar. ℥iij ſem. Cumin. ℥ß. f. Catapl. ad inflammationem ſcroti & teſtium, add. flor. cham. Mj.

℞. ⊖ ſtipit. fabar. gr. vij Oc. ♋. ℈j. M. f. pulv. diureticus.

SEMEN LINI.

Sunt Lini ſativi C.B.P. ſemina oblongo-rotunda compreſſa, lubrica lucida, in medio tumida, ex fuſco punicea.

Eſt planta multicapſularis non leguminoſa.

Ex mucagine farina & oleo conſtructa ſunt, unde leniunt, molliunt & ſuppurant.

Oleum recenter expreſſum per os ſumtum ſpecificum eſt in Pleuritide & tuſſi pro juvanda expectoratione, per clyſteres injectum in fœcibus alvi induratis, à quibus colica paſſio oritur. Oleum lini cum terra japonica & ſigillata exhibitum in dyſenteria arcanum. Externe applicatum valet ad ambuſta & dolores hæmorrhoidum. Oleum lini cum ♆a deſtillatum mire medetur Paralyſi.

Mucago & farina applicantur tumoribus emolliendis & abſceſſibus maturandis.

Decoctum ſeminis vim habet ſingularem ad fœtum mortuum expellendum [conf. Dav. Herlicius de cura gravidar. c. XI. p. 57. & ex eo Sim. Paulli Quadr. Bot. p.m. 77.]

Datur mucago & oleum.

℞. ☍ lini rec. expr. ℥iij amygd. dulc. ℥j cape ex juſculo ad pleuritidem.

℞. ☍ lini ℥iij dent. apri præpar. ʒj cape ex ▽ chærefol. calide ad pleuritidem.

℞. ☍ lini ℥vij infund. troch. alhand. ʒß. f. clyſter. injiciendus ad fœces emolliendas.

℞. ☍

℞. 🝆 lini, rapar. ana ℥iiij f. Clyſter. ad iliacam paſſionem.

SEMEN HYPERICI.

Sunt Hyperici vulgaris C.B.P. ſemina exigua, oblonga ex fuſco nigricantia, ſaporis ſubamari & reſinoſi, odoris picei. Planta eſt tricapſularis polyſperma.

Ob reſinam cui ⊖ blandum diureticum conjunctum convenit in calculo, ut & in vulneribus tam internis quam externis. Specifice laudatur in mania, melancholia, delirio atque in vermibus, unde ſi herbam inſpergis caſeo, omnes vermes exiliunt.

Datur ſubſtantia, eſſentia ſeu tinctura, 🝆 dſat. decoct. & balſam.

℞. Sem. hyperic. ʒj ſacch. q. ſ. f. conſerv. l. a. contra nephritidem.

℞. Sem. hyperic. ʒj conſerv. roſar. fl. malv. ana ʒj f. bolus diuret.

℞. Sem hyperic. ʒj conſerv. fl. nymph. ʒij f. bolus contra maniam & deliria.

℞. Sem. hyperic. affund. 🜁 vin. q. v. f. tinctura Vulneraria, vel affund. 🝆 thereb. 🜁uoſ. l. Balſ. 🜍is & fac balſamum vulnerarium.

Quidam ad vulnera conſolidanda & cicatriſanda capiebat Hyperic. recent. Mij affundebat vinum, idem vinum bis vel ter ſeminibus affundebat, & ſic deſtillabat, uſque dum 🜁us vini prodiens per alembicum totus reſinoſus adparebat, hoc penicilla tingebat & vulneribus applicabat.

SEMINA

* * *

SEMINA MUCAGINE VOLATILI PRÆDITA.

SEMEN HYOSCYAMI.

Sunt Hyoſcyami albi majoris C. B. P. ſemina exigua rotunda, compreſſa, ex cinereo ſpadicea, ſaporis aliquantum viſcoſi, odoris narcotici & ingrati. Semen toſtum ad △em omni virtute anodyna expers eſt, unde probatur in volatilitate illam conſiſtere.

Horum oleum craſſum quod per expreſſionem elicitur, includit quoque volatilem quandam mucaginem ♌us involventem eorumque ferociam ſiſtentem velut opium, unde inter anodyna & hypnotica refertur.

Suffitus tetri odoris eſt & vermibus exoſus, ſi dentibus cavis & auribus excipiatur elicit vermes. Cave tamen ne nimis utaris ſuffitu vel ſeminibus ipſis nam ſtuporem & amentiam inducunt.

℞. Sem. hyoſc. ℥j infund. vin. hiſpan. relinq. per 24 horas, cape hora j ante decubitum, ad ſomnum conciliandum.

℞. Sem. hyoſc. ℈j conſerv. fl. violar. vel pap. rh. ℥j. f. bolus ad ſputum ſanguinis.

℞. Sem. hyoſc. papap. ana ʒß. Coriandr. ʒj thur. gr. ij maſtich. gr. iiij hirund. præpar. ℈ß. ſacch. ʒiiij f. pulv. ad ſput. ſanguinis. doſ. ad ʒj.

℞. ☍ hyoſc. expr. ℈ß. extract. verben. ʒj f. liniment. ad cephalalgiam.

℞. ☍ hyoſc. gt. vij nucl. perſic. ℈ij ungt. alabaſtr. ʒj f. linim. ad leniendos dolores capitis.

℞. ☍

℞. ☍ hyoſc. inunge inguina & lumbos ad gonorrhæam & menſium nimium fluxum, inunge tempora ad compeſcendum ſanguinem.

SEMEN PÆONIÆ.

Sunt Pæoniæ folio nigricante ſplendido maris C.B.P. ſemina ſeu grana oblongo rotunda, nigra, lucida, ſub cortice craſſiuſculo membranaceo includentia pulpam albam ſaporis ſubdulcis, odoris nullius.

Emulſiones Epilepticæ cum aquis appropriatis à nonnullis inde parantur.

SEMEN PAPAVERIS.

Sunt papaveris hortenſis ſemine albo ſeu ſativi C. B. P. ſemina minuta rotunda, rugoſa, candicantia, ſaporis ſubdulcis. Nigri ſunt ejuſdem deſcriptionis.

Utrumque opii ſuccedaneum eſt, conſtans oleo craſſo cum mucagine ♌-uum motum compeſcente, acria quævis mulcente, ſanguinis fermentationem compeſcit, hinc tanquam anodynum, hypnoticum & lenitivum in catarrhis à lympha acri, in tuſſi, raucedine, hæmoptiſi, cephalalgia convenit.

Datur ſubſtantia, emulſio, ſyrupus & ☍ expreſſum tam interne quam externe.

℞. Sem. papav. alb. ℥ij. ▽ pap. rhœad. q. ſ. f. Emulſ. cui adde ſyr. ſcabioſ. q. ſ. ad tuſſim & lympham acrem.

℞. Sem. papav. alb. ℥ij decoct. hord. cum liquirit. fact. q.ſ, f. Emulſ.

℞. ☍ papav. expr. ℥j. ☍ N. M. expr. ℥ß. opij g. v. f. liniment. temporibus applicandum.

℞. ☍ papav. expr. ℥j. ☍ nucl. perſic. ℥iß. ☍ hyoſc. aneth. ana ℈j ceræ q. ſ. vel potius ☍ N. M. expreſſ. q. ſ. f. unguentum anodynum & hypnoticum.

***

SEMINA TERREIS CONSTANTIA.

GALLÆ.

Gallæ & glandes cum cortice & foliis quercus conveniunt. Cæterum in glandibus eſt vis alimentoſa, unde tempore annonæ caritatis vices alimentorum ſupplent, in gallis vero major vis adſtrictoria cum auſteritate.

℞. Rad. torment. biſtort. ana ℥j. ſem. papav. alb. ℥ij coq. in ▽ colatur. add. gall. ℥ij f. gargariſm. ad gingivas cruentas.

℞. Gallar. malicor. ana ℥j fol. ſalv. laur. chamom. ſummit. aneth. flor. balauſt. ana p. ij. coq. in vino & ▽ chalybeata, applica pro fotu ad procidentiam ani ex ſphinctere reſoluto.

SEMEN SUMACH.

Rhus folio Ulmi C.B.P. ſunt baccæ ſeu grana orbiculata, villoſa, parva, rubentia ſeu coccinea, includentia nucleum ſeu ſemen orbiculatum ſpadiceum, ſaporis adſtringentis & terreſtris.

Terreis & acido-auſteris conſtant, hinc conveniunt in fluxu alvi, uteri, ſeminis, ſanguiniſque ob vim ſumme adſtringentem. Proſunt etiam in procidentia ani, dum firmant ſphincterem, conveniunt etiam in dentium vacillatione, laudantur ſingulariter ad præcavendos oculos à variolis & morbillis.

Datur ſubſtantia, decoctum.

℞. Sumach. ℥j cape ex ▽ plantag. ad fluxum alvi.

℞. Cort. granator. ℥iiij ſumach. ℥ij Maſtich. ʒj coq. in vino rubro ad procidentiam ani.

℞. Sumach. ℥j infund. c. ▽ fœnicul. ad præſervandos oculos à variolis & morbillis.

MESPILUM.

Immaturum ob auſterum terreum potenter adſtringit in quovis alvi fluxu, vomitu, diarrhæa, dyſenteria. Matura facile fermenteſcunt, & ventriculo debili & inteſtinis nocent.

℞. Meſpill. immatur. crud. utere frequenter ad fluxus alvi, vel coque cum ▽ contra fluxus alvi.

NUX CUPRESSI.

Terreis adſtringentibus cum pauca reſina gaudet, & officium adſtringendi cum blanda vi enodyna abſolvit.

℞. Nuc. cupreſſ. n. vij fol. ebul. Miiij, Cort. aurant. Mac. Cinam. rad. Zedoar. galang. deſtilla ex cereviſia Brunſvvicenſi vel alia generoſa, cape ad crebrum vomitum ex crapula.

BACCÆ MYRTI.

In ſpecie ſtomachi fibras roborant, hinc quibuſdam ſtomachicæ audiunt, laudantur in gingivarum laxitate, præcipue ſcorbutica, junctis ſpecificis.

℞. Bacc. Myrt. ℥ij nuc. cupreſſ. ℥iß. coq. in ▽ f. apoz. add. ſyr. menth. q. ſ. ad ventriculum debilitatum.

SEMEN CYDONIORUM.

Auſteras & terreas particulas obtinent in mucagine in quam reſolvuntur ſemina, hinc in fluxu alvi, ſiſtendo vomitu, in temperando æſtu, ardore linguæ, faucium, cæteris mucaginoſis præſtantiora ſunt, conveniunt etiam partibus ambuſtis & inflammatis oculis.

Datur eorum ſubſtantia, emulſio & mucago.

℞. Sem. cydon. ʒiß. Cort. granat. ʒj add. conſerv. roſar. r. ℥ß. f. bolus in alvi fluxu & in ſiſtendo vomitu add. L. O. gr. ij cum ſyr. cydon. ſ. q.

℞. Sem. Cydonior. ℥ß. ▽ ſemperviv. plantag. nymph. q. ſ. f. mucago add. ⦶tri ʒj phlegm. ⊖li par. & ſyr. papav. q. ſ. detine in ore ad ariduram ſiccitatem & æſtum.

℞. Sem. cydon. fœnicul. hormin. f. mucilag. c. ▽ flor. cyan. add. ſacch. ♄ni ℈ß. impone oculis inflammatis.

℞. Pulp. pomor. redolent ℥iiij mucag. ſem. cydon. ſperm. ranar ana ℥iß. camph. ʒß. ſacch. ♄n. ʒiß. f. cataplaſma ad ambuſta & oculorum contuſiones.

℞. Mucag. cydonior. fact. c. decocto linariæ, applica hæmorrhoidibus dolentibus.

℞. Mucag. ſem. cydon. cape ad faucium præcluſiones ex ▽ forti factas.

CAPUT

CAPUT SEXTUM.
DE HERBIS, FOLIIS ET GERMINIBUS.

PLANTÆ CONSTANTES SALE VOLATILI OLEOSO ſeu AROMATICO.

ORIGANUM.

CReticum & Hiſpanicum vulgari longe præferendum, inter plantas 🜔e aromatico 🝆ſo volatili conſtantes penetrantiſſimum & ſpirituoſiſſimum, propterea in obſtructione contumaciſſima, in morbis ſoporoſis, catarrhoſis, hyſtericis ſpecificum eſt.

Datur infuſ. 🜄 decoct. 🝆 deſt. ingreditur clyſteres.

℞. Summit. origan. fol. thym. ſerpill. ana p. ij flor. origan. Mß. infund. vino ad pint. ij f. infuſum anticatarrhale vel apoplecticum, addita radice pyrethri ad ℥j f. infuſum antiparalyticum tam internum quam externum.

℞. 🜄 origan. deſt. c. vino, inſtilla auribus ad auditum amiſſum, ex cruditate tympanum obſidente.

℞. 🝆 d. origan. gt. iij excipe goſſypio & inde cavo dentis.

MAJORANA.

Mitior eſt origano, ſpecifica in catarrhis, obſtructionibus, doloribus capitis, vertigine & morbis ſoporoſis.

Datur infuſ. ▽ ☍ d. ingreditur quoque clyſteres.

℞. Majoran. ſerpill. thym. ana Mj infund. vino q. ſ. f. infuſum anticatarrhale.

SERPILLUM.

Coincidit cum majorana & datur ut majorana.

℞. Lign. ſaſſafr. ℥iij rad. helen. ℥ß. Zedoar. ʒiij Majoran. fol. ſerpill. ana Mj cort. Winteran. ʒj affund. vin. generoſ. Miij f. infuſum anticatarrhale.

℞. Majoran. Thym. ſerpill. ſaturej. ana q.v. affunde ♁um proprium harum plantarum, f. tinct. ſ. eſſent. contra catarrhos.

℞. ▽ ſerpill. ℥ij cape pro cienda urina, it. in menſibus obſtructis & pro Catarrhis.

℞. ▽ ſerpill. infund. rad. pyretr. f. infuſ. ad aphoniam.

℞. ☍ Nuciſt. expr. ℈j. ☍ ſerpill. gt. vj vel viij f. balſam. ad vertiginem.

THYMUS.

Gratior eſt ſerpillo, & tanquam aroma cibis crudioribus adjungitur.

℞. Thym. infunde ▽ propriæ deſt. & ſpirituoſæ, fac eſſentiam & detine in ore ad læſam deglutitionem, it. ad odontalgiam ex catarrhis.

SATUREJA.

Thymo reſpondet, & fabis additur, quarum flatulentiam præcavet, cæterum ♀ ſtimulat. Datur ut Thymus.

℞. Saturei q. v. affunde aquam ♁uoſam propriam f. eſſentia contra odontalgiam & catarrhos.

℞. Eſſent.

℞. Eſſent. ſaturej. ▽ Magnanim. ſ. ⌒us formicar. ana q. v. exhibe cum tinct. ambr. & elect. diaſatyr. ad excitandam ♀.

MELISSA.

Ob ⊖ia ſua aromatica valde ſpirituoſa, analogiam quandam cum ſpiritibus noſtris habentia ſummum cardiacum & analepticum eſt. Hinc in ſubitanea virium proſtratione & amiſſa memoria, in palpitatione cordis ſpecificum eſt. Commendatur & ad epilepſiam, maniam & ſterilitatem.

Febure primum ens Meliſſæ, mire prædicat, quo aniculam juventuti reſtituiſſe affirmat. [ſi credere fas eſt vid. ejus Chym. part.2. p.m. 221. ſeqq. ubi plura de renovatione & reſtauratione habet.]

Vis analeptica intenditur flavedine Corticum citri.

℞. Herb. Meliſſ. Mj. ▽ Meliſſ. c. v. ℔j. flaved. Cort. Citr. ℥ß f. infuſum ad palpitationem cordis.

℞. Fol. Meliſſ. Mj. incidantur minutim & infundantur in ⌒ vini ℥ij, deind. add. marg. præpar. ʒß. doſ. cochl. ij ex mente Riverii [l. 1. prax. Med. c. 13. p.m. 187. qui addit hoc in Illuſtr. quadam familia Monſpelienſi pro ſecreto gentilitio & hæreditario habitum.]

℞. ▽ Meliſſ. rub. id. vel fœnicul. ana ℥ij ⌒ Theriac. ℥ß. ſyr. flor. tunic. ʒvj Eſſ. Citr. ʒß. M. f. mixtura ad tremorem & palpitationem cordis.

℞. Aq. Meliſſ. ſpirituoſ. ℥j. Cinam. ʒiij borrac. ℈j f. mixtura ad menſium remoram.

℞. ▽ Meliſſ. rub. id. ana ℥ij Cinam. ℥ß. Conf. alk. ʒiij ſyr. rub. id. ſ. cinam. ℥ß. Julep. roſar. ℥ß. M. f. mixtura analeptica.

HORMINUM

HORMINUM SCLAREA DICTUM.

In colica & hyſterica paſſione ut & in fluxu albo mulierum & ♀e ſtimulanda ſingulariter prodeſt, aſt cave ne nimis utaris, quia ob nimiam ſui fragrantiam capiti inducit gravedinem. Datur ut Meliſſa.

℞. Hormin. roriſmar. ana Mij Vin. generoſ. Menſ. ij f. decoct. leni igne vel infuſum in cineribus contra fluorem album & hyſtericam paſſionem.

℞. Horminum, ova conquaſſata & flores lactis q. ſ. f. c. butyro placentæ pro ſtimulanda ♀e, addi poteſt aliquid eſſentiæ ambræ.

*
* *

PLANTÆ SALE AROMATICO VOLATILI CUM QUIBUSDAM FIXIS CONSTANTES.

ROSMARINUS.

Præter vires analepticas vim etiam habet in uteri. capitis & nervorum obſtructionibus, ut & in colica, fluore albo, memoriæ debilitate, ingreditur quoque cum majorana, hormino ſatureja & mentha inſeſſus & balnea contra ſterilitatem.

℞. Roriſmarin. matricar. ſabin. ana Mj bacc. junip. ℥iß. Cinam. ʒiij Croc. ʒj. ⊖ ☐ri ʒß infunde vini q. ſ. ad remoram menſium. Si alvus ſit adſtricta add. helleb. n. ʒj fol. ſenn. ℥j exhibe per vices.

℞. Fol. roriſmar. ℥ß. rad. caryophyllat. ℥j. lign. ſaſſafr. ℥ij affund. vin. generoſ. q. ſ. coque leniter in vaſe clauſo, cape hauſtum ſubinde inter cibandum ad catarrhos.

℞. Ro-

℞. Roriſmar. ſalv. abſynth. ana q. v. coque in vino chalybeato & applica utero contra procidentiam.

SALVIA.

Virtute rorimarino reſpondet, ſed in epilepſia, vertigine, paralyſi, virginum cachexia & leucophlegmatia conſenſum cum Utero habente, ut & podagræ & arthriditi medetur.

℞. Fol. ſalv. ſerpill. ana Mij roriſmar. Miij vel iiij bacc. Junip. ℔ß. coq. in aq. f. pediluvium contra podagram.

℞. Fol. ſalv. q. v. infund. 🜈 vini fac eſſentiam, quam cape cochleatim ad podagram.

℞. Conſerv. ſalv. ℥vj. Zinz. chin. cond. ℥ß. ſem. ſinap. l. eruc. ʒij cardam. cubeb. galang. ana ℈j. ⊖ vol. ſucc. ʒß. 🜈 Ceraſ. nigr. & ſyr. flor. tunic. q. ſ. f. Elect. ad vertiginem ex cruditate ventriculi.

℞. Salv. menth. abſynth. ana q. v. deſtilla c. vino & deſtillatum cape ad famem caninam.

MENTHA CRISPA.

Hujus ⊖ aromaticum ventriculum ejuſque fermentum auget & confortat, ut ſpecifica audiat in cholera, vomitu, colica, anorexia, ſingultu & vertigine, ob ſui penetrantiam etiam uterina eſt. Externe commendatur ad lactis concreſcentiam, it. ad vomitum & dolorem ventriculi. Datur decoct. conſerv. infuſ. aq. deſt. 🜈us, °oo deſt. Elix.

℞. Menth. q. v. affund. 🜈 vin. f. Elix. ſtomachic.

℞. Menth. ſummith. abſynth. ana Mj acori rec. ʒij Cinam. ac. ℥ß. f. nodulus in vino ad vitia Ventriculi.

℞. Aq, Menth. ℥iij Eſſent. Caſtor. ʒiij Laud. op. gr. iij. Elix. Citr. q. ſ. M. exhibe in colica convulſiva.

℞. Aq. Menth. ℥iß. ⁀ theriac. ʒiij Conf. alk. ʒj L. O. gr. ij ſyr. menth. ʒvj f. mixtura ad ſingultum.

℞. ℅ Menth. deſt. gt. ij L O. gr. ij. aq. Menth. ℥j Cinam. ʒij M. contra vomitum præcipue ex hypercatharſi.

℞. ℅ Menth. tacamahac. q. v. cum ℅ Nuciſt. expr. f. liniment. ad vomitum & cardialgiam add. L. O.

MENTHA AQUATICA.

Cum Mentha omnino conſpirat, ſed recipitur hodie inter balnea uterina & nervina.

MENTHASTRUM.

Nil differt à Mentha aquatica, niſi quod odor ſit paulo gravior, ingreditur etiam balnea uterina & nervina.

NEPETA.

In Uteri affectibus, in menſium & lochiorum remora tam interne quam externe ſpecifice uſurpatur, ingreditur quoque balnea & clyſteres uterinos.

℞. Nepeth. puleg. ſabin. matricar. ana q. v. f. infuſum uterinum ex vino ad menſium remoram.

CALAMINTHA.

Cum mentha conſpirat, ſed in flatibus diſſipandis & menſibus ciendis potens eſt, adeo ut & fœtum excludere dicatur, in artuum doloribus & orthopnœa appropriatum medicamentnm eſt.

Externe

Externe pubi ad inflammationem uteri, ut & capiti ad vertiginem, & ventriculo ad vomitum applicatur. Præterea ingreditur catapl. fomenta inſeſſus & balnea.

PULEGIUM.

Menthæ vel calaminthæ analogum eſt, aſt præſtantius, & in tuſſi clamoſa, in menſibus & lochiis pellendis nulli ſecundum. Externe in iſchuria renali ſingulare eſt.

℞. Puleg. Mij infund. vino dulci, add. ſyr. puleg. vel ſcabioſ. contra tuſſim clamoſam, vel loco vini affundatur aqua deſt. ♒uoſa.

℞. Decoct. puleg. ℥iiij Tinct. Caſtor. ʒj ſyr. puleg. q. ſ. ad gratiam, M. exhibe contra iſchuriam. it. injice in veſicam ad eundem affectum.

℞. Puleg. flor. chamom. ſabin. ana Mj pulp. colocynth. ʒß. f. decoct. injiciendum ad provocandos menſes.

℞. Decoct. Puleg. vel aq. ♒uoſ. ℥iiij ex quo cape borrac. Venet. ℈j pulv. Cinam. ℈ß. ☍ d. cinam. gt. j M. f. pulv. ad menſium remoram, ſi dolores adſunt add. L. O. gr. j.

℞. Aq. puleg. ℥ij hyſteric. ʒij Eſſent. Caſtor. ʒß. liq. C. C. ſuccinat. ʒj M. mixtura in paroxyſmo apoplexiæ.

℞. Aq. puleg. ℥iij Cinam. ʒiij M. pro vehiculo in quo medicamenta contra obſtructiones uteri exhibentur.

℞. Gumm. anime ʒß. aq. Puleg. ♒uoſ. q. ſ. redige in formam Emulſ. add. ſyr. puleg. contra tuſſim & aſthma.

EUPATHORIUM.

Salia aromatica obtinet cæteris rigidiora, propterea convenit in obſtructionibus & morbis chronicis ortis à craſſa, terreſtri acida viſcoſa materia. Datur decoctum & infuſum.

℞. Eupator. Mij agrimon. ſummit. abſynth. ana p. ij f. infuſum ex vino ad affectus melancholicos.

℞. Eupator. Mij coq. c. vino, adde camphoram & applica tumoribus hydropicis.

HYSSOPUS.

Ob vim aromaticam temperatiorem in ſpecie pectori dicatur, tuſſi & pleuritidi medetur. Datur decoctum infuſum. ſyr. ſimpl. & comp. aq. deſt.

℞. Hyſſop. Mij coq. in vino, add. colatur. ſyr. propr. contra tuſſim.

℞. Aq. Hyſſop. Menth. ana ℥ij aſthmatic. ℥vj. gumm. ammon. ʒiß. ſyr. Nicot. ℥iij M. cape cochleatim ad aſthma.

℞. Aq. Hyſſop. applica oculis ad ſuggillata, add. aq. puleg.

POLIUM.

Celebre eſt in Uteri vitiis ab obſtructionibus. Sylvius adſcribit vim antepilepticam [Prax. Med. l. 2. c. 20. §. 128. p.m. 427. ubi inter celebratiſſima medicamenta antepileptica poſt rutam, polio montano ſecundum locum tribuit] veteres vero vim contra maniam & noctambulationem.

℞. Summit. polij q. v. affund. vin. f. infuſ. uterinum.

℞. Polij q. v. f. cucupha imponenda capiti.

DICTA-

DICTAMNUS CRETICUS.

Polio accedit & cum pulegio ad partum promovendum commendatur.

℞. Dictamn. Cretic. ʒß. borrac. ℈j f. pulv. dandus ex vino rataceo ad partum promovendum.

℞. Dictamn. Cretic. puleg. polij, ſabin. f. ex vino infuſ. ad part. promovendum.

℞. Fol. dictamn. p. j. infund. ▽ fervent infuſum, ſorbillando, cape ad menſium remoram.

MATRICARIA SEU PARTHENIUM.

Cæteris rigidiore & penetrante ⊖e præfertur, & propterea ratione virtutis uterinæ polium & dictamnum ſuperat, habet quoque blandam vim anodynam, quæ ex odore ad chamomillam nonnihil accedente cognoſcitur. Succus in pluſcula doſi datus alvum laxat.

℞. Matricar. artemiſ. flor. chamom. ſabin. ana Mj f. decoct. ex ▽ & vino ad Uteri affectus. Ex eodem decocto f. etiam clyſter carminans. Etiam cataplaſmatis forma abdomini applicari poteſt poſt partum.

ARTEMISIA.

Virtutibus uterinis remiſſior eſt, quam matricaria. Carbo ad radicem repertus neque productum radicis eſt, neque antepilepticus, ex obſervatione propria.

℞. Artemiſ. Mij ſem. cicer. r. ℥j coq. cum ▽ propria deſtillata, exhibe contra dolores poſt partum, adde tantillum cinamomi ſi adſit virium proſtratio.

℞. ⊖ artemiſ ʒß exhibe ex decocto ficuum ad ſudores movendos, præcipue in variolis & morbillis.

* * *

PLANTÆ SALE AROMATICO VOLATILI OLEOSO CUM MULTO FIXO CONSTANTES.

SCHOENANTHUS ſeu JUNCUS ODORATUS.

[Juncus odoratus aromaticus C.B.P. ſeu Gramen dactylon aromaticum, multiplici pannicula, ſpicis brevibus tomento candicantibus, ex eodem pediculo binis. Eſt ſaporis acris amaricantis & ſuavis, odoris fragrantis.] Dicitur etiam fœnum ſeu gramen camelorum.

Gratius eſt acoro & cum hoc felici ſucceſſu præſcribitur, additur quoque tanquam confortans infuſis & decoctis diureticis contra hydropem v. c.

℞. Rad. gramin. ℥ij vincetoxic. ℥j ſchœnanth. Mij. f. decoctum ex ▽ cui adhuc ferventi pro ſtimulo ſi opus ſit add. rhabarb. ʒiij colatur. edulcor. ſyr. 5. rad. aperient.

℞. Schœnanth. Mj. ſantal. citr. ℥ß. rad. acori ℥j f. infuſum diureticum & ſtomachicum.

ABROTANUM MAS.

Hujus acredo aromatica graveolens pronunciat vim uterinam, fragrantia vero camphorata ſudoriferam in malignis, ob amaritiem vermes quoque expellit. Externe quoque laudatur ad alopeciam; junctum cum

cum floribus chamomillæ & ſeminibus calidis vim acquirit anodynam & uterinam.

℞. Abrotan. Matricar. ſabin. ana Mj rad. galang. ʒß. flor. chamom. p. ij f. dec. Uterin.

℞. ⊖ abrot. ℈ß. Camph. gr. v. C.C. phil. præpar. ℈j f. pulv. exhibendus ex decocto abrotani in febribus malignis.

RUTA.

Vim aromaticam penetrantem & ∽uoſam obtinet, cui quoque ⊖is fixa ſunt aſſociata, unde in morbis chronicis contumaciſſimis & malignis indubitatum promittit effectum. Præ cæteris laudatur in peſte, variolis, morbillis, epilepſia, paſſione hyſterica & affectibus ſoporoſis. Confert quoque in gonorrhœa, ſatyriaſi & priapiſmo. Externe gangrænæ & ſphacelo ſuccurrit Vis concentrata conſiſtit in capſulis ſeminalibus.

℞. Rut. ſabin. abrotan. ana Mj coq. c. cereviſia & cape cum tinctura caſtorei vel croci, contra omnes obſtructiones uteri.

℞. Capſul. ſeminal. rutæ q. v. affunde aquam propriam ∽uoſam f. eſſentia, exhibe in morbis malignis cum camphora, vel ∽u vini camphorato & theriaca.

℞. Rut. affund. acet. q. v. cape tempore peſtis pro præſervatione, add. rad. Zedoariæ, Camphor.

SABINA.

Salia volatilia oleoſa fixis juncta incluſa habet reſinæ, hinc vermes, urinam, præcipue vero menſes & & fœtum promovet, & ob reſinam inter vulneraria ponitur.

In

In verme umbilicali dicitur ſpecifica. [Si pulvis ſabinæ cum vitro veneto & melle ſubigitur & in putamine nucis dimidiatæ infantis umbilico per noctem applicatur. De hoc verme umbilicali vid. Sennert. Pract. l.3. part.x. c.4. p.m. 984. & Hœfer. Hercul. Medico l.7. p.m. 352.]

℞. ☍ deſt. ſabin. gt.ij extract. inſpiſſ. croc. myrrh. caſtor. ana gr. vij f. pil. ad menſes provocand.

℞. Rut. matricar. ſabin. ana Mß. Myrrh. ʒij Croc. ʒß. f. peſſus ſyndoni involvendus &c.

MARRUBIUM ſeu PRASSIUM.

Præter cæteras virtutes quas cum abrotano aliiſque graveolentibus communes habet, poſſidet quid ſingulare in tuſſi aſthmate & ictero flavo ex mucagine orto.

℞. Cichor. tarax. ana ℥ij chelid. maj. ℥j Endiv. Mij Marrub. Mß. ☐ri. ℥ß. coq. ex ▽ & vino, infunde fol. ſenn. ℥vj. contra icterum.

TANACETUM.

Convenit cum cæteris nempe ſabina, matricaria abrotano. Hodie in uterina balnea, cum nepetha pulegio, menthaſtro ingreditur, decocta minus.

ABSYNTHIUM.

Ob ſal volatile oleoſum cum multis fixis ſociatum, convenit in morbis chronicis tanquam polychreſtum, in ſpecie eſt ſtomachicum, antihydropicum, antifebrile, anticolicum, antihyſtericum anthelminticum, ob nimiam ſui graveolentiam accerſere dicitur ſomnum.

Datur

Datur infuſ. ♒. Eſſent. ⊖. ſyr. conſerv.

℞. Summit. abſ. p. ij menth. roriſmar. ana Mß. galang. Zedoar. Zinz. ana ℥j ſem. fœnic. ℥ß. cinam. ʒv Cort. aurant. ℥ß f. infuſ. cum vino ad cachexias quaſcunque.

℞. ⊖ abſynth. ℈j Oc. ♋. gr. xv. L. O. g. j. ☍ junip. gt. ij. M. f. p. contra febres tertianas.

℞. ⊖ abſynth. ℈j ſucci limon. cochl. j. M. exhibe in crebro vomitu. [Inſigniter laudat River. Prax. l. 9. c. 7. p. m. 284 ubi nominat remedium præſtantiſſimum, præſertim in vomitu, qui à febribus malignis ſolet contingere: & exemplis id confirmat Cent. 1. obſ. 15. & Cent. 2. obſ. 99]

℞. Elix. propr. ʒß. Eſſent. abſynth. ʒj f. mixtura in appetitu proſtrato.

℞. Vin. abſynthit. cape per totum tempus curationis febris tertianæ, it. ad anorexiam.

ABSYNTHIUM PONTICUM.

Virtute eſt remiſſius abſynthio vulgari, ſed vim habet diureticam ſpecificam ob quam hydropicis valde convenit.

NB ℞. Abſynth. pont. Mj Card. B. Mj Cent. min. Mß. lign. ſaſſafras ℥ij f. infuſ. in vino contra hydropem, ad diureſin movendam.

***

PLANTÆ SALE AROMATICO CUM PAUCA RESINA CONSTANTES.

CHAMÆPITYS.

Ob reſinoſum ſeu piceum illud, quod tam ex ſapore quam odore percipitur anodyna & nervina eſt

& in doloribus arthriticis & iſchiaticis ſpecifica eſt, in aſthmate convulſivo ſingulariter convenit, ab aliis in mictu cruento & catarrhis laudatur.

℞. Chamædr. chamæpit. ana Mj. Cent. min. Mß. coq. ex aqua ſeu vino dulci contra arthritidem, podagram & iſchuriam, ſi vinum ſuſpectum videatur decoq. in ſero lactis.

BOTRYS.

Virtute incidente diſcutiente & carminante potentior eſt quam chamæpithys, unde in flatulentia hypochondriaca, præcipue infantum expertiſſimum eſt remedium: laudatur quoque in tuſſi, aſthmate & vitiis pulmonum.

℞. Botrys Mij coq. in vin. add. Mell. q. ſ. ad orthopnœam ſingulis diebus capiendum.

Aqua botrys detur cochleatim ad inflationem hypochondriacam infantum.

LAURI FOLIA.

Reſpondent virtute baccis lauri, licet baccæ ob olei copiam vim potentiorem anodynam habeant, quas vide p.138.

℞. Fol. laur. Mj. cort. lign. ſaſſafr. ℥ij f. infuſum diureticum, hydropicis utile.

*
* *

PLANTÆ SALE BLANDO CUM PAUCO VOLATILI CONSTANTES.

AGRIMONIA.

Ob blanda ſua ſalia tam fixa quam volatilia egregium eſt deobſtruens in morbis chronicis, incidens ſangui-

ſanguinis impuritates acidas & terreſtres, eaſque per urinam expellens, unde quoque in mictu cruento ſpecifice laudatur. Externe convenit in uteri procidentia ut & tumore ſcroti. Datur decoct. infuſ. extr. eſſent.

℞. Agrimon. & hyperic. coque ex vino ſeu cereviſia contra mictum cruentum, à ſanguine juſto acidiore.

℞. Agrimon. fol. plantag ana Mj rad. liquir. ℥ß. coq. in aq. ferrarior. ℥xij add. mell. roſac. q. ſ. extr. aloës ℥ß f. injectio in penem, contra mictum cruentum.

℞. Agrimon. fumar. capill. ♀is ana Mß. rad. Cichor. fœnic. ana ℥j ir. flor. ℥ß. fol. ſenn. ʒvj rhabarb. ʒij Cort aurant. ʒiij. ⊖ ♀ri. ʒj M. f. infuſum contra hydropem.

THEA.

Hujus vis in ⊖ volatili blando potius quam fixo conſiſtit, quod acida quævis diſſolvit & per urinam educit, unde in cruditatibus mali hypochondriaci, podagra, arthritide & nephritide mire conducit, ♎us valde auget & confortat, eoſque in continua agitatione conſervat, proinde etiam cephalicum audit, & in confortanda memoria, arcendo ſomno & inducendis vigiliis elegans remedium eſt. A Chinenſibus laudatur in comate tam ſomnolento quam vigili.

BETONICA.

Ob ſubtilem ſuam acrimoniam in qua paucum alcali latet, diureticum & vulnerarium eſt, præcipue in vulneribus & affectibus capitis. Poterius decoctum herniariæ cum betonica in calculo renum & veſicæ commendat, alii laudant decoctum betonicæ ad ſiſten-

dum fluxum lochiorum poſt partum. Chirurgi addunt cataplaſmatibus ſuis cephalicis.

℞. Fol. beton. Mj. ſummit. abſynth. Mß. cubeb. ʒj M. & cum vino fiat catapl. cephalicum. Eligenda ſunt folia recentia.

CHAMÆDRYS.

Convenit cum chamæpithi aſt ſiccior eſt & validior in acidis infringendis, ideo à nonnullis in quartana & menſium remora laudatur.

℞. Chamædr. ʒj f. pul. ad quartan.

℞. Chamædr. Chamæp. ana Mj rad. ariſtol. eryng. ana ℥j M. f. decoct. in remora menſium.

RIBESIUM.

Urinam potenter movet, propterea in ſtranguria & variis morbis chronicis commendatur. Datur infuſum ex vino & cereviſia it. decoctum: plurima vis conſiſtit in novellis foliorum germinibus.

⁂

PLANTÆ SALE BLANDO FIXO CUM QUIBUSDAM TERREIS CONSTANTES.

VERONICA.

Ob ⊖ia temperata diureticum eſt & vulnerarium, chamædri reſpondet: cæterum incidit viſcidam pituitam pulmonibus, inteſtinis & renibus impactam; confert in phthiſi, tuſſi, colica, nephritide.

℞. Heder. terr. veronic. ana Mj, coq. in ▽ ex qua cape myrrh. ℈ß. contra colicam.

HEDERA

HEDERA TERRESTRIS.

Symbolizat cum veronica eam tamen ob ſalia ſua temperata quibus & nonnulla volatilia ſunt admixta, & ratione virtutis antiphthiſicæ ſuperat.

℞. Heder. terr. q. v. coq. c. ▽ ℞ Colatur. ℥viij add. ☍ laurin. ℥j f. Clyſter. ad colicam.

℞. Heder terr. q. v. infund. in ſ. q. ▽æ pro pot. ordin. in calculo & nephritide.

℞. Conſerv. heder. terr. roſar. ana ℥j flor. △is. antihect. Pot. Oc. ♋ ana ʒj f. Elect. contra pthiſin.

℞. Heder. terr. chærefol. veron. ana Mj helen. ir. flor. ana ℥ß. rad. ari. ʒj caricar. n. v. paſſul. min. ℥iß. liquirit. ℥ß. f. decoct. ex ▽ in tuſſi & phthiſi ad incidendam mucaginem pulmonum.

PIMPINELLA.

Cum hedera terreſtri, veronica, agrimonia ſimul præſcribitur & viribus etiam coincidit, ſingulare quid obtinet in dyſenteria, tum quod acidum dyſentericum infringat, tum quod laxatas inteſtinorum fibras blande adſtringat. Sanguinis quoque purgatorium polychreſtum audit.

EUPHRASIM.

Ob ſalem tenuem blandum acidas ſordes abſtergentem Oculis eorumque inflammationibus dicata eſt. Hoffmann. Clav. in Schrœd. Euphraſiam flore albo decoctam in vino contra icterum flavum utilem deprehendit.

℞. ▽ Euphraſ. deſt. cape intro & lava oculos inflammatos & caliginoſos, pro hoc ſcopo cape etiam conſervam.

HYPERICUM.

Hujus virtutes vid. in capite de ſemine hyperici.

℞. Virg. aur. Mj ſummit. hyper. p. ij. rad. ſymph. ℥ß. raſur. corn. cerv. ℥ß. coq. ex aq. ferrarior. ℞. Colatur. ℥xvj add. ſyr. dialth. ℥j aq. Cinam. ℥ß. pro vulnerario. it. in eroſis à dyſenteria inteſtinis.

℞. Summit. hyperic. herb. anagall. fl. purp. ana q. v. f. decoctum contra melancholiam, delirium, maniam, pro potu ordinario.

*
* *

PLANTÆ SALE AROMATICO OLEOSO PENETRANTISSIMO CUM ACIDO VOLATILI CONSTANTES.

COCHLEARIA.

Reſpondet aromaciæ & cortici Winterano, tamen multo penetrantior eſt & validior ut ejus acredo ubi ſuſpecta videtur temperanda ſit paſſulis, dulcibus aliis, & temperatis, ſcilicet lacte caprillo, ſero lactis. [Optime cochlearia cum acetoſa conjungitur & ſic evadit nobile remedium antiſcorbuticum. vid. Act. Hafnienſ. ann. 1671. obſ. 9. p.m. 35.]

Datur ſubſtantia, ſuccus. conſerv. infuſ. aq. 🜅us 🜔. Elix.

℞. Rad. helen. armorac. ana ℥j cochlear. naſturt. ana Mj paſſul. min. ʒvj f. infuſ. ex vino vel ſero lactis contra ſcorbutum.

℞. Cochlear. Naſturt. armorac. rad. ari, contunde omnia, affund. 🜅um ſambuci, digere & deſtilla ex balneo f. 🜅us antiſcorbuticus.

℞. Coch-

℞. Cochlear Mj fumar. Mj cortic. fraxin. ℥vj. baccar. Junip. ℥j ſummit. abſ. Mj rad. cichor. ℥j helen. ℥ß. ſtrobil. abiet. n. v, ſine hæc omnia efferveſcere cum cereviſia, & fiat cereviſia medicata contra ſcorbutum.

℞. ⏜ cochlear. impone membris ab epilepſia contortis

NASTURTIUM.

Virtute antiſcorbutica remiſſius eſt Cochlearia, cœterum potenter pellit urinam & in verme pericardii enecando propriam dicitur habere vim. Datur ut cochlearia.

℞. Naſturt- ▽tic. exprime ſuccum, ſine efferveſcere cum fermento, miſce cum novo naſturtio & fac maſſam, deſtilla per balneum, cujus cochl. j. vel ij exhibe in ſcorbuto.

⁂

PLANTÆ SALE VOLATILI ACRI CONSTANTES ABSQUE OLEO VOLATILI.

ERYSIMUM.

Convenit in tuſſi pro incidenda mucagine, it. in raucedine ex lympha acida viſcida in faucibus hærente.

℞. Eryſim. Eruc. ana Mj coq. in hydrom. contra raucedinem.

PERSICARIA.

Ratione ⊖is volatilis penetrantis, aperientis & reſolventis ſumme commendatur in obſtructione hypochondriorum, meſenterii & aliorum viſcerum, & ob nimiam ſuam acrimoniam temperatur paſſulis.

▽ deſt. c. ☿io dulci contra vermes poteſt dari, commendatur etiam contra calculum.

℞. Perſi-

℞. Perſicar. Mj paſſul maj. ℥ij coq. ex ▽ ſ. cereviſ. contra obſtructiones varias.

Vis magnetica quam in curandis ulceribus & vulneribus obtinere dicitur, docente experientia merito explodenda eſt. [Conf. hac de re Helmont. d. Magnet. Vuln. curat. §.29. p.m.459. Fonſeca conſult.37. p.m.252. & Marc. Marci. Phil. Vet. reſtitut. Part. v. Sect. I. ſubſect. vj. p.m. 495. ſeqq. de magnetiſmo Perſicariæ.]

Externe in tumoribus reſolvendis magni uſus eſt, ut & in ulceribus ſordidis mundificandis ſuccus cavo dentis inditus dolorem ejus tollit.

℞. ▽ deſt. Perſicar. cape cochleatim contra calculum, tempera cum ſyr. d. althæa.

CUSCUTA. EPITHYMUM.

Notæ ſunt plantæ ob vim reſerandi obſtructiones præcipue in ſcorbuto & hydrope, quæ in ⊖e volatili acri, mitiori tamen quam in perſicaria & eryſimo conſiſtere videtur, ſpecifice laudatur in viſcoſa bile incidenda. Datur ▽ deſt. ſyr. & infuſ.

℞. Rad. helen. ℥j. raphan. ruſtic. ʒvj ſummit. cent. min. Mj cortic. Winteran. ʒj. Cuſcut. p. ij. paſſul. min. ℥j f. infuſ. add. fol. ſenn. ℥j rad. hell. n. ʒij contra malum hypochondriacum.

*
* *

PLANTÆ SALIBUS FIXIS SEU AMARIS CONSTANTES CUM PAUCO VOLATILI SALE.

CENTAURIUM MINUS.

Ob intenſam acredinem & amaritiem in qua ⊖ fixum latet, potentiſſime deobſtruit, acida deſtruit, unde

unde in obſtructionibus chronicis & febribus intermittentibus primum locum obtinet, aperit quoque ſuppreſſas hæmorrhoides & menſes promovet. Extra ulceribus applicatum ea mundat, additur quoque clyſteribus contra morbos ſoporoſos.

Datur ſubſt. decoct. infuſ. extract. aq. ⊖.

℞. Cent. min. ℥ß. ⊖ propr. ℈ß f. pulv. exhibendus c. aq. propria.

℞. Cent. min. Card. B. ſummit. orig. ana Mß. Croc. or. ℈j fruct. alkek. ℥j coq. in aq. ſimpl. ad conſumt. mediet. ad ℔ij hoc decocto curatus fuit ſenex quidam febre quotidiana cum anorexia & ictero afflictus.

CARDUUS BENEDICTUS.

Virtute antifebrili reſpondet centaurio minori, quo tamen paulo mitior eſt, cæterum ſanguinem coagulatum mire diſſolvit, & propterea in pleuritide confert. Decoctum obſervante Ballonio urinam reddit craſſam & fœtidam.

Datur ſubſt. decoct. infuſ. aq. deſt. ſyr. & ⊖.

FUMARIA.

Salibus nonnihil ſubtilioribus volatilioribus & penetrantioribus gaudet quam Centaurium & carduus benedictus, & propterea ſanguinis acidam dyſcraſiam corrigendo, in morbis chronicis, malo hypochondriaco, ſcorbuto, ſcabie, quartana, ſingularis eſt efficaciæ.

℞. Fumar. rec. Mj. rad. Chin. ℥ß. M. f. decoct. contra ſcabiem, ut & malum Hypochondriacum.

℞. Cort. ſaſſafras ℥iß. fumar. rec. Mj. f. infuſ. ex vino ad obſtructionem hypochondriorum.

℞. Succ. cochlear. fumar. ana q. v. M. c. ſero lactis exhibe contra ſcorbutum, tanquam medicamentum temperatius.

℞. Pil. d. fumar. ʒß. ☿ d. ℈ß. cape ad ſcabiem ſcorbuticam, it. ad ſanguinem purificandum.

SCORDIUM.

Ob ⊖ia penetrantiſſima tam fixa quam volatilia omne peregrinum præcipue ad putredinem inclinans per ſudorem expellit, hunc in finem eſt alexipharmacum, antipeſtilentiale & polychreſtum. In malignis, it. in curando morbo Hungarico, in lumbricis. [Qua occaſione vis putredini reſiſtendi in ſcordio reperta fuerit vid. ap. Galen. de antidotis l. 1. c. 12.]

Dat. ſubſt. decoct. Elect. diaſcordium dictum, ſtimulatur optime camphora, ſed à plebejis allio.

℞. Scord. abſynth. cent. min. ana Mj. Myrrh. ℥j camph. ʒij vel iij affund. ♒ vin. f. infuſum pro lavamento ad curandam gangrænam vel ſphacelum.

℞. Scord. ſummitat. hyperic. ana Mj f. decoct. pro lumbricis.

℞. Scord. ſummit. hyperic. fol. ſenn. ana ℈j f. pulv. contra vermes.

℞. Diaſcord. ʒj camph. gr. iij ⊖ volatil. C. C. ℈ß. M. f. bolus in malignis ad ciendos ſudores.

LUPULUS.

Salem ſuum utrumque tam fixum quam volatilem in quadam quaſi reſina & glutinoſitate incluſum habet, ob quam rationem non tantum craſſa & viſcida incidit & vel per ſudorem vel per urinam expellit, ſed

&

& blande ad ſomnum componit: in malo hypochondriaco, ſcorbuto, febribus chronicis, ſcabie aliiſque cutaneis affectibus laudatur. Turiones etiam vim alvum laxandi obtinent.

℞. Capit. lupul. herb. fumar. ana Mj paſſul. min. ℥ij f. decoct. contra malum hypochondriacum, cui in alvi adſtrictione infunde rad. hell. n. rhabarb. ana ʒij.

*
* *

PLANTÆ PURGANTES.

SENNA.

Blandum, polychreſtum & catholicum, humores quoſvis movens, & per alvum blande educens, jungitur variis ſpecificis, morbo cuique appropriatis, vis potior in ⊖e volatili particulis terreis & nonnihil gummoſis videtur incluſa, quare potius in infuſo, quam in decocto exhibendum eſt. In gummoſitate autem potior vis conſiſtere videtur.

Dantur folia pediculis & ſtipitibus rejectis, corrigitur & ſtimulatur ſale ♀ri ut & rhabarb. in ſubſtantia datur ad ℈ij vel ʒj in infuſo ad ℥ß. vel ex aq. fumariæ ♀iſata vel ſero lactis, vel decocto prunorum, vel melius decocto paſſularum.

℞. Fol. Senn. ℥ß. rhab. ʒß. ⊖ ♀ri. ʒj ſem. aniſ. fœnicul. ana ℈j cinam. ʒß. infunde aq. fumariæ pro infuſo laxante.

℞. Fol. ſenn. ʒiij ⊖ ♀ri. ℈j. infund. aquæ ſimplici ferventi. ℞. Colatur ℥ij vel iij edulcor. ſyr. d. pom q. ſ.

℞. Fol. ſenn. ʒj caryoph. arom. gr. vij. ſacch. cand. ℈j. M. f. p.

℞. Fol. ſenn. ℥ß. ſem. aniſ. ʒi affund. aq. ferventiſſ. ſtent in infuſo, infuſum cape ſorbillando ut thea, pro laxanda alvo.

SOLDANELLA.

Gallica præſtantior quam noſtra Belgica, blande aquas præcipue hypochondriorum expurgat per alvum, hinc antiſcorbuticis, & antihypochondriacis medicamentis pro laxante adjungitur.

Datur pulv. ʒj ſuccus ad ℥ß. in decoct. Mj.

℞. Fol. ſoldanell. ℥j f. decoct. vel infuſ. cui infund. rhabarb. ʒj exhibe hypochondriacis ſimul hydrope correptis.

℞. Fol. ſoldanell. cochlear. ana Mj contere & extrahe ſuccum, quem miſce cum paſſularum decocto; exhibe ſcorbutico ad laxandam alvum.

GRATIOLA.

Aquas hypochondriacorum & hydropicorum fortiter tam per ſuperiora quam inferiora expurgat, in enecandis & expellendis vermibus efficaciſſima eſt.

Externe hydrocephalis medetur.

Corrigitur Zinzib. Mac. Cinam. aniſ. ſpecificatur in malo hypochondriaco cochlearia. Datur pulv. ad ℈ij. infuſ. ad ʒij. ſuccus inſpiſſatus ad ʒj & aliquando ad ʒiß. optimum eſt extractum cum ſp.u vini factum, quod datur à ℈j ad ʒß. conſerva folior ad ʒij, paratur quoque ⊖ fixum, quod hydropicis ſpecificum & diureticum eſt.

℞. Gratiol. ℥ß ſem. aniſ. ʒij paſſul. min. ℥j coq. leniter in vino, propinetur colatura ad educendas aquas hydropicorum.

℞. Sol-

℞. Soldanell. gratiol. fol. ebul. aſari, ſambuc. flor. geniſt. perſicor. cort ſambuc. frangul. rad. ireos. germanicæ, eſul. ana q.v. add. pulp. colocynth. p.j. coq. in urina, applica totum hydrocephalo.

℞. Gratiol. Cent. min. ana Mj ſem. tanacet. ſanton. ana ℥ß. coq. in ▽ f. Clyſter. ad enecandos & expellendos vermes, hærentes præcipue in inteſtinis craſſis.

NICOTIANA.

Hujus vis potenter incidendi, aperiendi & reſolvendi ut & purgandi aquas tam ſupra quam infra conſiſtit in ⊖ etam fixo quam volatili, vis ſomnum inducendi & ſtuporem, dolorem mulcendi, conſiſtit in pinguedine ſeu glutinoſitate volatili.

Foliorum infuſum vel decoctum purgat per ſuperiora & inferiora, idem faciunt folia in pulverem trita, quæ maſticata ſalivam potenter movent, dentium quoque dolorem tollunt, naribus attracta ſternutationem inducunt. Decoctum mundificat ulcera ſordida.

Fumus accenſorum foliorum in quo ⊖ volatile ☍ſum ſeu mucoſitas ſtuporem inducit, ac ſomnum accerſit.

Oleum deſtillatum ſanguini immediate permixtum præſentaneam infert mortem, unde Viganus in Chym. p.36. oleo hoc fœtido feles enecari dicit, dummodo una guttula ſtilletur in linguam. [Circa hanc rem plura experimenta inſtituit Franc. Redi in Experim. circa res diverſas naturales pag. 13. ſeq. it. p. 54. & Joh. Jacob. Harder. in Apiario obſ. 1. uſque ad ſeptimam, qui in varii generis animantibus experimenta fecit.]

Succus ad ʒij datus purgat ἄνω & κάτω, injicitur quoque fiſtulis ad eas mundificandas & curandas.

Balſamus Nicotianæ Vulnerarium eſt externum Unguent. ex Nicot. factum necat pediculos; quod etiam præſtat ☍ Nicot.

Syrup. & ▽ Nicot. in aſthmate & catarrho ſuffocativo valent. Proſtat quoque Syrup. Nicot. emeticus in aſthmate & aliis thoracis affectibus deprædicatus.

℞. Fol. Nicot. ʒiß infund. vino ſeu Cerev. ad ℔ß. cape infuſ. pro purgante ἄνω & κάτω.

℞. Fol. Nicot. ʒj fl. roſ. pall. ʒij infund. ſ. q. vini pro Vomitorio & purgante.

℞. Syr. Nicot. heder. terr. d. ſymph. Fern. ana ℥ß. M. exhibe contra aſthma.

℞. ▽ deſt. Nicot. ℥j. ☍ ♆ per camp. gt. v. cape contra aſthma.

℞. Succ. Nicot. inde vulneribus ad ea mundificanda, add. quoque myrrh. aloë ad vulnera & fiſtulas curandas.

⁂

PLANTÆ CONSTANTES SALE NITROSO LAXANTE.

MERCURIALIS.

Ob nitroſum ſuum laxantem ſalem fœces emollit & promovet, propria quoque virtute fluxum menſium uterumque purgat, ob quam rationem forſitan ſterilitatem à vitio Uteri arcere dicitur.

Adhibetur interne & externe, in decoctis, cataplaſmat. peſſariis, Clyſteribus.

℞. Mer-

℞. Mercurial. parietar. violar. coque in juſculo pulli, vervecis vel alterius ad alvi adſtrictionem.

℞. Myrrh. ℥ß. Croc. or. ʒß. ſucc. Mercurial. vel mell. Mercurial. M. f. peſſus ſindoni involvendus, & utero intrundendus ad provocandum menſes & purgandum uterum, ſtimuli loco adde troch. alhand. ʒß.

PARIETARIA.

Reſpondet mercuriali & in ſpecie ratione nitroſi ſalis copioſioris movet ſimul urinam, ejuſque ardorem temperat, hinc commendatur in hydropicis. Externe applicata pubi, iſchuriæ infantum medetur.

℞. Parietar. Mij chæref. Miß coq. cum butyr. rec. ℥ij add. ℅ ſcorp. ℥iß. applica pubi ad ſuppreſſionem urinæ, ut & ad tormina ex calculo orta.

VIOLARIA.

Virtute emolliente laxante cum parietaria & mercuriali convenit, juvat quoque ad expectorationem ſi cum propriis floribus coquatur & exhibeatur.

℞. Violar. Mj. coq. c. ▽ colaturæ ferventi infunde flor. violar. p. ij edulcora ſyr. violar. pro expectoratione.

*
* *

PLANTÆ CONSTANTES PARTICULIS OLEOSIS AQUOSIS CRASSIS.

ALTHÆA.

Reſpondet ſeminibus & radici, ingreditur decocta clyſteres, cataplaſmata emollientia.

℞. Fol. Malv. alth. coq. ex ▽ add. camphor. & applica ſcirrhis emolliendis & maturandis.

℞. Fol.

℞. Fol. Malv. alth. flor. chamom. ana Mj. melilot. Mß. ficu. n. iij liquir. ℥ij vel iij coq. in lacte f. cataplaſm. emolliens, colatura inſervire poteſt in angina pro gargariſmata.

MALVA.

Convenit cum althæa, aſt debilior eſt.

TUSSILAGO.

Hujus folia tenuiorem habent mucaginem quam althææ folia, hinc ſpecifica ſunt in tabe, phthiſi, ulcere pulmonum vel aliorum viſcerum, ut & in vomica pulmonum, it. in catarrhis à ſalibus acribus. Horum decoctum Lindanus laudat in aſthmate & orthopnœa, ut & in pleuritide à quibuſdam.

Foliorum fumus hauſtus per fiſtulam à Boileo commendatur in phthiſi & catarrhis ſalſis.

Datur pulvis foliorum, ſuccus expreſſus, Syrupus ut & decoct. cum propria radice factum, addita China, ſaſſafras, aliiſque appropriatis. Vulgus recipit folia ad placentas contra tabem & phthiſin.

℞. Fol. tuſſilag. ſuccin. ſem. aniſ. ana q. v. hauri fumum per fiſtulam contra catarrhos ſalſos.

℞. Tuſſilag. flor. 🜍is, ſuccin. fol. nicotian. hauri fumum per fiſtulam contra phthiſin.

VERBASCUM.

Ob mucaginem emollientem terreis junctam convenientiſſimum eſt in teneſmo, dyſenteria, dolore hæmorrhoidum in ſpecie ad procidentiam uteri & ani.

℞. Fol. verbaſc. querc. coq. in lacte contra teneſmum.

℞. Verbaſc. Mij ſem. hyoſc. coq. in ▽ contra teneſmum & dolores hæmorrhoidum.

PLANTÆ

***

PLANTÆ TERREIS CONSTANTES.

EQUISETUM ſeu CAUDA EQUINA.

Virtute fibras Ventriculi & inteſtinorum roborante, adſtringente, ut & incraſſante ſanguinem cæteris ſui generis præpollet, & propterea in ſiſtendis hæmorrhagiis, in ſedando mictu cruento & cæteris generibus fluxuum ſanguinis laudatur; recipitur hoc nomine inter vulneraria.

℞. Equiſet. Millefol. plantag. ana Mj coq. in 🜄 colat. add. ſucc. vel ſyr. granator. q. ſ. ad ſiſtenda profluvia ſanguinis.

℞. Equiſet. q.v. coq. in 🜄 ad ulcera quævis pulmonum ſanguinolenta.

MUSCUS QUERNUS.

Equiſeto & cæteris quercuum partibus longe anteponitur ob inſignem ſui vim deſiccativam, & humorum motum ſiſtentem, ſpecifice laudatur in tuſſi à tenui & acri lympha proveniente, pulmones erodente & nonnunquam phthiſin inducente. Externe quoque cum phlegmate 🜕li, aceto vel alio acido applicatus mire ſiſtit ſanguinem fluentem. Ingreditur quoque capillares pulveres ſeu cyprios ad deſiccandas humiditates.

℞. Muſc. Quern. ℥j. ⚇ aniſ. gt. iij. ſacch. cand. ℈j. M. f. pulv. contra tuſſim.

℞. Muſc. ℥ij lac. 🜍is ℈ij ſem. aniſ. ℈j M. f. pulv. contra tuſſim.

℞. Muſc. quern. uſneæ cran. human. ana ʒß cape c. ▽ burſ. paſtor. in hæmorrhagiis internis etiam periculoſiſſimis.

℞. Equiſet. Millefol. muſc. quern. ana Mj. coq. in ▽ cape contra mictum cruentum.

PULMONARIA.

Muſco querno inferior eſt virtute, cæterum in ulcere pulmonum & ſanguinis ſputo, hectica & phthiſi ſingulariter commendatur, uſurpatur ut muſcus quernus.

℞. Pulmonar. fol. tuſſilag. ana Mj f. decoctum contra phthiſin ex ulcere viſceris nobilioris cum alvo adſtricta.

SOLIDAGO SARACENICA. VIRGA AUREA.

Harum plantarum particulis terreis, ⊖is ⦶troſi diuretici aliquid eſſe incluſum docet earum vis diuretica, & quia impedimenta quæ vulnerum curationes morantur removent, inter vulneraria recenſentur. Datur decoctum & pulvis.

[Quam mira præſtare dicatur virga aurea in reſolutione calculi renalis vid. ap. Wedel. Amœn. Mat. Medic. l. 1. ſ. 3. c. 9. p. m. 185. ex Barclai Satyrico p. 1. p.m. 75.]

℞. Virg. aur. ʒij cape ex vino ſeu decocto proprio pro diuretico & nephritico.

℞. Alchymill. virg. aur. conſol. ſaracen. ſummit. hyperic. ana Mj rad. conſolid. ʒij glycyrrh ℥ß. coq. ex ▽ vel Cerev. col. add. ſyr. d. ſymph. Fern. q.ſ. ad grat. f. apoz. vulnerarium & diureticum.

BURSA

BURSA PASTORIS.

Tam intra quam extra pro adſtringente & vulnerario habetur, convenit quoque in ſputo ſanguinis & hæmorrhagiis quibuſvis, nonnulli prædicant etiam in pollutionibus nocturnis.

℞. Burſ. paſtor. Mij. f. dec. ex ▽ ad ſiſtendum hæmorrhagias, & præcipue nimium menſium fluxum, mictum & vomitum cruentum, vel cape ſucci expreſſi ℥ß ex decocto proprio contra eundem affectum.

VERBENA.

Vim adſtringentem cum cæteris ſuæ ſortis obtinet, ab hac an dependeat vis ſpecifica contra dolorem capitis à tenui & acri lympha ortum aliis judicandum relinquo, interim omnibus cephalicis medicamentis impune addi poteſt. [Etiam ex collo ſuſpenſam miraculi inſtar dolorem capitis ſedaſſe exemplo demonſtrat Foreſt. lib.9. obſ.52. p.m. 270.]

℞. Extr. verben. inſpiſſ. ☍ hyoſc. papav. q. ſ. applica temporibus ad dolores capitis tollendos.

℞. Santal. rubr. ʒij flor. roſar. fol. verben. ana Mj camph. ʒß. ſacch. ♄ni ʒj acet. q.ſ. f. epith. egregie ſiſtens dolorem capitis febricitantium, temporibus applicatum.

℞. Aq. deſt. verben. ad ℥iiij cum aliquot guttulis ♎us ⊕li aſſumta Hoffmanno Clav. in Schrœd. in dolore capitis mira præſtare dicitur.

℞. ▽ verben. cape bon. hauſt. cum ℈j Oc. ♋ ad abortum præcavendum.

FOLIA QUERCUS.

Cum corticibus coveniunt, in ſpecie laudantur in procidentia ani & uteri.

PYROLA.

Virtute vulneraria, quæ ſola de hujus foliis innotuit cæteris omnibus congeneribus præfertur.

℞. Pyrol. ſanicul. virg. aur. ana Mij hyperic. p. ij rad. ſymph. ℥iij coq. in vino f. decoct. vulnerarium.

SANICULA.

Cum cæteris ſuæ ſortis ad vulneraria tam interna quam externa refertur.

℞. Sanicul. alchym. beton. ana Mj rad. ariſtol. ℥j ſymph. ℥iij f. decoct. vulnerarium tam internum quam externum, pro interno add. ſyr. d. ſymph. Fern.

ALCHYMILLA.

Ob vim validam adſtringentem, additur tam decoctis quam clyſteribus, convenit vulneribus medendis, dyſenteriæ curandæ & laxis pudendis conſtringendis, laudatur quoque ad flaccidas laxaſque mammas duriuſculas & minores reddendas.

℞. Alchymill. fol. ſymph. cum toto q. v. f. inſeſſus vel balneum vel fotus ad mammas & pudenda ſtrictiora reddenda. [De vi adſtrictoria radicis ſymphiti in hoc caſu vid. ridiculam hiſtoriam apud Phil Hœchſtetter. Decad. 3. caſ. 5. p. m. 236.]

PILOSELLA.

Inter vulneraria temperatior eſt, & propterea infantibus ad herniam curandam datur. Laudatur quo-

quoque in temperanda lympha acida, à qua aſpera arteria moleſte afficitur & inflammationem faucium producit.

℞. Piloſell. Mij coq. ex Cerev. f. gargariſm. ad dentium dolores ortos ad humore acido acri.

POLYGONUM.

Præter virtutem vulnerariam habet quoque ſpecificam in hæmoptyſi & quavis hæmorrhagia. Externe applicatur carpis manuum ad æſtus febriles tollendos.

℞. Succ. polygon. ℥j vin. auſter. ℥iiij cape contra vomitum ſanguineum.

℞. ▽ Polygon. melius decoct. in quo exhibe rhab. toſt. N. M. toſt. ana ℈ß. contra hæmoptyſin.

***

PLANTÆ PARTICULIS TERREIS TEMPERATIS ET BLANDIS VOLATILIBUS CONSTANTES.

HERBÆ CAPILLARES.

Sunt Adiantum album ſeu ruta muraria. Adiantum nigrum ſeu Capillus ♀is. Polytrichum aureum. Aſplenium ſeu Ceterach ſeu Scolopendrium.

In terrea ſubſtantia habent quidpiam volatilis, ob quod blande incidunt, reſerant hinc viſcerum obſtructionibus quibuſvis, in ſpecie ſcorbuto, malo hypochondriaco, ſcirrhis & obſtructionibus lienis conducunt, proſunt quoque in vitiis pectoris & renum, ad incidendum mucum illis impactum. Ob ſubſtantiam terream tubulos & poros conſtringunt ne defluant capilli.

Pulvis aureus qui parti foliorum averſæ adnaſcitur cum ſuccino albo & ſucco portulacæ datus mire ſiſtit fluxum gonorrhææ.

Datur decoctum, ſyrupus tam ſimpl. quam comp.

PHYLLITIS ſeu LINGUA CERVINA.

Convenit cum plantis capillaribus & cum his ſimul & ſemel præſcribitur, frequenter ingreditur vulneraria decocta.

℞. Phyllit. Mij virg. aur. Mj rad. helen. ℥iij rad. Cichor. ℥vj bacc. juniperi ℔ij ſine efferveſcere cum Cereviſia recenti f. potus diureticus & vulnerarius, cachecticis & hydropicis utiliſſimis.

CAPUT SEPTIMUM.

DE FLORIBUS. SALIBUS VOLATILIBUS OLEOSIS SEU AROMATICIS CONSTANTES.

FLORES RORISMARINI OFFICIN. ANTHOS.

Recentiores ſunt ſubtiliores & ſpirituoſiores quam folia, cum quibus cæteroqui conveniunt. Ex his fiunt ♎. Eſſent. ▽ reg. Hung. ☍ deſt. conſerva cum ſaccharo facta, in qua tanquam appropriato vehiculo cephalica, uterina & nervina exhiberi poſſunt.

℞. Con-

℞. Conſerv. anth. Menth. ana ʒß. ⊖ vol. ſuccin. gr. xv. M. f. uterin. & cephalic. pro ij. doſ. ſi addideris petrol. alb. gt. ij vel iij habebis medicamentum contra vertiginem.

℞. Conſ. anth. ʒß. ☍ cinam. gt. j. borrac. Ven. ℈ß. f. bolus in menſium remora.

℞. ♌us vin. rectif. ſ. melius ♌ roriſmar. per ferment. fact. ſ. ▽ propr. deſt. ſæpius cohob. affundendo floribus roriſmarini & deſtilla ex balneo, hunc laborem continua donec deſtillatum fragrantiſſimum acquiſiverit odorem & ſaporem, hoc repone pro ▽ Reginæ Hungariæ.

℞. ▽ fragr. roriſmar. & ♌uoſ. cohoba cum ☍ deſt. roriſmar. recenti, deſtilla blando △e, donec imbiberit tantum ☍ quantum commode ſuſcipere poteſt, hoc applica partibus paralyticis.

℞. ▽ Reg. Hung. applica vertici capitis ad tollendos dolores.

FLORES LAVENDULÆ
ſeu SPICÆ.

Virtutibus cephalicis uterinis & nervinis roriſmarini floribus multo præſtantiores & penetrantiores eſſe, probat primo ☍ deſtillatum, deinde ſalivatio quam flores & folia maſticata excitant, hinc in ſoporoſis & catarrhalibus morbis valdopere laudantur. Horum odorem fugiunt pediculi, blattæ & alia inſecta, propterea in his fugandis vel enecandis conducunt.

Datur ▽ deſt. dec. conſ. infuſ. ☍ deſt.

℞. ▽ deſt. lav. ♌uoſ. ℥ij Eſſ. Caſtor. ʒij. ♌ Ceraſ. n. ℥j detin. in ore contra aphoniam & amiſſam loquelam.

℞. ☍

℞. 🝆 d. ſpicæ, 🝆 thereb. ana gt. v. 🝆 laur. expr. ʒj camph. gr. v. M. f. balſamus, cujus gr. v. in juſculo calido exhibita mire conducunt convulſis, externe applicatur paralyſi, ſtupori, & omnibus convulſionis generibus citiſſime medetur.

℞. Conſ. flor. tunic. ʒj add. 🝆 ſpic. gt. v. f. bolus contra aphoniam.

℞. Flor. & fol. lavend. ana q. v. affund. ▽ ∽uoſam propriam, add. rad. pyrethri, ari q. v. f. infuſum contra aphoniam.

℞. Decoct. vel lixivium lavendulæ, lava caput pro enecandis pediculis.

FLORES SALVIÆ.

His folia multo anteponuntur.

CHÆMEMELI FLORES.

Licet Romani flores habeantur penetrantiores quam vulgaris, tamen in his potior vis anodyna eſt; cæterum utriuſque flores aliquid balſamici volatilis in ⊖ibus tam fixis quam volatilibus fovent, à quo pendet ſingularis vis acria quævis mulcens & obtundens, hinc in cardialgia, colica ſimplici & convulſiva & nephritica, ut & flatulentis omnibus diſcutiendis nulli cedunt, promovent quoque lochia, menſes, dolores poſt partum tollunt, conducunt etiam in lacte cocti contra dyſuriam. Externe emolliunt, & dolorem præcipue teneſmi leniunt, ingrediunturque clyſteres ad flatus & tormina.

Datur eorum pulv. decoct. ▽ deſt. 🝆 d. eſſent. c. ▽ propria f. ſpirituoſa facta.

℞. Flor. chamom. Mij coq. in vin. rhenan. ℔ij leniter exprime & expreſſum novis floribus affunde, laborem aliquoties reitera, exhibe cochl. iij contra nephritidem.

℞. For. chamom. Mj coq. ex vino leniter, exhibe contra dolores poſt partum, ſtimuli loco add. eſſent. artemiſ. q. ſ. vel coq. in lacte ad dyſuriam & aromatiſa nuce moſchata.

℞. ▽ deſt. chamom ℥iij. ⁓ theriacal ʒij L. O. gr. j. 🝆 d. chamom. gt. ij ſyr. è Cort. aurant. ℥ß. f. mixtura in colica flatulenta.

℞. ▽ Menth. Chamomill. ana ℥ij. ⁓ Carm. ſylv. ℥j Elix. menth. ʒij. ⁓ ⦶tr. dulc. q. ſ. ad acorem L. O. gr. j. vel ij f. mixtura colica.

℞. 🝆 d. chamom. gt. v. 🝆 laur. expr. ℈j f. balſam. quo inungatur regio ventriculi ad cardialgiam, add. L. O. gr. iiij exhibe cum juſculo calido hujus balſami ℈ß interne contra cardialgiam.

℞. Flor. chamom. ʒj cape ex decocto proprio ad febres puerorum, idem confert in lumbricis.

℞. Flor. chamom. Mij coq. in vaſe clauſo, igne leni ſufficienter ℞ Colatur. ℥viij add. 🝆 laur. expr. ℥j f. clyſter. ad colicam quamcunque, ſi nullæ exeant fœces, add. ⊖is ʒij injice denuo clyſterem, in torminibus cum convulſione add. Eſſent. caſtor. fact. cum 🝆 ſubtili therebinth. ℥ß.

CROCUS ORIENTALIS.

Crocus optimus ex Anglia ad nos transfertur, in India orientali ipſe non creſcit, ſed ad Indos e noſtris regionibus transfertur.

Conſtat ⊖ibus fixis & volatilibus fragrantiſſimis & penetrantibus junctis blando oleo ſeu unctuoſitati, hinc aromaticus eſt, anodynus & hypnoticus, diſcutit mirifice, & in tuſſi dyſpnœa, aſthmate convulſivo, in phthiſi, roborando & exhilarando corde, in menſium lochiorum & fœtus remora ut & in ictero in uſu habetur, item in ſyncope & palpitatione cordis, externe ventriculo applicatus eum roborat. Frontalibus admotus phrenitidem, maniam & cephalalgiam à frigido ortam tollit, membris paralyticis & induratis impoſitus mire diſcutit.

–n–us vini crocatus grumos ſanguinis ſub cute, ſolvit & diſcutit, immoderatior uſus vitandus eſt, quia ſtuporem, ſomnolentiam nimiam & aliquando etiam amentiam inferre ſolet.

Datur ſubſtant. ad ℈ß Eſſent. Extract. proſtat etiam emplaſtrum oxycroceum.

℞. Croc. or. gr. vij Cinam. ac. ℈ß. Myrrh. gr. viij Zedoar. ℈ß. ♁ diaph. ℈j. M. f. pulv. ad lochia promovenda.

℞. Croc. or. gr. vij borrac. ℈ij Myrrh. gr. xv. cinam. ac. ℈ß. M. f. p. pro ij doſ. in remora menſium & lochiorum.

℞. ▽ Meliſſ. ℥ij Croc. or. ℈ß. exhibe ad cordis palpitationem, idem quoque applica regioni cordis.

℞. Cro. or. ℈ß. ☐r. ⊕lat. gr. vij ſem. aquil. ʒß. cape ex vino contra icterum.

℞. Croc. or. ℈ß. Moſch. gr. v. cape ex vino in aſthmate convulſivo.

℞. Eſſ. croc. Caſtor. ambr. ▽ –n–uoſ. meliſſ. ana ʒj exhibe ad gt. xxx ex decocto vel ▽ Meliſſ. contra palpitationem cordis.

℞. Ex-

℞. Extract. Croc. ʒj. ▽ roſar. ℥j. ▽ ſolan. ℥iiij f. Epith. contra cephalalgiam, addita aq. nymphææ prodeſt contra maniam & phreneſin.

FLORES STOECHADOS.

In ſpecie inter aromata cephalica, pulmonica nervina & uterina recenſetur. Datur eorum conſerva & ſyrupus, in quibus tanquam vehiculo ſpecifica ad ſupra dictos affectus exhibentur.

℞. Aq. flor. til. ℥ij ſyr. ſtœchad. ℥j conſ. alk. ʒß. ⩪ Ceraſ. n. ʒij f. mixtura ad roborandum caput in Apoplexia.

℞. Conſ. fl. Stœchad. ʒj. ⊖ vol. C. C. ℈ß. f. bolus in capitis & uteri obſtructione.

FLORES SCHOENANTHOS.

Conveniunt cum planta propria.

FLORES TUNICÆ.

Caryophylli hortenſis flores, officinis flores tunicæ dicti, ⊖e blando temperato volatili aromatico ad caryophyllos aromaticos accedente gaudent, & propterea in roborando capite, ut & à Sim. Paulli Q. B. aliiſque in morbis malignis prædicantur, proſtat horum conſerva, aq. deſt. ſyrupus multiplici infuſione factus, quæ omnia ſtimulantur caryophyllis aromaticis.

℞. Conſ. flor. tunic. ʒj caryoph. ar. gr. vj ☍ d. caryoph. gt. j. M. f. bol. ad roborandum caput, add. ambr. gr. iij vel iiij in ⩪ roſar. ſolut.

℞. Decoct. flor. tunic. exhibe contra febres ex obſervatione multorum practicorum.

FLORES VIOLARUM.

Sunt violæ martiæ purpureæ flores pentapetali oblongi purpurei, poſterius obtuſo calcare notati. Temperatum & blandum volatile aroma in tenui laxante viſcoſitate poſſident, propterea conveniunt in tuſſi, aſperitate faucium, ariditate linguæ, ⊖ibus acribus, catarrhis, it. in phthiſi, pleuritide, in fallenda ſiti, in affectibus cordis ab acido ⊖e terreo ſeu melancholia ortis, it. in febrium nimio æſtu temperando. Inter cardiaca roborantia numerantur tam ob gratum odorem ſeu blandum volatile aroma, quam ob tenuem viſcoſitatem terrea rigida ſolventem & lenientem.

Colligendi ſunt matutino tempore dum rore adhuc madent. Calices in quo vis nauſeoſa eſt rejiciendi, ſin autem vis potior laxans expetitur, calices cum floribus in uſum ſunt vocandi.

Datur conſerv. tinct. ſyr. ſimplex. & comp. it. ſucc. aq.

℞. Mann. q.ſ. affund. ſucc. violar. ad conſervæ conſiſtentiam add. novos flores recentes & fac conſervam cujus ʒj ad ʒij data laxat alvum.

℞. Conſerv. violar. ʒj Mann. ℥j ſyr. violar. q. ſ. f. laxans.

℞. Flor. violar. infund. aq. borrag. bugloſſ. q. ſ. add. tantillum ♎ ♆is f. tinctura.

℞. Syr. violar. ex multiplic. infuſ. aq. violar. cum ſacchar. ſ.q. f. tabulatum ad morbos pulmonum, tuſſim, aſthma.

FLORES ROSARUM PALLIDARUM.

Florum vis temperata alvum laxans, bilem mulcens & aquas evacuans conſiſtere videtur tum in rore, quo flores matutino tempore madent, tum in odore ſeu particulis volatilibus, quæ à rore tanquam menſtruo ⊖ino ⌽troſo ſuſcipiuntur, reliquum vero totum terreum eſt & adſtrictorium. Cæterum in volatili illo odore reſidet quoque vis analeptica, propterea tota ſubſtantia florum laxat, adſtringit & ſimul roborat, in ſpecie vero convenit bilioſis, hydropicis, & in fluxu albo mulierum.

Aq. deſt. habet vim roborantem & refrigerantem, quæ ex roſis damaſcenis eſt aqua, laxat ſimul alvum ſi paulo largiore doſi exhibeatur. Datur conſerva, ſyrupus, ♎ ardens. Aq. deſt. decoct. ung. ⁂ deſtil. & infuſ. Ob vim purgantem non exhibentur in decocto, quia vis purgans conſiſtit in volatili, frequentius dantur in infuſo.

℞. Roſar. pallid. rec. matutino tempore collectar. q. v. adde mannam & tantillum roſarum pallidarum, fac conſervam purgantem.

℞. Conſerv. roſ. pallid. ℥ß. pulv. hepat. rubr. ℈j ſcammon. roſat. gr. v. ♎ roſar. q. ſ. f. bolus laxans, convenit in febribus intermittentibus, præcipue in declinatione.

℞. Aq. deſt. roſ. ℥j. verben. ℥ij acet. roſ. ʒij ſantal. rubr. ʒß. f. epith. fronti applicandum.

℞. Aq. roſar. additur mixturis & emulſionibus cardiacis pro lubitu,

FLORES TILIÆ.

Aroma ſuum volatile in temperata tenui viſcoſa ſubſtantia fovent, unde capiti, nervis & ſpiritibus dicantur, & in ſpecie conveniunt in vertigine, convulſione, epilepſia, ſuffocatione hypochondriaca. Externe commendatur eorum decoctum pro cataplaſmate in teneſmo.

Præſcribitur ▽. decoct. & ∽us.

℞. ∽ lil. convall. flor. til. ana ℥j aq. Cinam. ʒiij conf. alkerm. ʒj ſyr. flor. tunic. ℥j M. f. mixtura ad epilepſiam add. ∽ Ceraſ. nigr. ʒiij.

LILIUM CONVALLIUM.

Virtute coincidunt cum floribus tiliæ ſed præſtantiores ſunt, externe naribus admoti ſternutationes provocant. Dantur ut flores tiliæ.

℞. Aq. flor. til. convall. lavend. roriſmar. primul. veris. ∽ ceraſ. n. ana q. v. cape pro uſu interno & externo in morbis convulſivis.

PRIMULA VERIS.

Tiliæ & lavendulæ floribus reſpondent, & in ſpecie laudantur in paralyſi, iis quoque qui tenerioris ſunt ætatis & complexionis hypnotici ſunt & anodyni. Dantur ut flores tiliæ.

℞. Aq. prim. veris flor. til. lil. convall. ana ℥ij aq. apopl. ∽ ceraſ. n. ana ℥ß. add. ⊖ vol. ſuccin. ʒß. ſyr. flor. tunic. ℥j f. mixtura contra apoplexiam & paralyſin.

FLORES

⁂

FLORES MUCAGINE TENUI POLLENTES.

FLORES MELILOTI.

Blandum diſcutiens ⊖ volatile temperatum quaſi unctuoſitate ſociatum habent, hinc interne inter expectorantia recenſentur, potior tamen eorum uſus eſt externus, diſcutiunt, emolliunt, dolores mitigant, hinc in inflammationibus quibuſvis præcipue uteri, ani, teſtium, ut & in doloribus nephriticis & podagræ conveniunt.

Præſcribuntur in cataplaſmate, fomentis emplaſtris; junguntur floribus chamomillæ.

℞. Flor. melil. chamom. fol. alth. ana Mj rad. alth. ℥j ficuum n. vij coq. in aqua vel lacte pro gargariſmate in inflammatione faucium maturanda, reſiduum applica partibus induratis emolliendis.

FLORES SAMBUCI.

Hujus flores ⊖e diſcutiente adeo potenti gaudent, ut ſimul ſudores moveant, ſi recentes ſunt, & frixi cum butyro exhibentur alvum blande ſubducunt, & ſi ex ſero lactis coquuntur hydropicis conducunt. Horum decoctum commendat Langius in augendo lacte mammarum. Externe in eryſipelate applicantur feliciter ut & in cephalalgia & nimiis vigiliis, nam ob demulcentem virtutem blandum ſomnum accerſunt.

Dantur in decocto vel ex aqua vel ſero lactis facto, aq. deſtill, laudatur in Variolis. Externe ingrediun-

diuntur clyſteres & cataplaſmata; inſpergitur pulvis cum creta & ſacch. ♄ni in eryſipelate.

℞. Flor. chamom. Sambuc. Melil. coq. c. aq. ad ileum, reſiduum applica ad calculum renum, idem quoque conducit pro clyſtere in colica paſſione.

***

FLORES SALE FIXO CUM PAUCIS VOLATILIBUS CONSTANTES.

FLORES CHEIRI.

Sunt Leucoji vulgaris flores lutei, ſaporis ſubacris ſubamari. In ſpecie utero dicantur, in pellendis menſibus & lochiis, laudantur quoque tanquam nervinum & diſcutiens in paralyſi & apoplexia. Datur conſerva & 🝆 ex florum infuſione, quod in Italia ad facilitandum partum regioni pubis inungitur.

℞. Conſerv. flor. cheir. cape ex aqua propria deſt. ad apoplexiam & paralyſin præſervandam, add. 🜔 volat. ſuccin. gr. v. & 🝆 d. caryoph. gt. j.

CALENDULA.

Sunt Calthæ vulgaris flores aurei diſcoidei radiati, ſaporis acris nauſeoſi, conſpirant cum floribus Cheiri, ſed in menſibus pellendis & fœtu tantæ efficaciæ cenſentur, ut & menſes calendularum odorem ſpiraſſe nonnulli affirment. Cæterum virtutem alexipharmacam habent, & in variolis, morbillis, peſte aliiſve malignis, ut & ictero conducunt.

Externe applicantur verrucis mollioribus tollendis, aliàs additur ☿ ♌ti gr. j. interim vicinæ partes muni-

muniantur emplaſtro diapalm. vel de gratia Dei; in hoc caſu etiam poſſunt applicari c. urina vel ▽ propria.

Datur conſerva. decoct. acet. calend. quod frontalibus pulſui & plantis pedum applicatur. it. ad capitis dolores in peſte commendatur.

℞. Succ. flor. calend. ℥j pulv. lumbr. terr. ʒj exhibe contra icterum.

℞. Conſerv. fl. calend. ʒj. ⁂ cinam. gt. j. f. bolus in remora menſium.

* * *

FLORES AQUEIS CONSTANTES.

BORRAGO & BUGLOSSA.

Utriuſque flores cardiacis annumerantur, dum ſua aquoſitate ⊖ia acida terreſtria ſeu Melancholica ♌us ſuffocantia diſſolvunt, hinc melancholicis & hypochondriacis conveniunt; proſunt quoque ſpecifice ad menſium ſuppreſſionem.

Datur conſerva ex floribus recentibus parata, ut & ▽ deſtillata & ſyrupus.

℞. Conſerv. bugl. borrag. ana ʒj cape c. ▽ propria ad melancholiam, add. tantill. aq. cinam.

℞. Conſ. borrag. ℥j cape cum vino albo ad menſium emanſiones.

℞. Flor. bugloſſ. infund. vino ad deſperatam epilepſiam ex mente Williſij.

FLORES CYANI.

Ejuſdem ſunt virtutis cujus ſunt flores borraginis & bugloſſæ, potentius tamen urinam pellunt, qua-

propter hydropicis cenſentur utiles. Camerarius pulv. ʒj feliciter exhibuit in ictero. Decoctum florum cyani & calendulæ datur ad urinam promovendam.

℞. Aq. deſt. fl. cyan. chærefol. ana ℥iij add. 🜍 vin. camphorat vel tinct. Croc. tantill. applica linteis exceptum oculorum ſuffuſioni.

⁂

FLORES MUCAGINE CRASSA CONSTANTES.

FLORES MALVÆ ROSEÆ ſeu HORTENSIS.

Ob viſcoſitatem & vim adſtringendi quam in receſſu relinquunt in ſpecie medentur inflammationi & exulcerationi faucium, gingivarum & uvulæ. Commendatur quoque ad eryſipelas ſub forma cataplaſmatis, in faucium vero affectibus præſcribitur ſub forma gargariſmi.

℞. Flor. Malv. coq. in aq. add. par. aceti. & ſyr. violar. pro gargar. ad faucium inflammationes add. par. ⦶tri q. ſ. ad gratiam.

℞. Pulv. flor. malvæ ʒj laccæ, myrrh. ann ℈j mell. q. ſ. & tantillum conf. alk. f. liniment. contra putredinem gingivarum ſcorbuticam, add. quoque tantillum 🜍us cochleariæ.

VERBASCUM.

Reſpondent foliis, quibus tamen præſtantiores ſunt ℞. Flor. verbaſc. pone in cella ut liqueſcant, cape 🝆 & applica omnibus tumoribus, præcipue lienis, hæmorrhoidum arthritidis & podagræ.

POEONIA.

POEONIA.

Sunt flores rubentes, ſaporis fatui, odoris nullius. Conveniunt cum radice & ſeminibus. Conſtant particulis aquoſis mucilagine volatili & craſſa, ob mucilaginem volatilem antepileptica eſt, raro decoctum vel infuſum paratur, ſæpius conſerva utimur pro vehiculo.

℞. Conſerv. flor. pœon. ʒj. ⊖ volatil. C. C. gr. v. ſyr. pœon. q. ſ. pro bolo ſ. Elect. epileptico.

Contra vertiginem ℞. conſ. flor. pœon. ʒj. rad. galang. gr. xv. caſtor. gr. ij ſyr. pœon. q. ſ.

PAPAVER ERRATICUM.

Ob mucaginem volatilem cui quidpiam acredinis & amaritiei eſt conjunctum, blande diſcutiunt, moderate incraſſant, leniunt, mulcent, & ſomnum inducunt, propterea conveniunt in pleuritide aliiſque pulmonum & thoracis affectibus, ortis ab acri tenui materia, magis tamen in principio ubi dolor urget quam in ſtatu & declinatione conferunt, ſiſtunt quoque nimias hæmorrhagias & nimium menſium fluxum.

℞. Conſ. flor. papav. ʒj. mandub. luc. piſc. ʒß. ⊖ volatil. C.C. gr. vij. L. O. gr. j. ſyr. papav. q. ſ. f. bolus contra pleuritidem.

℞. Aq. flor. papav. nymph. ana ℥j ſem. papav. alb. ʒij f. Emulſ. add. ſyr. violar. vel papav. contra nimias vigilias.

℞. Flor. beton. papav. ana p. ij ſaſſafr. ℥ij f. decoct. ex aq. colatur. add. ſyr. papav. q. ſ. contra phthiſin.

℞. Flor. papav. infunde aquæ propriæ deſt. exhibe contra nimium fluxum menſium.

***

FLORES TERREIS CONSTANTES.

FLORES BALAUSTIORUM.

Vi adſtringente & roborante conveniunt cum corticibus querc. granator. aliiſque congeneribus.

℞. Flor. balauſt. ʒj pulv. Malicor. ℈ij conſ. roſar. ⊖lat. ℈j ſacch. ♄n. gr. v. ☍ N. M. expr. ʒß. L. O. gr. ij ſyr. granat. q. ſ. f. boli in diarrhæa.

℞. Pulv. flor. balauſt. inſperge intertrigini & cuti exulceratæ.

FLORES NYMPHÆÆ.

Conveniunt cum radice propria.

℞. Sem. pœon. ℥ij. aq. Nymph. bugloſſ. q. ſ. f. Emulſ. add. ⦶tr. præpar. ℈j & tantillum ſyr. nymph. ad nimiam ſalacitatem.

℞. Conſ. flor. nymph. ʒj camph. gr. v. L. O. gr. j. ſacch. ♄ni ℈ß. ſyr. nymph. q. ſ. f. bolus contra maniam.

FLORES ROSARUM.

RUBRARUM.

Sunt roſæ rubræ C. B. P. flores ampli, modo ſimplici modo multiplici petalorum ſerie conflati, ſaporis ſi ſiccati adſtringentis & auſteri, ſi recentes ſubacris ſubamaricantis ingrati. Quia minus odorati ſunt terreis conſtant, & ubicunque adſtrictione opus eſt mire conducunt, præcipue ſi cum acido exhibeantur, laudantur quoque ad compeſcendum vomitum, it. in phthiſi ad conſolidanda ulcera, ut & in hæmoptyſi.

Para-

Paratur nobile balſamum roſarum ex 🜔 proprio deſtillato apud Perſas.

Conſtant quibuſdam particulis ⊖inis volatilibus cum prædominio tamen terrearum.

Datur conſerva, ſyrupus ut & pulvis.

℞. Conſerv. roſar. r. ℥j. 🜍 ⊕li g. v. L. O. gr. j. 🜔 N. M. expreſſ. ʒj f. bolus in omnibus fluxibus. In nimio menſium fluxu conſerva roſarum ad ʒj cum ſyrupo cydoniorum commendatur, præcipue ſi menſes doloroſi ſunt.

℞. Conſ. roſar. rubr. ʒj ſem. pap. alb. ℈j ſyr. roſar. f. bolus contra hæmoptyſin.

℞. Corall. r. præpar. ʒj Crem. ♀ri. ʒß. pulv. roſar. r. ℈j. 🜍 ⊕li ſantal. rubr. tincti q. ſ. f. pulvis ad æſtus vagos & phlogoſes reprimendas.

℞. Flor. roſar. r. affund. acetum, exhibe in quibuſdam dyſenteriis, præcipue ſi à bile acri oriuntur.

℞. Pulv. roſar. rubr. ʒij cape ſucceſſive ex aqua plantag. ad contumaciſſimam diarrhæam & dyſenteriam.

℞. Flor. roſ. r. balauſt. ana ʒß. N. M. toſt. ℈ß. L. O. gr. j. in dyſenteria maligna.

℞. Conſerv. roſ. ʒij pulv. Zedoar. ℈j ſucc. cydonior. q. ſ. f. Elect. in pica add. parum 🜍us ⊕li vel ſucc. limon. vel acetoſitat. citri.

Aqua deſtillata interne blande refrigerat, hinc in ſiti febricitantium. ℞. Decoct. hord. ℔j aq. roſar. ℥iiij ſyr. acetoſ. Citr. q. ſ. add. ⏀tr. depur. vel lap. prunell. ʒj ſyr. roſar. ℥j pro Julapio.

Externe in inflammationibus oculorum. ℞. Aq. roſar. fœnicul. ana ℥j ſacch. ♄ni ʒj ſacch. cand. ʒiß camphor. gr. iij extr. croci gr. ij f. collyrium.

CAPUT OCTAVUM.

DE SUCCIS RESINIS ET GUMMATIBUS.

SUccus dicitur, quod prælo ex planta vel arbore exprimitur. Reſina eſt corpus oleoſum craſſum quod tamen in oleoſis & ſpirituoſis ſolvitur. Gummi eſt pars aquea congelata in mucilaginem craſſiorem, quæ tamen in aqua ſolvitur. Dantur ceu amphibia, dicunturque gummi reſinæ, quæ partim in aqueis, partim oleoſis & ſpirituoſis ſolvuntur; reſina autem ex iis extracta, aliarum ſtatim genium induit reſinarum, tantumque in [illegible]uoſis ſe ſolvi patitur.

SEQUENTIA HABENT PARTICULAS SALINAS VOLATILES OLEOSAS CUM MUCILAGINOSIS TENUISSIMIS SUBTILIBUS.

CAMPHORA.

Prima eſt Camphora noſtra officinarum, Chinenſis vulgo dicta, licet nulla in China inveniatur, eò quod ab his olim ad nos tantum profecta ſit: Eſt ſubſtantia candidiſſima, facie ſalis vel nivis, pinguis & oleoſa, vi ignis ſublimata, ex ligno arboris camphoriferæ Japonenſis & compacta in glebas; ſaporis acris amaricantis, aromatici, pinguis & lenti, odoris fragrantis & gravis. Eſt ſcilicet arborum camphoriferarum in Japonia, quæ ſunt folio rigido virideſcente & fere lauri-

laurino, flore pentapetalo, baccis bifidis: radix, truncus, rami, folia fiſſa conjiciuntur in ahenum, cui imponitur capitellum ſeu conus ex ſtramine qui uſitatior, vel ſtoreis, vi ſubſtrati ignis quicquid volatile eſt expellitur, & lateribus coni ſe applicat, quæ dum levis inſtar fuliginis decidit, quæ deinde in maſſam cogitur & ad nos transfertur.

Secunda ſpecies eſt, quæ in inſula Zeylon conficitur, paratur ex radice ſeu arctius cortice radicis cinamomi, hic nempe ſeparatus à materia lignea immittitur veſicæ inſtructæ ſuo capitello, vaſe ſerpentino & recipiente, affunditur ▽ & deſtillatur, hinc obtinetur 1. aqua. 2. oleum & 3. oleo ſupernatans camphora, quæ ab oleo optime expreſſione manuum ſeparatur.

Tertia ſpecies ex variis plantis Indiæ orientalis elicitur, ſic ex radice Zedoariæ recenti, modo ſupra in ſecunda ſpecie deſcripto obtinetur aqua, oleum & huic ſupernatans Camphora, quæ eadem methodo ſeparatur. Hinc radix Zedoariæ deſtituta ſuo ſale volatili cum ad nos defertur optime Camphora ſtimulatur. Sic ex Menthæ quadam ſpecie camphoram mihi quoque comparavi. Atque hæ tres ſpecies ſunt camphoræ factitiæ, quarum duæ poſteriores non niſi à curioſioribus aſſervantur & poſſidentur.

Quarta nativa eſt in Inſula Borneo, quæ in arboribus mole tiliis, configuratione florum, foliorum & baccarum arboribus camphoriferis paribus, cæſis truncis in partes diviſis in ſpaciolis quibuſdam forma minutiſſimarum lamellarum lenticularium, quæ unciæ Indis vocantur reperiuntur in rara ſatis copia: duæ namque vel tres ſæpe arbores cœſæ vix ℥ij vel iij ſuppeditant. In Oriente maxime ad Japonas, quibus domeſtica

meſtica ſordet Camphora diſtrahitur, tantum pro libra una 100 ſolvuntur floreni, ad nos ob pretium non transfertur.

Blandam ſpiritibus requiem ob ſalem ſuum ſubtiliſſimum inducit, & ſimul acida, viſcoſa, terrea craſſave potentiſſime diſcutit, hinc ſudoriferum eſt & anodynum univerſale, quibus nominibus ſpecifice convenit in peſte, malignis, deliriis, melancholia, paſſione hyſterica & motibus convulſivis, gonorrhœa, &c. doſ. in ſubſtantia à gr. v. ad vij & x.

Jungitur tam internis quam externis medicamentis tanquam ſtimulus. Sic e. c. in Catarrhoſis cum capitis dolore & alvo adſtricta ℞. reſin. jalap. gr. x. extr. troch. alh. gr. iij ☍ ſuccin. gt. j. camph gr. v. balſ. Peruv. q. ſ. f. pil.

Singularis eſt in febribus malignis omnibus, item tam continuis quam intermittentibus: ſic in febre acuta cum alvi fluxu. ℞. C. C. præp. ʒß. L. O. gr. j. Camphor. gr. v. f. pulv.

℞. ♁ diaph. ℈j. ⊖ vol. ſuccin. gr. v. Camphor. gr. ij f. pulv. contra febres malignas, exhibe ex ▽ Card. B. vel theriacali.

Specifica eſt cum Laud. Op. & extracto Croci, in mania, deliriis, melancholia. Sic ℞. ▽ flor. ſambuc. hyperic. ana ℥j camphor. gr. v. extr. Croc. gr. iij. L. O. g. j. M. vel in forma pulveris: ℞. Arcan. duplicat. ♁ diaphor. ana ℈j. ſacch ♄ni ℈ß. camphor. gr. v. L. O. g. j. M. f. pulv. vel excipe conſerv. flor. nymph.

℞. Laud. op. gr. ij. camph. gr. iij f. pil. ad præſervationem epilepſiæ.

℞. Camph. ℈ß. Myrrh, ʒß. gum. tragac. cum aceto in mucagin. redact. f. trochiſci antipeſtilentiales.

Exter-

Externe additur unguentis ceu ſtimulus pro incidente & anodyno. ℞. Ung. dialth. ℥j camphor. ʒj pro tumoribus emolliendis.

℞. Empl. diapomph. ℥j. Croc. or. ℈j camph. ʒß. 🝆 philoſoph. q. ſ. f. emplaſtr. in tumoribus diſſipandis.

Paratur etiam Epithema egregium manuum febrifugum contra tertianas agyrtis commune, quod ſæpe effectu ſuo non caruit.

℞. Fulig. ſplendent. ℥iß. thereb. ʒvj. tel. aranear. ʒj camphor. ʒß. 🝆 aranear. vel ſcorpion. q. ſ. pro epithemate, applicatur carpis manuum.

Utimur etiam ∽u theriacali Camphorato it. eſſentia ejus in omnibus febribus malignis. ℞. ▽ flor. ſambuc. Card. B. ana ℥ij. ∽ theriac. camphorat. ℥ß. L. O. gr. ij ſyr. acetoſ. Citr. ℥j.

℞. ▽ roſac. ℥ij. ∽ vin. camphorat. ʒj. f. colyr.

OPIUM.

Eſt ſubſtantia partim reſinoſa, partim gummoſa, dum recens eſt mollicula dum inveterata durior, collecta ex papavere & in maſſam redacta, ſaporis ſubacris amari, odoris gravis viroſi narcotici & quaſi ſtupefacientis.

Aliud eſt Thebaicum, aliud Indicum, prius eſt ſubtilius & in Ægypto circa Cairo & confines tractus colligitur, poſterius eſt craſſius, fœculentius & in Inſulis Bengalenſibus colligitur. Scilicet utrinque agri, imo integri tractus conſeruntur ſemine papaveris hortenſis, ubi per æſtatem ad maturitatem vergunt diſcuſſis floribus ſcalpello 5 vel 6 locis ſauciant capita, è quibus lacteus ſtillat humor, qui idem in maſſam

coactus ad nos transfertur. Unum caput papaveris ultra ij vel iij grana vix ſuppeditat.

Thebaicum fere tantum apud nos officinale, uti in oriente proprio ſolo enutritum. Opium album vulgo Maſlac ſeu magnanimum, eſt opium ab omnibus ſordibus depurgatum, quod carioris eſt pretii. Meconium aliud non eſt quam papaver coctum expreſſum, filtratum & inſpiſſatum.

Univerſale eſt remedium in omnibus morbis ubi opus eſt ſomnum vel torporem inducendi, ubi conjuncti ſunt dolores, ubi acredo obtundenda, ſiſtit etiam functiones animales impeditque ſpiritus quo minus in organa influere poſſint.

Datur per ſe à gr. j. ad iij in aſſuetis ℈j imo ʒß in India non nocuit.

Vis ejus narcotica ſuſpecta corrigitur optime toſtione, ſic enim multum 🜔is volatilis perit, ſed nota ſtatim eſſe exhibendum, alias ad genium & malignitatem ſuam redit, reanimatum ab aëre. Alias etiam ſucco Citri vel cydoniorum caſtratur.

Nonnulli præcipue recentiores corrigunt caſtoreo, quod quaſi eſt opii antidotum. Item ℞. opii ʒj. Caſtor. ʒß. Camph. ℈ß. ſpirit. vin. 🜄iſat q. ſ. f. tinctura, quæ filtretur & inſpiſſetur. Omnium vero mitiſſime agit opium leviter toſtum.

Additur etiam clyſteribus in colica, ſed decoquitur in nodulo incluſum, ne ſe applicans inteſtinis tormina excitet.

Additur purgantibus validioribus pro correctorio præcipue in colica. Sic ℞. Extr. agaric. ℈j. Extr. troch. alhand. gr. iij. 🝆 expr. flav. cort. aur. gt. iiij Caſtor. gr. iij L. O. gr. ij f. c. Balſ. Peruv. pil. vel cum conſ. fl. anthos bolus. ℞. Opij

℞. Opii toſt. gr. ij Nuc. Moſch. toſt. ʒj f. pulv. in dyſenteria.

℞. Opii toſti gr. iij. ☍ Nuciſt. expr. ℈ß. Corall. r. præpar. ℈ß. rad. tormentill. ℈j conſerv. roſar ⊕lat. q.ſ. f. bolus in dyſenteria.

℞. Opii præp. gr. j. Camph. gr. iij conſ. Nymph. ʒj f. bolus in delir. & mania add. ſacch. ♄n. gr. ij vel iij.

℞. Reſin. jalap. gr. xv. extr. lign. aloës ℈ß. ☍ ſuccin. gt. ij opii gr. iij. balſ. Peruv. q. ſ. f. pilulæ purgantes & anodynæ, poſt alvi evacuationem placidum inducentes ſomnum.

Externe poma in manu calefacta naribuſque admota narcoſin inducunt. v. c. ℞. ſtyrac. calam. L.O. ana ʒiß. lign. aloës rad. ir. flor. ana ʒiij.

In hæmorrhoidibus exulceratis inflammatis, teneſmo dyſenterico ſequens unguentum profuit. ℞. Ung. alb. camphorat. ℥j L. O. ʒj ſacch. ♄ni. ʒij. ☿ viv. ʒß. ☍ roſac. vel quod ſæpe præſtat lini q. ſ. quod exceptum ſtupis vel filamentis linteaminum, præmiſſis univerſalibus intruditur ano ad partem uſque affectam.

***

CONSTANTIA PARTICULIS SALINIS VOLATILIBUS OLEOSIS CUM VIRTUTE LAXANDI.

AMMONIACUM.

Eſt ſubſtantia gummoſa & reſinoſa ex variis denſis ſolidiſque grumis modo flaveſcentibus modo lacteis, qui optimi, collectis ex ferula ammoniacifera, quæ mole fere arborem æquat & lacteſcens eſt, ſplendidis & nitidis in glebam coactis, dum manibus tra-

ctantur lentore molleſcentibus, ſaporis acris, amaricantis, viſcoſi, lenti & paululum nauſeoſi, odoris fragrantis gravis, alliaris fere.

Ratione ⊖is volatilis oleoſi incidit & diſcutit & aperit, ratione reſinæ lenit & mollit, utriuſque vero ratione, in largiori doſi datum, alvum ſolvit, & proinde in tuſſi, aſthmate, hyſterica paſſione, hypochondriaca & ſcorbutica ſpecificum eſt. Externe tumores & ſcirrhos reſolvit, diſcutit & emollit.

Datur in ſubſtantia ʒj ſub forma emulſionis, electuarii, boli vel pilularum, ſtimulatur ♀ro, oleo ♀ri, ⊖e ♀ri, borrace &c. pro ſcopo vero alvum laxandi exaltatur trochiſcis alhandali & diagrydio. Præparant ad impuritates removendas ſolvendo in aceto, filtrando & inſpiſſando, in pilulis, bolis, emplaſtris, in potionibus vero melius ſolvitur in ipſo vehiculo non inſpiſſandum, alias periret multum ⊖is volatilis, v. c. in aſthmate: ℞. Gum. ammon. ʒß. ▽ Cinam. ſimpl. vel puleg. ℥ij ſyr. hyſſop. puleg. vel cinam. ʒiij.

℞. Gum. ammon. purgat. ʒj ſolv. in ▽ Cinam. ℥iiß. f. hauſtus contra tuſſim, vel loco ▽ Cinam. cape ▽ puleg. vel menthæ.

℞. Gum. ammon. purg. ʒj ♀r. ⊕lat. ℈j ▽ fumar. Meliſſ. ana ℥iß ſyr. cochlear. ʒij, f. hauſtus in paſſione hypochondriaca & ſcorbutica.

℞. Gum. ammon. ſal. ammon. ana q. v. deſtilla ex retorta, cape ⌒um ad gr. x. xx cum ▽ deſtillata vel ſyrupo ad tuſſim, aſthma, ſcorbutum.

℞. Gumm. ammon. ʒj ⚭ caryoph. & tantundem ⚭ amygdalarum f. linimentum ad reſolvendos tumores frigidos.

℞. Gum

℞. Gum. ammon. ʒß. troch. alhand. ℈ß. Elix. propr. q. ſ. f. pil. laxantes, pro aliquot doſibus in ſcorbuto & paſſione hypochondriaca.

Habentur in officinis pilulæ de ammoniaco, quæ ſunt aſthmaticæ hypochondriacæ & uterinæ, pro cujus obſtructionibus tollendis egregie maritantur purgantibus. ℞. Pil. d. ammon. ʒß. extr. troch. alhand. gr. j. ⊖ vol. ſuccin. gr. v. ℅ d. ſuccin. gt. iij f. pil. Emplaſtrum de ammoniaco impoſitum in dolore iliaco, iſchiatico lumbarique commendatur junctum cum emplaſtro de cicuta anatica portione.

Externe in doloribus dentium à cavitate, pro iis extrahendis ſequens paro unguentum: ℞. Gum. ammon. ſem. hyoſc. ana ʒiß ſucc. hyoſc. cum pinguedine ranarum viridium ana q. ſ. f. unguentum, quod exceptum ſtuppis intruditur cavo dentium, ubi circumjectam carnem quaſi ſolvit, ac proinde extractionem dentis multum facilitat.

GALBANUM.

Eſt ſubſtantia gummoſa & reſinoſa, pinguis, ceræ inſtar ductilis, ammoniaco ſimilis, aſt ex variis albentibus & ſplendentibus grumis in maſſam coacta, collecta ex ferula Galbanifera Syriaca, ſaporis ſubacris, ſubamari nauſeoſi, odoris fragrantis gravis, odor & ſapor idem qui ammoniaci, mitior tamen, tantumque gradu differunt.

Virtute nonnihil inferior eſt gummi ammoniaco, ſed in ſpecie valet in colica hyſterica & affectibus convulſivis, datur ut ammoniacum.

℞. Maſſ. pil. d. galban. Sylv. ʒß diagryd. gr. v. f. pil. purgantes hyſtericæ.

℞. Galban. q. v. ſolv. in + ſeu ⁔u vini, ſoluto add. extract. Caſtor. Croc. ana ʒiij. ☍ d. ſuccin. ſpic. ana ʒiß. M. applica umbilico ad paſſionem iliacam.

BDELLIUM.

Eſt ſubſtantia gummoſa & reſinoſa, pinguis, ceræ inſtar tenax, glutinoſa, coloris ex flavo ferruginei, collecta ex arbore Americana ſpinoſa, ſaporis acris, pinguis, lenti, amaricantis, aromatici, odoris fragrantis.

Conſtat ⊖ibus volatilibus cum reſinoſis multis. convenit cum ammoniaco in virtute niſi quod tanta vis catharctica non inſit, ſed potius adſtrictoriam in receſſu relinquat.

℞. Pil. d. bdell. ʒß. ſacch. ♄ni ℈j balſam. Peruv. q. ſ. f. pil. in hæmorrhagia hæmorrhoidum.

Externe paratur emplaſtrum, linimentum, pro mollienda duritie & nodis nervorum, pro membris convulſis. ℞. g. bdell. myrrh. ana ʒj ſolv. in ☍ laurin. expreſſ. ſuper carbon. add. ☍ ſuccin. ſpic. ana ʒß. ☍ ſcorpion. lumbricor. & ceræ q. ſ. pro conſiſtentia linimenti vel emplaſtri.

Odoramenta uterina etiam ingreditur. ℞. Bdell. myrrh. ſuccin, ana ʒj injiciantur carbonibus & fumus recipiatur per muliebria, ad lochiā, fœtum, menſes pellendum. Oleum deſtillatum ſpecificum eſt contra obſtructiones Uteri.

ASA FOETIDA.

Eſt ſubſtantia gummoſa & reſinoſa, aridior, ceræ inſtar ſequax, ex variis ſplendentibus grumis, modo albeſcentibus, modo flaveſcentibus modo elegantiſſime incarnatis collectis ex tithymalo Perſico fœtido ſarmen-

ſarmentoſo, ferula Perſica rectius dicto, in maſſam coacta, ſaporis acris mordacis lenti, aliquantum nauſeoſi, odoris fragrantis gravis allium mentientis. Indi loco aromatis in cibis uſurpant.

Ammoniaco & Galbano potentior eſt & ſpecifica in colica & hyſterica paſſione, ſudorifera eſt in variolis & morbillis, febribus malignis, odor naribus exceptus hyſtericam paſſionem ſiſtit. Datur à ʒß ad ʒj.

℞. Aſ. fœtid. ℈j. ⊖ vol. ſuccin. gr. xv. borrac. ℈ß. f. cum balſ. Peruv. vel Elix. propr. pil. in menſibus lochiis & fœtu pellendo.

℞. Aſ. fœtid. q. v. infund. in ♒ vin. ⊡iſat. q. ſ. f. tinctura antihyſterica, cujus guttas xx vel xxx exhibe in aqua florum ſambuci.

℞. Ung. alcis ʒij aſ. fœtid. ʒj verruc. equor. ʒj vel ij f. pulv. injiciendus carbonibus candentibus ad ſuffocationem uteri.

℞. Aſ. fœtid. ℈j. ⊖ vol. ſuccin. ℈j L. O. gr. j. conſerv. flor. ſtœch. ad ʒj f. bolus uterinus.

℞. Galban. ʒj. aſ. fœtid. ʒß. Caſtor. ℥ß. °° ſuccin. cer. alb. q. ſ. f. pomum auro obducendum, quod naribus ſæpius admotum paſſionem hyſtericam impedit.

℞. Aſ. fœtid. q. v. album ovor. q. ſ. applica panaritio.

SAGAPENUM.

Eſt ſubſtantia gummoſa & reſinoſa lacteſcens, in grumos foris ex rufo flaveſcentes intus candidos concreta, collecta ex ferula Ægyptiaca ſagapenifera, ſaporis acris mordacis nauſeoſi, odoris viroſi ſi recens, fragrantis gravis.

Conſtat virtute laxandi cæteris majore, tum datur ad ʒiij alias ubi alterare debet ad ʒj.

Mitius

Mitius eſt aſa fœtida, & interne in retentis lochiis, fœtuque mortuo pellendo uſurpatur.

℞. Ammoniac. opopan. ſagap. galban. ana ℥j. ☿ philoſophor. q. ſ. f. Emplaſtr ad ſuppurandas hernias carnoſas, additur magnes arſenicalis & bdellium.

⁂

RESINÆ SPISSIORES ET SOLIDIORES ARIDIORES, SALE OLEOSO CONSTANTES.

GUMMI ANIME ET CANCAMUM.

Eſt ſubſtantia reſinoſa, arida, forinſecus retorrida, alba, ſubinde farina aſperſa, intus ex candido flaveſcens, ſuccino ſimilis, facile friabilis, collecta ex arbore Jetoiba in occidentali & orientali India. Sponte naſcitur in inſula Minæa, unde antiquitus tantum ad nos transfertur, hinc forta anime dicitur. Saporis prima fronte nullius evidentis, hinc pinguis reſinoſi, odoris dum accenditur fragrantis jucundi, accedentis ad odorem thuris vel olibani.

Ratione odoris cerebro & ſpiritibus gratiſſimo annumeratur cephalicis, nervinis & anticatarrhalibus, quoniam vero ſuum volatile habet in reſina duriſſima in ventriculo vix ſolubili, ideo præſcribitur tantum in ſuffitu, ſi interne velimus uſurpare volatilibus ſtimulatur vel oleis.

℞. Anim. ℥j Maſtich, benz. ana ℈ij Ladan. ʒß. M. f. p. pro ſuffitu in Catarrhis.

℞. Lign. al. ʒj anim. benz. ana ʒß. f. pulv. pro ſuffitu ad corrigendam aëris malignitatem.

Parantur

Parantur & flores ex hac reſina, ejuſdem virtutis cum floribus benzoë, egregie nempe thoracici. ℞. Flor. g. anim. ℈j benz. gr. vij exhibe in ▽ Cinam. vel ſyr. aſthmatico.

SANDARACHA.

Eſt ſubſtantia reſinoſa, arida, pallide flaveſcens, ſplendida, in guttas vel ſtrias modo rotundas, modo oblongas maſtichi ſimiles concreta, collecta ex truncis juniperi vulgaris fruticoſæ, ſaporis reſinoſi, odoris dum accenditur fragrantis & ſuavis.

Conſtat ⊖ volatili 🝆ſo incluſo mucilaginoſis craſſis, cum acido volatili remiſſiore tamen quam in ſuperioribus gradu.

Cum g. animeconvenit, & contra catarrhos in ſuffitu tantum præſcribitur. Cum 🝆 ſuccini vel lini ſoluta medetur doloribus hæmorrhoidum, cocta cum 🝆 lini externe applicatur vulneribus. In multis etiam convenit cum maſtiche, pauciora tamen habet ⊖ia volatilia, hinc in defectu maſtichis in ſuffitu eo poſſumus uti.

℞. Sandarac. ʒiij Maſtich. ʒj benz. ʒß. raſ. ſuccin. ʒij f. pulv. pro ſuffitu in catarrhis & coryza.

MASTICHE.

Eſt ſubſtantia reſinoſa arida, dura, pellucida ex flavo palleſcens, granulata, modo oblonga, modo rotunda, profluens ex lentiſco Chienſi, odoris quaſi obliterati, dum accenditur fragrantis. Salia ſua volatilia reſinæ molliori facilius ſolubili & nonnihil conſtringenti habet incluſa, hinc in ventriculi debilitate, diarrhæa & dyſenteria convenit. Ratione acidi ſui

volatilis, ventriculi acidum ſuppreſſum erigit, unde prodeſt hypochondriacis ſcorbuticis; externe convenit in ſuffitu ad procidentiam uteri & ani.

Datur in granis iij vel iiij vel coct. ut & deſtillat. it. ſub forma emplaſtri & unguenti.

℞. Maſtich. gr. iij integra deglutiantur ad debilem ventriculum roborandum.

℞. Maſtich. ʒß pulv. Caryoph. ʒj roſ. rubr. p. ij vin. rubr. adſtr. ℔ß coq. in vaſe clauſo ad medietatem, ad ſiſtendum vomitum & diarrhœam. add. L.O. gr. j.

℞. Pil. Maſtichin. ℈j horis veſpertinis poſt cœnam ad movendam alvum in doloribus capitis à catarrhis, quibus add. pro ſtimulo diagryd. gr. v. troch. alhand. gr. ij.

℞. Medull. lapid. ʒiij ligni aloës reſinoſ. ʒiß f. pulv. cujus ʒß exhibe in incontinentia urinæ puerorum, & fiat ex ſolo maſtiche ſuffitus.

Paratur etiam ▽ Maſtichina capiendo ollam recentem, ſuſpendendo ſuper fumos maſtichis ita enim egregie imprægnabitur, huic deinde infunde vinum vel aquam utere pro potu ordinario, vel in liquoribus hiſce extingue accenſam maſtichen.

Si per retortam deſtillatur, phlegma prodit ſubacidum, ſi vero per veſicam aqua aromatica.

THUS ſeu OLIBANUM.

Eſt ſubſtantia reſinoſa ex flavo palleſcens & ſplendeſcens, in guttas maſtichi ſimiles ſed majores concreta & indurata, collecta ex arbore ſpinoſa Arabica thurifera, ſaporis amaricantis reſinoſi, odoris fragrantiſs.

Ratio-

Ratione figuræ eſt vel indicum, quod eſt in glebis, vel mammoſum, quod eſt in guttis, vel maſculum, quod eſt in granis.

Maſtiche fragrantius & ſuavius, cæteroquin cum ipſo convenit.

— ℞. Maſtich. Oliban. ana gr. v. Ocul. ♋. gr. xv. Corall. r. præpar. C. C. præpar. ana ℈ß. ☍ d. fœnic. gt. iij M. f. pulv. ad abſorbenda acida, ex lacte dandus, præcipue contra tormina puerorum.

— ℞. Thur. ʒj Maſtich. ʒß. ⊖✱ci. ℈j f. pulv. pro eroſis gingivis, ſuperſumendum decoctum ſalviæ.

LACCA.

Lacca in ramulis eſt ſubſtantia reſinoſa ſuperficie inæquali veluti granulata, dura, rufa pellucida, intus cavernoſa, elaborata à formicis alatis & circa extimos arborum pingueſcentium ramulos congeſta, ſaporis reſinoſi, odoris dum accenditur fragrantis ſulphurei, nonnihil acidi, & ſpiritibus noſtris mire grati.

Lacca in granis, ſunt grana reſinoſa, parva dura rubicunda, lapillis granatorum ſimilia, pellucida, collecta ex arbore Indica Kapethia dicta, floribus villoſis, ſemine tricocco, ricini ſpecie, ſaporis & odoris præcedentis.

Utraque in reſina ſua adſtringente fovet ⊖em volatilem qui cum accenditur ſeu incaleſcit ſe prodit, cujus ratione blandum diureticum, & præcipue in morbis chronicis, variolis & morbillis blandum & elegans diaphoreticum eſt. Externe tanquam ſpecificum ingreditur compoſitiones ad gingivarum & dentium vitia ſcorbutica. Datur ſoluta in ſpiritu vini,

℞. Lacc. q. v. affund. ⌢ vin. rectificat. relinque donec ⌢us vini ſufficienter tinctus fuerit, hanc tincturam exhibe cum ſyr. Card. B. vel emulſ. ſem. aquileg. ad expellendum variolas & morbillos add. quoque tinct. myrrhæ, eſſent. Caſtor. ⊖ volat. ſuccin. &c.

℞. Crem. 🜿ri ℥iij Lacc. ʒj. ⊕l. cypr. gr. vij f. pulv. dentifricius.

℞. Lacc. ʒiß Myrrh. alum. uſt. ana ʒj. phlegm. ⊕li q. ſ. ſtent in digeſtione f. tinctura, huic admiſce ſyr. cochleariæ, vel ſyr. gran. cherm. q. v. f. linimentum, ad gingivas carie ſcorbutica infectas.

SUCCINUM.

Eſt ſubſtantia reſinoſa in lapideam duritiem concreta, lucida, magnitudinis & figuræ variæ, tranſparens, flaveſcens, nata in terræ viſceribus Pruſſiæ mari Balthico vicinæ, ſaporis ſubacris & reſinoſi, odoris fragrantis dum accenditur. Eſt vel album vel flavum vel nigrum, præſtat omnibus album.

Habet ⊖ia volatilia oleoſa cum acidis volatilibus incluſis reſinoſis craſſis & terreis. Utrumque & album & flavum, polychreſtum medicamentum eſt, tum ob ſalem volatilem oleoſum incidentem & diſcutientem, tum ob reſinam temperantem, & adſtringentem, utriuſque nomine in catarrhis, morbis ſoporoſis, convulſivis, alvi fluxibus, gonorrhœa ſpecifice convenit. Datur à ℈j ad ʒß. in ſubſtantia, pulvere. Externe ſumitur ad ſuffimigia contra catarrhos, coryzam.

℞. Succin. præp. terr. ſigil. ana ʒß. Ocul. ♋ ℈ij diaſcord. Fracoſt. & ſyr. myrtin. q. ſ. ſumatur cum ∇ plantag. contra dyſenteriam. addi poteſt L. O. gr. ij vel iij.

℞. Troch.

℞. Troch. d. charab. ☍ N. M. expr. ana ℈j L. O. gr. ij conſerv. roſar. ⊕lat. ʒj M. f. bol. contra diarrhœam.

℞. Succin. præpar ʒß borrac. Venet. ℈j. ☍ cinam. gt. j. conſerv. fl. anth. ʒj f. bolus in menſium remora. Vel R. ſuccin. præpar. ʒß. borrac. calcinatæ ℈j omitte ☍ cinamom, cape cum conſerva flor. nymph. ad nimium menſium fluxum.

℞. Succin. præpar. oſ. ſepiæ ana ℈j datur ex ▽ flor. nymph. contra gonorrhœam.

℞. Succin. præpar. ℈ij vin. auſter. q. ſ. cape ad lochiorum fluxum.

℞. Eſſent. ſuccin. gt. xv. ▽ Cinam. ℥j f. lac. in menſium remora & paſſione hyſterica & aliis debilitatibus nervorum & ſpirituum.

℞. Eſſent. ſuccin. Croc. caſtor. ana cape ad menſes.

Contra ſputum ſanguinis & mictum cruentum: ℞. ſuccin. præpar. ʒj terr. ſigill. lap. hæmatit. ana ℈ß conſerv. heder. terreſtr. ℥ß L. O. gr. j ſyr. de ſymph. q. ſ. pro bolo.

Externe parantur etiam linimenta pro partibus convulſis egregia, ut & balſama antapoplectica, quæ naribus induntur. Sic ℞. ☍ expr. bacc. laur. ℥ß. ☍ ſuccin. lavend. ana ℈j camphor. ʒß. M. f. liniment.

℞. ☍ junip. ſuccin. ana ʒj. ☍ caryoph. gt. v. camphor. q. ſ. ſcil. gr. xxx vel xL f. balſamum.

Ex diutina coctione cum oleo lini fit vernix pro incruſtandis ſcriniis.

GUMMI ELEMI.

Eſt ſubſtantia reſinoſa pinguis ex flavo albicans, ſplendida, ad viridiuſculum inclinans, in maſſam co-

acta, dum recens mollecula, dum inveterata durior, collecta ex arbore indica prunifera, ſaporis reſinoſi & pinguis, odoris fragrantis, cum ſuavitate graviuſculi.

Hujus uſus externus frequentiſſimus eſt præcipue in confortandis nervis & roborando cerebro, quidam interne commendant pro diuretico.

In uſum domeſticum parant Indi impinguando ſtoreas cannabinas vel paleas, quibus deinde utuntur, pro elychniis, diutius enim alit flammam.

℞. Elem ʒß. detur ex vitello ovi ad gonorrhœam ſimplicem, ex debilitate nervorum à cauſa paralytica.

℞. Elem. q. ſ. ſolv. in ꝏ deſt. elem. vel ſucccin. f. ung. contra dolores artuum à frigore ortos.

℞. Elem. ℥j. ꝏ d. caryoph. ʒß. laurin. ℥j f. linimentum quo inungatur abdomen.

℞. Tacamahac. Caran. Elem. ana ℥ij ceræ ℥j picis ℥ß. ꝏ deſt. elem. q. ſ. ad diſſolutionem omnium, add. ꝏ mac. ʒß. ꝏ caryoph. ℈j f. emplaſtrum ad dolores ſcapularum, dorſi, in catarrhis, aliiſque affectibus à frigida cauſa ortis, applica etiam partibus dolentibus in colica.

CARANNA & TACAMAHACA.

Caranna eſt ſubſtantia reſinoſa, concreta, laxa, dum recens tenax, picis inſtar ductilis, ſine tamen agglutinatione vel colliquatione, in veterata dura ac friabilis, foris ex cinereo nigricans, picei coloris, collecta ex arbore Mexicana Caragna, ſaporis viſcoſi amaricantis, odoris dum accenditur fragrantis.

Tacamahaca eſt ſubſtantia reſinoſa, arida & retorrida, ex cinereo & rufo nigricans, modo figura minori, modo in glebas coacta, modo foliis tubuli in modum

dum convolutis incluſa, ex arbore Mexicana Tacamahaca collecta, ſaporis aromatici reſinoſi, odoris fragrantiſſimi.

Utrumque ob ſuaviſſimum odorem nervinum eſt & cephalicum potentius quam elemi. In ſpecie medetur externe podagræ, arthritidi, iſchiatico dolori, capitis item & dentium catarrhis. Egregie ſtimulatur ol. Nuc. moſch. expr.

℞. Carann. Tacamah. ana q. v. diſſolv. in balſ. Peruv. ſ. q. f. emplaſtr. temporibus applicandum ad prædictos affectus, etiam in debilitate ventriculi & doloribus poſt partum.

℞. Ol. deſt. caran. tacamahac. ana. q. v. ſuccin. ℥j f. linim. ad affectus prædictos.

℞. Gum. Carann. ſem. naſturt. ana q. ſ. M. c. albumine ovorum, & fac Ceratum, impone herniæ inteſtinorum.

Interne etiam uſurpari poteſt in affectibus pectoris, licet apud Practicos non præſcribatur. Sic ℞. Gum. Carann. Tacamah. ana ʒj affund. ol. thereb. ad eminent. iij digitorum f. eſſentia à gt. ix. ad xx. ſumenda in ▽ Cinamomi vel ſyrupo pectorali.

Oleum deſtillatum acre & rubicundiſſimum, ſpecificum uterinum eſt, confortatque egregie, odoriferum etiam ſummum eſt.

BENZOE ſeu ASA DULCIS.

Eſt ſubſtantia reſinoſa arida & indurata ex variis ſplendentibus micis modo flaveſcentibus, modo albicantibus in maſſam coacta, collecta ex arbore indica Benzoifera, ſaporis pinguis cum quadam dulcedine, odoris ſuavis.

Reſpon-

Reſpondet ſtyraci, & ob ſui mollitiem & blandam dulcedinem Pulmonibus dicatur, externe fiunt linimenta, emplaſtra, balſama, ſuffitus. Proſtant flores, tinctura ſ. eſſentia.

℞. Benz. ℈j Maſtich. thur. ana ʒj anim. ʒij M. f. ſuffitus in catarrhis & coryza.

℞. Flor. benz. ℈j. flor. 🜍is ℈ij balſ. Peruv. q. ſ. f. pil. contra tuſſim ab hydrope pectoris, quibus adde ol. dauci vel fœnic. ana gt. ij vel iij. Flores benz. ſoluti in ol. Thereb. balſamum præbent ſumme aſthmaticum, in ♒u vini tincturam rubicundam ſumme thoracicam ex ſyrupo pectorali.

PIX.

Eſt ſubſtantia reſinoſa nigricans, ſplendeſcens, ignis calore in furnis, vel deſtillatione per deſcenſum ex foliis & arboribus reſinoſis & piceis, ut pini, abietis &c. collecta & prolecta, primo liquida, hinc igne inſpiſſata, ſaporis reſinoſi, odoris gravis. Conſtat ⊖e volatili 🜔ſo incluſo reſinæ & craſſis terreis.

Robuſtiſſimis etiam in dyſenteria aliquando datum una cum cera, in forma pilularum. Externe virtute diſcutiendi, temperandi, mulcendi cum reſinis aliis convenit, ingreditur emplaſtra & cerata, ob vim conglutinantem, it. ad tineas capitis & alopecias.

℞. Pic ℥iij reſin. pin. ℥j virid. æris. ⊕li Romani ana ℥ß. 🜍r. viv. ʒj coq. c. ✝to f. emplaſtrum contra tineam applicandum, & biduum relinquendum, poſtea removendum, ut hujus beneficio radices pilorum ſimul evellantur.

STYRAX CALAMITA.

Eſt ſubſtantia reſinoſa, pinguis, mollecula, ex flavo rufeſcens, in grumos vel micas variæ magnitudinis collecta, ex trunco arboris ſtyracis foliis mali cotonei in inſulis Maſſilienſibus & locis calidioribus, ſaporis ſubacris lenti aromatici, odoris ſuavis. Habetur etiam ſtyrax liquida, quæ ſponte fluit, eſtque ſubſtantia liquida, reſinoſa, terebinthina craſſior & ſpiſſior, ex flavo ferruginea, collecta ex corticibus arboris ſtyracis, folio mali cotonei, ſaporis acris aromatici oleoſi, odoris fragrantis gravis.

Convenit cum tacamahaca ſed cum odoris ſit fragrantioris, conducit in corroborandis nervis, cerebro, ſpiritibus ut & utero, convenit & cum benzoë ſed abeſt dulcedo, interne datur velut maſtiche, externe recipitur inter ſuffimigia.

℞. Styrac. calam. benz. ana ℈j Maſtich.| oliban. ana ℈ß L. O. gr. j. ſucc. liquir. ſol. in elix. propr. q. ſ. f. pil. in obſtructionibus cerebri cum doloribus, in coryza.

Sublimantur etiam flores, qui exterius in doloribus pudendorum ſpecifici ſunt, vel ſoli cum modico ⊕li applicati.

*
* *

RESINÆ LIQUIDÆ SALE VOLATILI OLEOSO CONSTANTES.

TEREBINTHINA.

Eſt ſubſtantia reſinoſa, tranſparens mollecula, reſina laricea craſſior, coloris flaveſcentis ad cœruleum

vergentis, collecta ex trunco arboris Terebinthi ad imam medullam ſauciato, ſaporis acris amaricantis reſinoſi, odoris fragrantis & gravis, & hæc quidem eſt Cypriæ deſcriptio. Vulgaris colligitur in Germania, Norvvegia ex truncis pini & abietis junioribus ſauciatis, quæ dum lavatur aqua, donec limpida tota iterum poſſit decantari depurata dicitur.

[Terebinthina vera ex Cypro vel Chio inſula allata nunc raro eſt in uſu apud nos, loco hujus uſurpatur Terebintina Veneta dicta, & eſt reſina liquida collecta ex arbore laricis, conſiſtentia mellis novi, coloris flaveſcentis, ſaporis acris & amaricantis.]

Reſpondet ſuperioribus reſinis & gummatibus ſiccis, aſt cum gaudeat ⊖e volatili copioſo balſamico oleo ſociato & temperato, vim ſuam per diureſin exerente, multivarii uſus eſt: inprimis medetur tuſſi, aſthmati, pleuritidi, nephritidi, ſtranguriæ, gonorrhææ, fluori albo, & ſanandis vulneribus tam internis quam externis. Alvum facit apertam in majori doſi.

Externe in nervorum vitiis & tumoribus diſcutiendis nulli inferior eſt, additur quoque clyſteribus tam vulnerariis quam anticolicis & antinephriticis, veſicæ utero & inteſtinis injiciendis.

In genere datur à ʒß ad ʒj datur lota cum vitello ovi, vel in Emulſione.

℞. Tereb. ʒß. cape ex vitell. ovi add. tantillum ſacchari contra tuſſim, addito ſucco Citri contra pleuritidem & nephritidem, addito ſyrupo ſymphiti ad vulnera & ulcera interna, præcipue renum & veſicæ.

℞. Sem. papav. ʒij Melon. ℥ß. f. ex ▽ Portulac. Petroſel. ana ℥iij emulſ. add. 🝆 thereb. gt. vij. in nephritide & lumborum dolore.

℞. Tereb.

℞. Tereb. ʒij rhabarb. ʒj Merc. virid. gr. iij ſacch. ♄ni ℈ß cap. cum vitello ovi contra gonorrhæam virulentam, pro duabus doſibus.

℞. Rhabarb. toſt. Nuc. Moſch. toſt. borrac. probe calcinat. oſſ. ſep. terræ alum. dulc. ana ℈j Merc. virid. gr. iij tereb. q. ſ. f. pil. in gonorrhæa ſiſtenda, pro quatuor vicibus.

Poſt curatam maxime pleuritidem reſiduis tamen adhuc doloribus. ℞. Pic. naval. ℥iß. flor. ♀is ʒß ceræ ʒvj tereb ℥j 🝆 chamom. q. ſ. f. emplaſtr.

Fiunt etiam clyſteres in paſſionibus colicis non ſpernendi. ℞. Urinæ human ℔j thereb. venet. ℥j vitell. ovor q. ſ. f. clyſter injiciendus parari etiam hoc modo poteſt cum lacte bubulo.

BALSAMUM PERUVIANUM ET TOLUTANUM NIGRUM.

Balſamum Peruvianum eſt liquor oleoſus viſcoſus tenax, balſamicus nigricans ex arbore Hoitziloxilt Hernandez dicta, ramis cortice & ligno excoctus & ſaporis & odoris balſamo albo debilioris & ignobilioris. Fluit illud etiam partim ſponte, partim ex arbore ſauciata, eſt ſacchari inſtar coagulabile.

Spirituoſius eſt & ſubtilius terebinthina & in ſpecie confert in phthiſi. In vitello ovi ſolut. datur à gr. v. ad ʒß. exhibetur ut terebinthina, additurque pro ſpecifico clyſteribus in eroſione inteſtinorum.

℞. Balſ. Peruv. ℈ß. 🜕 Junip. ʒiij vitell. ovi q. ſ. ad ſtomachum confortandum.

℞. Balſ. Peruv. ʒij 🝆 ſuccin. ℈ß. f. liniment. ad epilepticas convulſiones.

In phthiſi conjungitur cum elix. p. p. ſ. acido ad ʒj cujus gt. vj ad x exhibentur, & decocto vel ſyrupo thoracico vel aqua hordei.

In majori quantitate ſudorem movet, ſic in morbis malignis ♀eis. ℞. reſ. lign. guajac. ʒß. ☍ Guajac. vel lign. rhod. gt. ij Bez. min. ℈ß. Balſ. Peruv. q. ſ. f. pil. ſudoriferæ nocturnæ.

BALSAMUM PERUVIANUM ALBUM.

Et liquor oleoſus, tenax, limpidus ex albo flaveſcens terebinthina conſiſtentia tenuior, ſtillans ex arbore Americana eadem, ſaporis acriuſculi aliquantum amari, odoris fragrantis & grati.

Eſt potentius nigro, ſpecifice commendatur in diarrhoea & gonorrhœa, ubi cum adſtringentibus formantur pilulæ.

Oleum deſtillatum ſtatim concreſcit in ſalem inſtar ſacchari, candidum velut camphora, quod nobile eſt in obſtructionibus uteri, tuſſi & aſthmate.

BALSAMUM DE COPAIVA.

Eſt liquor oleoſus, viſcoſus, tenax, terebinthinæ conſiſtentia, balſamicus limpidus, ſubflavus, ſtillans ex arboris Americanæ Copaiva Bontii trunco ad medullam uſque vulnerato, ſaporis acriuſculi amaricantis aromatici, odoris fragrantis. Nihil differt à Peruviano niſi quod in fluore albo & gonorrhœa ſit ſpecificum, datur ut Peruvianum à ℈ß ad ℈j.

℞. Rha-

℞. Rhabarb. ʒj balſ. copaiv. ʒiß. ☿ vir. gr. ij ſolv. in vitell. ovi, ſingulare eſt in fluore albo.

℞. Extr. lign. ſaſſafr. cui proprii olei aliquid admixtum eſt ℈ß Balſ. Copaiv. gr. xv. vitell. ov. q. ſ. cape pro gonorrhœa & fluore albo.

℞. Succin. præpar. ℈j. balſ. Copaiv. q. ſ. f. pil. ſtipticæ ex ▽ Hormini.

AMBRA LIQUIDA

Eſt liquor reſinoſus, oleoſus, pinguis craſſiuſculus, terebinthina conſiſtentia tenuior & dilutior, coloris ex flavo rufeſcentis, ſtillans ex arbore Mexicana, ſaporis acris aromatici, odoris ad ſtyracem accedentis. Styraci reſpondet, & potius externo quam interno uſui dicata eſt, adhibetur ut terebinthina.

℞. Liquid. ambr. ʒij Moſch, gr. j f. linimentum ventriculo applicandum.

℞. Ambr. liquid. ℈j tacamahac. ʒj f. emplaſtrum temporibus ad capitis dolorem applicandum.

OPOBALSAMUM.

Eſt ſubſtantia liquida balſamica oleoſa ac craſſiuſcula limpida, albicans, collecta ex arbuſcula Syriaca & Ægyptiaca, Balſamum dicta, ſaporis acris aromatici, odoris fragrantis, fluit hoc ſponte ex arbuſcula. Fructus dicitur carpobalſamum, lignum vero xylobalſamum.

Recentiſſime collectum & deſtillatum oleum præbet fragrantiſſimum, ubi annoſius fit terebinthinaceum fere habet odorem.

Copaivæ reſpondet, ſed ſpirituoſius & ſubtilius eſt, in ſpecie laudatur in viſcerum debilitate; Alpinus contra ſterilitatem in Ægypto adhiberi refert.

Externe in nervorum vitiis applicatur.

℞. Opobalſ. ℈j ſpic. nard. q ſ. f. bol. vel diſſolv. in vitell. ovi & cape in debili ventriculo, add. rad. helenii pro appetitu excitando.

℞. Opobalſ. ℈j ſuccin. præpar. q. ſ. f. bolus vel cum vitello ovi dilue & cape pro fluore albo & gonorrhœa & Alpino teſte contra ſterilitatem.

LADANUM.

Eſt ſubſtantia partim reſinoſa, partim gummoſa pinguis, indurata ex gryſeo & fuſco nigricans, multis arenulis permixta, collecta ex Ladani Cretici, vel cyſti Ladaniferæ foliorum ſuperficie, & in ſpiras contorta, ſaporis amaricantis, odoris, dum accenditur fragrantis & grati: nec in aqueis nec oleoſis, ſed tantum ſpirituoſiſſimis ſolvitur, principia activa ſunt ⊖ia volatilia oleoſa cum multis partibus terreis, propter quas conſtringit, fibras confortat, cephalicis nervinis maritatur. Datur ſolut. in 🜄u vini à ℈j ad ʒß.

Ingreditur etiam candelas odoratas & balſama apoplectica, tum ratione virtutis, tum ratione nigredinis quia tingit, ſolvitur nempe ladanum prius in 🜄u vini ſolutum inſpiſſatum, additur 🝆 menth. caryoph. &c.

STYRAX LIQUIDA.

Ambræ liquidæ ſuccedanea eſt, & frequentius externe quam interne uſurpatur. v. ſtyr. calam.

RESINA

***

RESINA SALE TAM VOLATILI QUAM FIXO CUM INTENSISSIMA AMARITIE CONSTANS.

MYRRHA.

Specificam vim obtinet reſiſtendi putredini, reſerandi obſtructiones uteri, lumbricos expellendi, variolas & morbillos, & curandi morbos malignos quoſcunque.

Eſt reſina arida, dura, ex flavo ferruginea & nitens, unguibus ſubinde pallidis permixta, in glebas compacta. Collecta ex arbore ſpinoſa Bola dicta, cui aliquando adnatæ inveniuntur glebæ magnæ, ſaporis ſubacris amari & aromatici, odoris fragrantis. Myrrha Stacte eſt, quæ ſua ſponte effluit.

Datur à ℈ß ad gr. xv. in pilulis convenientius, quam in cæteris formulis, datur tantum in parva doſi ob inſignem amaritiem; in majori doſi alvum laxat, ſimulque vim adſtrictoriam habet in receſſu. Famoſiſſimæ ſunt illæ pilulæ de tribus dictæ, ſeu peſtilentiales Ruffi, quæ elixir proprietatis ſiccum dici merentur:

℞. Myrrh. gr. xv. borrac. ʒß cinam. ℈j. f. pulv. excipiendus vel conſerv. menth. vel balſamo Peruviano pro formandis pilulis, ☉i foliis obducendis ad menſium remoram.

℞. Myrrh. C. C. præpar. ♉ diaph. ana ℈ß f. pulv. pro ij doſibus. Vel cum theriaca in forma pil. exhibendus.

℞. Tinct.

℞. Tinct. Myrrh. eſſ. Caſtor. eſſ. aſ. fœtid. ana ℈ß cape ex ▽ flor. ſambuc. ad promovendam maturationem variolarum & morbillorum.

℞. Sabin. Mß Myrrh. ʒij lign. aſphalt. troch. alhand. ana ʒj f. ſuffitus ad remoram menſium.

In halitbus uteri putridis. ℞. Myrrh. gr. iiij croc. orient. gr. iij. rad. Zedoar. ℈ß. conſ. anthos q. ſ. f. bolus.

℞. Aqu. theriacal. ℥j. ♎ theriac. ʒij camph. gr. v. Myrrh. ℈j in malignis.

Convenit etiam me experto in tertiana & quartana, quas aliquando feliciter fugavi h. m.

℞. Myrrh. ℈ß. ⊖ abſynth. ℈j extr. gentian. vel abſynth. cent. min. ana ʒß f. pil. paulo ante paroxyſmum, etiam ſi placuerit diebus vacuis aſſumendæ.

Singularis eſt etiam in defectu bilis ab obſtructo poro vel veſicula à calculo. ℞. Rhabarb. ʒij myrrh. ℈ß rad. helen. ʒß f. cum pauxillo ſacchari pulvis, vel cum ♎u vini aucta helenii doſi tinctura, guttatim ex vino generoſo vel juſculo exhibenda.

Externe vel per ſe vel cum Euphorbio inſpergitur oſſibus denudatis carioſis, vel cum ♎u vini extrahitur tinctura, quæ in ſphacelo & gangræna ſpecifica: egregio ſæpe cum ſucceſſu ingreditur etiam linimenta contra gingivas putridas, ſanguineas ſcorbuticas, ſic ℞. Tinct. Lacc. ʒj Myrrh. ʒß alum. ʒj mell. anthoſ. ſyr. gran. cherm. q. ſ.

RESINÆ

***

RESINÆ GUMMOSÆ ET SUCCI PURGANTES FORTIORES.

EUPHORBIUM.

Eſt ſubſtantia gummoſa & reſinoſa concreta in guttas vel lachrymas ex pallido flaveſcentes, ſplendidas, modo rotundas, modo oblongas, ramoſas & cavernoſas, collecta ex Tithymalo Mauritanico ſpinoſo ſarmentoſo, ſemine tricocco, ſaporis acerrimi exurentis nauſeoſi, odoris nullius.

Principia activa ſunt ⊖ia volatilia cum aliquibus fixis, quæ maxime maſſæ noſtræ ſanguineæ adverſantur, cum illa mixta excitant efferveſcentiam purgatricem. ⊖ia fixa ſunt in decoctis ſatis acribus.

Fortiſſime purgat, hinc ſpecificam & pôtentem habet vim educendi hydropicorum aquas tam per ſuperiora, quam per inferiora, prodeſt ſeroſis melancholicis, nocet bilioſis ſanguineis. Vix ſe patitur cicurari, quod quidam tentant ſolvendo in ⏜u ⊕li, aceto, & hinc iterum inſpiſſando, ſubſequente tamen nonnunquam ubi exhibetur inflammatione ventriculi & inteſtinorum excoriatione. Corrigitur amygdalis vel ſucco portulacæ.

Datur in ſubſtantia à gr. v. vij. ad x. extract. c. ⏜u vini à gr. iij in robuſt. ad v. vij. Nunquam in pulvere datur propter cauſin, ſed in pilulis, addito ſtimulo alcalico vel etiam purgante. Sic ℞. Euphorb. gr. v. extr. coloc. gr. iij. ⊖ ☿ri gr. j caryoph. arom. gr. v. f. c. balſ. peruv. vel elix. propr. P. pil. antihydropicæ.

℞. Euphorb. ℈ß extr. abſynth. ℈j mell. q. ſ. f. bolus ad quartanam cum hydrope curandam.

℞. M. P. fœtid. ℈j Euphorb. gr. iij ſucc. limon. q. ſ. f. pil. in morbis chronicis, præcipue hydrope.

℞. Euphorb. ʒj rad. ir. flor. ℥j ari ℥j f. pulvis inſpergendus denudatis oſſibus & carioſis, addi poteſt ☍ deſt. caryophyllorum.

℞. Extr. euphorb. admiſce gr. iij cuilibet purganti pro ſtimulo, in hydrope.

Pro excitandis ſoporoſis. ℞. ▽ ſerpill. origan. lil. convalli. ana ℥ij Euphorb. ʒij f. decoct. quod naribus admoveatur.

Ad tumores ſcirrhoſos ſæpe cum fructu maximo D. Hermannus diſſolvit euphorbium in oleo, tribuſque vel quatuor diebus illos ſuſtulit.

ELATERIUM.

Eſt ſuccus ille viridis luridus & viſcoſus ex fructibus maturis cucumeris ſylveſtris aſinini C. B. P. vel ſua ſponte vel leni compreſſione emiſſus, collectus & inſpiſſatus, qui veterum eſt elaterium. Vel ex tota planta expreſſus coctus inſpiſſatus coloris fuſci, inſtar extracti, ſaporis amaricantis, nauſeoſi & vinoſi.

Datur in affectibus deploratis chronicis, hydrope, cachexia, leucophlegmatia ad hydropem vergente, ſurſum enim & deorſum humores viſcidos & aquoſos expurgat.

Elaterii veterum doſ. eſt à gr. iij ad v. recentiorum ad ℈j, quod ſi debite correctum ſit ſucco limonum vel ♎u ⊕li etiam delicatulis ad ℈j poteſt exhiberi.

℞. Elat,

℞. Elat. vulg. gr. xv. reſin. Jal. gr. iij extr. troch. alhand. gr. j. f. pil. antihydropicæ.

℞. Elater. veter. gr. iij conſ. flor. perſic. ʒj f. bolus antihydropicus.

GUTTA GAMBA.

Eſt ſubſtantia gummoſa & reſinoſa indurata, flaveſcens ex arbore pomifera Chinenſi, Indis Coddampulli dicta & in horto Malabarico [Tom. I. p. m. 41. ubi etiam D. Hermanni noſtri literæ Columbæ datæ de hac arbore producuntur] deſcripta. Ex ſaucio enim trunco manat humor flaveſcens, qui collectus purgatur à fœcibus ſolutione in 🜄 pluvia & inſpiſſatione, ſed leniſſimo 🜂e alias nigreſcit, in maſſam uniformem flaveſcentem, ſaporis primum reſinoſi, hinc acris cauſtici. Transfertur potiſſimum ad nos ex China, ubi affabre norunt tractare.

Virtutem ſuam purgantem paucæ reſinæ, ſed largiori gummati habet incluſam, hydropicis ſpecificam, quarum aquas tam per ſuperiora quam inferiora educit. Datur etiam in ſcorbuticis, potiſſimum in ſubſtantia, rarius in potionibus, fiunt enim acres valde & inflammant fauces.

Datur à ℈ß ad ℈j poteſt & delicatulis dari, licet tormina excitet, hoc enim omnibus fere accidit purgantibus.

℞. Gutt. gamb. ℈ß. extr. troch. alh. gr. ij ſyr. roſ. ʒv. 🜄 Cinam. fœnic. ana ℥j M. f. mixt. purgans contra hydropem.

℞. Reſin. jalap. gr. v. gutt. gamb. gr. v. vel vij diagr. gr. iij 🜔 ♀ri gr. v. ☍ d. caryoph. gt. ij ſyr. rhab. q. ſ.

[Optime exhibetur cum anatica quantitate ☿ri ⊕lati & ſic raro vomitum excitat.]

SCAMMONIUM.

Eſt ſubſtantia reſinoſa nigricans, cum frangitur fulva ſplendens, collecta ex convolvuli ſpecie ſcammonia Syriaca C. B. P. dicta, ubi vel in excavatam radicem, quæ aliquando ad capitis magnitudinem accedit, deſtillat ſuccus, vel ex radice in taleolas ſciſſa exprimitur & inſpiſſatur, ſaporis acris nauſeoſi, odoris viroſi.

Etiamſi inter purgantia cholagoga ſit relatum, tamen uſus comprobavit eſſe catholicum, polychreſtum, & in omnibus morbis conducere, ubi purgatione opus eſt. Hujus vis conſiſtit in reſina, cui acredo ⊖ina indiſſolubili nexu eſt involuta.

Datur in ſubſtantia & extracto à ℈ß ad gr. xv. robuſtiſſ. ad ℈j reſin. à gr. vij ad xv. in vitell. ovi.

℞. Scammon. crud. gr. vij. extr. troch. alhand. gr. iij. ⚯ Cinam. gt. j. ſyr. cinam. ʒj ▽ borrag. fumar. ana ℥j f. hauſtus purgans.

℞. Reſin. Jalap. gr. vij ſcammon. gr. v. ☿ dulc. gr. iij ⚯ cinam. gt. j. balſ. peruv. q. ſ. f. pil.

℞. Scam. ♎at. ℈ß ☿ dulc. ℈j mucag. tragac. q. ſ. f. pilulæ Riverio Calomelanos dictæ. v. plura in Jalappa.

ALOE.

Hepatica eſt Aloës Aſiaticæ ſuccus gummoſus & reſinoſus, foris fuſcus intus flaveſcens, hepatis inſtar, expreſſus ex foliis Aloës, imo & plantis tranſverſim radicem verſus inciſis, extillanſque in patinas ampliores, in ſole concretus & condenſatus, ſaporis aris amariſſimi,

riſſimi, nauſeoſi pinguis oleoſi & adſtringentis, odoris fragrantis ad myrrham accedentis. Colligitur Hepatica in China non in Socotra.

Succotrina eſt Aloës Aſiaticæ ſuccus gummoſus & reſinoſus ex fuſco nigricans, dum fractus in lamellas ſplendens & pellucidus ex foliis imo & planta tranſverſim radicem verſus diſſectis, ſua ſponte in vere ubi ſucci nutrititii luxuriant in ſuppoſitas patinas amplas deſtillans, vel etiam expreſſus, in quibus à Sole condenſatur & exſiccatur; hinc ulterius inſpiſſatur & quo diligentius hæc fiunt, eo pellucidior & nobilior fit, ſaporis & odoris cum Hepatica ejuſdem. Colligitur in inſula Succotra, quod in fundo ab illa fœculentum remanet, id Aloën caballinam conſtituit, differuntque utræque plantæ, ex quibus colligitur Hepatica & Succotrina, nonniſi loco natali.

Caballina eſt Aloës Aſiaticæ ſuccus nigricans gummoſus, prioribus ponderoſior craſſior ac impurior expreſſus ex foliis Aloës diſſectis, in Sole concretus & ſaporis ſubacris amariſſimi nauſeoſi, odoris gravioris quaſi picei & myrrhati, conficitur etiam in omnibus tractibus Indiæ Orientalis maritimis.

Hepatica quæ mihi optima videtur vim purgantem habet in ſubſtantia gummoſa, cui quippiam mucaginis reſinoſæ in qua vis ſtyptica & balſamica juncta eſt, purgat blande & in ſpecie humores bilioſos. Specificum habetur in affectibus capitis, in præcavenda putredine, in omnibus morbis à bile oriundis.

Ob ſingularem acredinem excitat hæmorrhoidum fluxum, venaſque hæmorrhoidales aperit, quod tamen impeditur Aloë roſata & violata facta. Externe vulnera & ulcera mundificat, toſta & virtute pur-

 gante

gante orbata vim adſtringentem habet, & propterea in fluxibus quibuſvis ſanguinis ſpecificum eſt.

Tardior vis corrigitur diagrydio, 🝆ro ⊕lato &c. correctorio aromatico vix opus habet.

℞. Aloës ʒj myrrh. ʒß. Croc. or. gr. xv. oxym. q. ſ. f. pil. contra malignos morbos præcavendos.

℞. M. P. aloëtic. ℈j ambr. gryſ. gr. iij ⊖ vol. ſuccin. gr. v. °oo d. ſerpill. gt. iij balſ. peruv. q. ſ. f. pil. catarrhales & cephalicæ.

℞. Al. toſt. rhabarb. toſt. borrac. calc. ana ℈ß. f. pulv. exhibendus cum ▽ plantag. ad fluxum quemlibet, & præcipue ſanguinis, hæmoptyſin.

Externe fiunt emplaſtra ſic: ℞. Myrrh. ʒß Aloës ʒij fell. taur. q. ſ. f. emplaſtrum, quod applicatum umbilico vermes necat.

Caballinæ uſus Medicus non eſt, Indi utuntur pro incruſtandis navibus picis loco, norunt nunc eam expurgare à ſordibus, ita ut à Succotrina etiam experientiſſimus vix contradiſtinguere valeat.

MANNA.

Sunt grana ex albo flaveſcentia, pinguia, figuræ variæ, quaſi ex Salina quadam ſubſtantia concreta, collecta ex arboribus orni & fraxini in Calabria, Sicilia, agris Romanis, ubi ſæpe coluntur, ſaporis ſubacris, ſubdulcis, pinguis & nauſeoſi, odoris nullius vel melliti.

Meminere Theologi Mannæ cœleſtis, Philoſophi aëreæ, Medici frumentaceæ quæ in Polonia ex ſeminibus alicujus graminis colligitur [etiam ſuper folia borraginis reperitur. v. Hœfer. Herc. Med. p.m. 385.] item factitiæ à formicis alatis candicantibus in India

Orien-

Orientali [vid. Joh. Ott. Helbig. C. N. C. Dec. 1. an. 9. & 10. obſ. 194. p. 459. & Herm. Nic. Grimm. C. N. C. Dec. 2. ann. 1. & 2. obſ. 151. p. 369.] ſed hæc fori noſtri jam non ſunt. Loquimur jam de illa ſpecie arborum in Calabria &c. ubi diebus canicularibus potiſſimum ſucco ſeu nutritio ſeu alio luxuriantes, ab hujus copia fiunt tubercula, quibus hiſcit cortex, ex quibus deinde profluit ſuccus viſcidus ſubdulcis, qui pomeridianis horis à Sole nonnihil condenſatur, colligitur, & in clibano ulterius exiccatur. Non omnes indiſcriminatim illæ arbores fundunt hanc mannam, ſed quæ talis eſt, per annos quadraginta & plures mannam depromit.

Calabrina, quæ optima, fovet in ſubſtantia dulci gummoſa virtutem laxantem, & in ſpecie evacuant bilem, ſed cave exhibeas iis qui bilem habeat viſcoſam & ad icterum vergentem, ut & Hypochondriacis, Melancholicis & Scorbuticis, cum enim ſalia tam blanda ſint, facile obtunduntur, & in corpore remanentia putreſcunt. Vis ejus torminoſa corrigitur cinamomo, tarda vero exaltatur ⊖ e ♀ri, ſcammon. &c.

℞. Mann. ℥iß cape ex ꝏ amyd. d. rec. expr. ad ℥iiß ex juſculo ad laxandam alvum, præcipue in colica, add. ſem. fœnic. vel cinam. pro correctorio.

℞. Mann. ℥ij Diagryd. gr. iij Crem. ♀ri. ℈j M. cape ex vino.

Datur infantibus per ſe ad ℥j. vel ʒj cum gr. ii vel iij diagrydii.

SARCOCOLLA.

Eſt ſubſtantia gummoſa in grumos vel ſabulum concreta ex albo rufeſcens, quaſique mannæ colore, ſpon-

ſpongioſa, lucentibus ſubinde micis permiſta, levi concuſſu in pollinem abiens, collecta ex tithymalo Perſico ſpinoſo ſarcocollifero, ſaporis amaricantis cum obſcura quadam dulcedine, odoris dum incenditur ingrati.

Convenit cum gummi ammoniaco, galbano, niſi quod major inſit vis catharctica, quod acredo major teſtatur, ob dulcedinem ſuam ſpecifice convenit in vitiis pulmonum & thoracis. Externe convenit in conſolidandis vulneribus.

℞. Sarcocoll. ʒß extr. troch. alh. gr. iij Zinz. gr. vij ſyr. roſ. q. ſ. f. pil. purgantes ſerum & pituitam.

℞. Myrrh. Aloë ſarcocoll. ana q. v. f. pulv. pro conſolidandis vulneribus.

OPOPANAX.

Eſt ſubſtantia reſinoſa gummoſa ex rufo & albo flaveſcens mollecula, quaſi ex grumis vel ſabulo craſſiori in maſſam redacta, collecta ex radice Panacis Sphondylii C. B. P. ſaporis acris amaricantis ingrati, aliquatenus nauſeoſi, odoris fragrantis gravis.

Ad galbanum accedit, & ob intenſam acredinem habet vim potentiorem alvum deponendi, ſerum & pituitam educendi. Externe laudatur in herniis, præcipue carnoſis diſcutiendis & ſuppurandis, ut & in ſcirrhis, nodis & gangliis reſolvendis.

℞. Opopan. ʒß. Caſtor. ℈j. borrac. gr. v. troch. alhand. gr. vj. ſyr. ſtœchad. q. ſ. f. pil. uterinæ in menſium obſtructione & lochiorum ſuppreſſione.

℞. Opopan. ſagap. galban. ammon. ana ʒij. 🜍 lumbr. vel philoſoph. q. ſ. f. emplaſtr. ad hernias carnoſas ſuppurandas.

SUCCI

SUCCI CONCRETI MUCAGINOSI DULCES.

SACCHARUM.

Eſt ſubſtantia gummoſa, facie ⊖is contuſi, excocta ex expreſſo ſucco arundinis ſacchariferæ. Ita fit in India Orientali & Occidentali; agri probe aratro perfoſſi conſeruntur articulis Cannæ ſacchariferæ, exſurgunt hinc arundines digitos quatuor circiter craſſæ, quæ dum ad altitudinem hominis aſcendunt, revelluntur à radice, in conſtitutas vehuntur molendinas, ibidemque ſuccus omnis ex iis exprimitur, qui niſi intra 24. horas coquatur, aceſcit, nec ulla arte reduci poteſt. Coquitur autem primo lixivio forti, ut viſcoſitas hæc melius incidatur ad ſpiſſitudinem mellis, hinc iteratur coctio ſine calce, ſi nimium ebullit, fervor hic ſupprimitur affuſo oleo vel tantillo butyri, vel continua debet fieri agitatio; coctus infunditur in vaſa conifera & exponitur Soli ut indureſcat, talis defertur ad nos, ubi varias adhuc ſubit ſolutiones & cryſtalliſationes.

Conſtat mucagine temperata dulci cum blanda acredine, quibus nominibus tam alimentum, quam medicamentum eſt, temperans & obtundens quævis acida, reſolvit quævis rigida & aſpera. Indi coquunt in aqua cum pauxillo farinæ ad nutritionem.

Promovere urinam experientia docuit, ſpecificum eſt in tuſſi, raucedine, aſperitate aliiſque pulmonum vitiis. Melancholicis, hypochondriacis, hyſtericis, febricantibus &c. ob facilem ſui aceſcentiam

ſacchari largior uſus non convenit. Externe vulnerarium & ophthalmicum eſt.

℞. Sacch. q. v. coq. c. ▽ q. ſ. in form. ſyr. vel julepi add. ſacch. ♄ni tantillum, cape contra inflammationem faucium cum raucedine.

℞. Spir. vin. vulgar. q. v. M. c. ſacchar. & accende ♌um, deflagret, cape hanc maſſam cochleatim pro raucedine & catarrhis.

Fiunt etiam clyſteres cum ſaccharo ſic pro infante, ℞. juris ping. ℥vj ſacchar. ℥j ſtimuletur ⊖e gemmæ pro alva laxanda.

SUCCUS GLYCYRRHIZÆ.

Eſt glycyrrhizæ ſiliquoſæ vel Germanicæ C. B. P. radicis ſuccus expreſſus, ignis calore inſpiſſatus & in glebas redactus, coloris nigricantis, ſaporis ſubacris & dulcis, odoris debilis. Convenit cum propria radice è qua conficitur.

℞. Succ. glycyrrh. ſolv. in ▽ Puleg. f. hauſtus contra tuſſim.

℞. Succ. glycyrrh. ʒj ſacch. cand. in ▽ puleg. ſolut. f. morſuli pectorales.

℞. Aur. foliat. volum. v. vel vj ſucc. glycyrrh. in ▽ fœnic. ſolut. f. pilulæ in fluxu nimio ſalivæ à ☿libus, quibus ſi alvus foret adſtricta add. reſin. jalap. gr. v. ⊖ ♀ri gr. ij, ſic enim ubi erunt imprægnatæ globulis ☿ij per alvum ſecedent è corpore.

MEL.

Eſt ſubſtantia liquida craſſiuſcula ex albo flaveſcens, nata in variis vegetabilium floribus & ab apibus in iiſdem collecta.

Cum

Cum ſaccharo convenit, aſt temperatius, ⏜uoſius & magis anodynum eſt.

℞. Mell. q.v. camphor. par. applica vulneribus mundificandis, vel add. ⏜ vin. ▽ reg. Hung. vel balſ. Peruv. pro eodem uſu add. virid. æris ſi ulcus ſit malignum, add. ſacch. ♄ni ſi ſit cum inflammatione.

℞. Mell. deſpum. ℥iij paſſul. major. ℥iiij liquirit ℥j ſacch. cand. ℥v vin. Hiſpan. ℥vj bulliant invicem f. conditum, add. ad gratiam ⏜ 🜍is per campan. in aſthmate & tuſſi expertiſſimum.

℞. Mell. ℥ß. ▽ Cinam. ℥j. ⏜ ⊖✱ci ʒj f. mixt. add. etiam flor. benz. ℈j.

* * *

GUMMOSA AQUOSA CRASSA.

GUMMI ARABICUM.

Eſt ſubſtantia in grumos variæ magnitudinis & formæ condenſata, foris ſuperficie granulata, pallide flaveſcens vel omnino albicans, intus pellucida vitri inſtar, collecta ex arbore Ægyptiaca Acacia dicta, ſaporis aquei viſcoſi lenti, odoris nullius.

Gummi Arabicum optimæ notæ eſt vermiculatum, purius enim illud & candidius.

Ratione ▽eæ ſubſtantiæ in quam totum reſolvitur, temperat, incraſſat lympham acriorem & tenuiorem, & proinde in raucedine orta à catarrhis tenuioribus mulcet: in ſtranguria & acri ſanguine ſpecificum eſt, recentiores nonnulli commendant ad ſtranguriam & urinæ ardorem h.m. ℞. Gumm. Arab. ʒß. Nuc. Moſch. ℈j f. pulv.

℞. Gum. Arab. ʒß ſacch. cand. ʒj. ▽ ſperm. ran. ℥ij f. gargariſma ad faucium aſperitatem; adde ſacchar. ♄ni ſi inflammatio adſit.

℞. Gumm. tragac. Arabic. ana ℈j ſucc. glycyrrh. ʒß ſacch. cand. ℥j ▽ roſar. q. ſ. f. l. a. trochiſci vel rotulæ ad raucedinem & faucium aſperitatem.

TRAGACANTHA.

Eſt ſubſtantia gummoſa, lucida, alba, tenax, levis gracilis, collecta ex radice tragacanthæ, quæ in agris Maſſilienſibus copioſe alitur, ſaporis ſubdulcis & mucilaginoſi.

Craſſiore mucagine conſtat, quam Gummi Arabicum, cum cujus virtute cæteroquin convenit.

℞. Succ. glycyrrh. ʒj. terr. Japon. ʒj ſacch. cand. ʒiij opij gr. v. mucilag. tragac. q. v. f. trochiſci ſublinguales ad excipiendos humores catarrhales.

*
* *

RESINÆ OLEO CRASSO CONSTANTES.

BITUMEN JUDAICUM.

Naturam & eſſentiam picis imitatur.

℞. Bitum. ʒj ladan. ℈j. ☍ d. caryoph. gt. vj. ☍ ſuccin. q. ſ. f. balſam. apoplecticum.

COLOPHONIUM.

Eſt ſubſtantia reſinoſa, arida & indurata flaveſcens, ex reſina vel laricis vel abietis vel terebinthi excocta, ſaporis pinguis oleoſi & reſinoſi.

Lenit,

Lenit, temperat, emollit, ſimulque urinam provocat, ejuſque acredinem temperat, pro quo ſcopo ſolutum cum -♎-u vini exhibetur, uſus externus tamen frequentior eſt interno.

℞. Colophon. q.v. Spir. vin. rectific. q.ſ. f. tinctura, nonnullis ☉ potabile dicta, contra morbos chronicos ab obſtructionibus dependentes.]

℞. Colophon. q v. ꝏ lil. alb. q.ſ. f. liniment. ad mammas induratas, add. quoque emplaſtr. de ſperm. ceti.

CERA.

Eſt ſubſtantia oleoſa flaveſcens ex favis apum vel ſolis vel ignis calore extructa, ſaporis lenti & ꝏſi, odoris debilis.

Cera alba eſt ſubſtantia ceræ flavæ ſimillima, coloris albi, ex cera flava multiplici ▽æ infuſione & in Sole exiccatione in alborem converſa, ſaporis cum cera flava ejuſdem, odoris debilioris.

Particulis oleoſis mollioribus conſtat quam colophonium, hinc quoque potentius emollit, interne autem à nonnullis commendatur in fluxu alvi & ictero. Externus uſus frequentior eſt in linimentis & emplaſtris pro baſi.

℞. Ceræ, vitell. ovi ad duritiem coct. croc. or. gr. vij ſyr. abſynth. q. ſ. f. pil. ad icterum.

℞. ꝏ ceræ deſt. ad gt. v. exhibe cum ▽ petroſel. in paſſione nephritica & urinæ ſuppreſſione.

*
* *

RESINÆ ET SUCCI TERREIS CONSTANTES.

SANGUIS DRACONIS.

Eſt ſubſtantia gummoſa & reſinoſa, rubicunda, ex parvis quaſi grumis modo in panes ſeu offas cocta, modo folliculis foliaceis incluſa, collecta ex trunco arboris Mexicanæ Draco dictæ, non quod illam dracones circumvolitent, fabuloſum enim illud, ſaporis adſtringentis & viſcoſi. Arbores hæ naſcuntur etiam copioſe in inſulis Canarienſibus. Sunt arbores palmarum inſtar, nec fructus multum abludit ab his. Colligitur etiam ex aliis Indiæ arboribus. [conf. Dale Pharmacolog. ſupplement. p. m. 263. qui tradit tres dari ſpecies arborum, quæ ſanguinem Draconis præbent.]

Cum terreis virtute fluxus quoſvis ſiſtente convenit.

℞. Sang. dracon. ʒß. corall. r. ℈j. L. O. g. j. c. ▽ plantag. ℥ij. ▽ Cinam. ʒiij acet. ʒij ſyr. myrt. ℥ß. f. mixtura pro ſiſtendis menſibus.

℞. Sang. dracon. myrrh. ana ʒj gumm. helen. ʒiij thereb. venet. ʒvj. 🝆 hyperic. par. f. balſamum ad rupturas.

Datur in ſubſtantia à ʒß ad ʒj cum ſucc. acac. rob. rib. ſyr. cydon. &c.

℞. Sang. dracon. maſtich. myrrh. ana ʒß alum. ℈j Mell. roſat. q. ſ. f. linimentum in vacillatione dentium à cauſa ſcorbutica.

TERRA

TERRA JAPONICA
ſeu CATECHU.

Lycium veterum: Eſt ſubſtantia reſinoſa, gummoſa indurata, ex rufo nigreſcentis obſcuri & obſoleti coloris, expreſſa ex fructibus immaturis Arecæ & corticibus arboris Indicæ Catſchu, quæ eſt certa ſpecies Acaciæ & in maſſam calore △is redacta, ſaporis adſtringentis & auſteri, poſtmodum dulcis & grati, odoris vel nullius vel debilis.

Et fructibus & corticibus ſeorſim affunditur ▽ impraegnata calce, quam Indi parant ex conchyliis, utuntur loco ⊖ ☐ri, quo melius terrea virtus extrahatur, confundunt ſolutiones & inſpiſſant.

Conſtat adſtringentibus ſimul & abſtergentibus partibus, ob quas in alvi & ſanguinis fluxibus, ut & in catarrhis ſalſis tenuibus, in aſperitate, inflammationibus tollendis commendatur. In mundificandis ulceribus ſordidis eſt etiam ſpecificum.

Sumitur vel interne, vel detinetur in ore donec liqueſcat ad tollendas aſperitates.

℞. Terr. Japan, q. v. diſſolv. in ▽ add. aliquid ſucci liquirit. & leni △e denuo inſpiſſa, adde inſpiſſato aliquid ambræ & moſch. cum ⏜u roſar. ſolut. forma inde pilulas vel trochiſcos, vocatur Catechu præparatum.

℞. Terr. Japon. ʒß lap. hæm. præpar. ℈ij conſ. roſ. ⊕lat. ʒj L. O. g. j. f. bolus in fluxibus. [De terr. Japon. conf. G. W. Wedel. C. N. C. Dec. 1. ann. 2. obſ. 128.]

SUCCUS

SUCCUS ACACIÆ.

Ægytiacæ eſt ſiliquarum immaturarum acaciæ foliis ſcorpioides leguminoſæ C. B. P. ſuccus expreſſus coctus & induratus, ſaporis acerbi auſteri adſtringentis. Acaciæ in Ægypto duæ ſunt ſpecies, alia ſiliquis rotundis teretibus inarticulatis, aliæ articulatis ſiliquis iſthmo interjecto, ex hoc poſteriori colligitur copioſum Gummi Arabicum vermiculatum. Deſcripta habetur ab Alpino.

In utriuſque ſiliquis craſſioribus reperitur ſubſtantia quædam viſcida ſubdulcis aromatica, quam ego credo eſſe acaciam veterum, has enim vulgo adſcribunt ſaporis differentias. Datur in ſubſtantia à ʒß ad ʒj. Conſtat particulis acidis auſteris, cum aliquibus reſinoſis.

Germanicæ eſt fructuum pruni ſylveſtris immaturorum ſuccus expreſſus coctione induratus, nigricans, ſaporis acidi auſteri & adſtringentis.

Uterque notus eſt vi adſtringente & incraſſante, ob quam in ſiſtendis fluxibus laudatur. Germanicus differt tamen ab Ægyptiaco, vel nullam vel nonniſi parvam reſinam obtinet, modus utendi item.

℞. Succ. acac. ℈j L. O. g. j. ▽ plantag. ℥iiß. cinam. ʒij ſyr. acetoſ. ℥ß. f. hauſtus adſtringens in fluxibus.

℞. Acac. Germ. ʒj ſolv. in decocto cortic. gran. ʒij contra nimios menſes.

Veteres tribuere vim alexipharmacam ſed hoc intelligendum de medulla intra ſpinarum cavitates.

Pilulæ de ſucc. acac. cum ☿io viridi dantur in gonorrhœa.

HYPO-

HYPOCISTIDIS SUCCUS.

Eſt plantæ ſpuriæ quaſi ex Coſti radice enaſcentis & hypociſtis dictæ ſuccus expreſſus inſpiſſatus, niger, ſaporis auſteri acidi adſtringentis; naſcitur ut plurimum in Inſula Creta. Ex Ciſti foliis colligitur Labdanum.

Convenit cum ſuperioribus, vim etiam habet levem anodynam datur à ʒß ad ʒj.

℞. Succ. acac. hypociſt. ana ℈j. roob. rib. q.ſ. f. bolus vel cum ▽ roſar. portulac. ℥ij f. hauſtus in fluxibus alvi.

***

SUCCUS SALE SALSO CONSTANS NONNIHIL LAXANTE.

TARTARUS ALBUS ET RUBER.

Eſt ſubſtantia ſabuloſa, indurata, modo ex rufo cinerea, modo ex rufo candicans, intermixtis ſalinis quibuſdam micis, nata ex vini recentis vel muſti particulis terreſtribus fermentationis ope præcipitatis & coagulatis, ſaporis ſubacris acidi & nauſeoſi, odoris nullius.

Ob ſal ſalſum ex alcali multo & acido pauco conſtans virtutem tam incidentem diureticam & laxantem alvum habet, in ſpecie medetur obſtructionibus morborum chronicorum, præcipue ubi ſimul opus laxante.

Datur cremor à ℈j ad ʒj imo ʒiij, ſolvitur enim cito in ventriculo etiam debili. ♀rus ⊕latus à ℈j ad ʒß.

℞. ♀r. præpar. ʒß. ♀r. ⊕lat. ℈ß. M. f. pulv. in febribus ad laxandam alvum.

℞. ♀r. præpar. ʒj coq. cum juſculo pulli, cape ad alvum laxandam in obſtructionibus morborum chronicorum.

PARS SECUNDA.

CONTINENS REGNUM ANIMALE.

PARTES ANIMALES SALE VOLATILI OLEOSO CUM MULTIS TERREIS CONSTANTES.

CORNU CERVI.

HUjus ſubſtantia vel cruda vel præparata ⊖e alcalino & multa terra conſtans, vi incidente & abſorbente omnia acida & viſcida pollet, decoctum hujus temperat, eſt anodynum & blandum ſimul ſudoriferum, hinc confert in malignis, acutis, variolis, morbillis, diarrhœa, dyſenteria, epilepſia & pluribus aliis affectibus ab acido.

Datur crudum & præparatum à ʒß ad ʒj.

℞. C. C. præp. gr. xv. ⊙r. foliat. par. exhibe contra tormina infantum.

℞. C. C. præp. ʒj. ☍ Nuc. Moſch. expr. ℈j. L. ○. gr. ij conſ. fl. roſ. ʒiß. ſyr. acetoſ. citr. q. ſ. f. bolus in quolibet fluxu.

℞. C. C.

℞. C. C. præp. ʒß. Caſtor. gr. vij ⊖ vol. C. C. gr. v. M. f. pulv. exhibendus ex ▽ Card. B. & Cent. min. contra variolas & morbillos.

℞. Raſur. C. C. ℥ij coq. in ▽, cape decoct. ad temperandam ſanguinis acredinem, cui add. ſyr. roſar. vel violar. vel rub. id. ad ℔j. ℥j. f. Julepus.

℞. Raſur. C. C. ℥iiij rad. ſcorzon. ℥iij raſ. lign. ſaſſafr. ℥j f. decoct. ſudorif. in malignis pro potu ordinario.

♌us C. C. datur à gt. v. ad xxx in convenienti vehiculo.

℞. ♌us C. C. ſuccinat. ʒß. cape ex ▽ cinam.

℞. ▽ Cinam. ℥ß, ♌ C. C. ſuccinat. eſſent. caſtor. ana ʒß. L. O. gr. j. pro uteri vitiis.

Pulv. dentifricius: ℞. C. C. uſt. Corallor. oſ. ſepiæ, pumic. ana ℈j. ☍ lign. rhod. gt. ij M.

CRANIUM HUMANUM.

Reſpondet virtute Cornu Cervi, etiam ſpecificum habetur in epilepſia, datur in ſubſtantia à ℈j ad ʒj.

℞. Cran. human. ʒß cap. cum ▽ ♌uoſ. meliſſ. vel. fl. lil. ad Epilepſiam.

℞. Cran. human. ʒj uſneæ cran. hum. gr. x. L. O. gr. j. N. M. toſt. ℈j f. pulv. ad dyſenteriam.

EBUR.

Plus terræ, ſed minus ⊖is volatilis habet quam Cornu Cervi, unde potentius eſt in abſorbendo acido, commendatur etiam ſpecifice ad nimium menſium & album fluxum & præcavendum abortum, datur ut cornu cervi.

℞. Raſ. ebor. ℈ij camphor. gr. vij f. pulv. cape ex ▽ roſar, nymph. ana ℥iß ad menſes decolores albos.

℞. Raſ. ebor. ſuccin. præp. C. C. præp. ana ʒj maſtich. ʒß lign. aloës ℈j Nuc. Moſch. ʒß f. pulv. ad fluorem album, cape vel ex decocto vel vino ſaſſafr.

UNICORUM VERUM.

Cum ebore convenit, decantatum tamen eſt in epilepſia puerorum item torminibus, & convulſionibus.

℞. Ebor. unic. cryſtall. mont. præpar. ana ʒj L. O. gr. v. N. M. toſt. ℈j f. pulv. in cholera dandus ex ▽ Menth. Cinamom. & ſyr. portulac.

℞. Unic. ʒß. ⊖ vol. C. C. gr. vij datur ex ▽ chærefol. ℥iß. c. ſyr. Card. B. ʒij contra variolas & morbillos.

UNGULA ALCIS.

Cornu Cervi per omnia reſpondet, epilepſiæ vero unice videtur dicata eſſe. Datur ut cornu cervi, incenditur quoque ad arcendos inſultus epilepticos.

℞. Ungul. alc. rad. pœon. ſem. pœon. viſc. quern. ana ʒj. ⊖ vol. C. C. ʒß. f. pulv. ad epilepſiam, cape ex ♒u ceraſorum nigrorum.

℞. Ung. alcis. caſtor. ana ℈j M. f. pulv. ad epilepſiam fœminarum pro ij doſibus.

BLATTA BYZANTINA.

Eſt ſubſtantia teſtacea ad corneam accedens, oblonga ſpadicea, pinguis, nata in conchis rubri maris, orificium vel aperturam conchæ claudens, ſaporis & odoris dum accenditur caſtaneam æmulantis. Ab ungula alcis nil differt, attamen paulo majore doſi data ſolvit alvum, datur ut ungula alcis.

℞. Blatt.

℞. Blatt. byzant. ung. alc. troch. d. charab. ana ʒß. injice prunis, hauri fumum ore & naribus ad arcendum epilepticum inſultum.

Recentiores hanc blattam pro ungue odorato veterum habent, æmulatur enim figuram unguis animalis.

MANDIBULÆ LUCII PISCIS.

Sunt oſſa mandibulæ inferioris, ſaporis & odoris nullius evidentis, integra capita proſunt, palmam oculis ♋ præripiunt in pleuritide & peripneumonia.

℞. Mandib. luc. piſc. C.C. præp. ana gr. xv. ⊖ vol. C. C. gr. iiij f. pulv. cap. ex ▽ flor. ſambuc. contra pleuritidem, add. eſſent. caſtor. ʒj. ▽ flor. chamom. ℥iß. ſyr. artemiſ. ʒiij exhibe contra colicam convulſivam.

℞. Mand. luc. piſc. dent. apri. ♄ diaph. ana ℈j. ⊖ vol. C. C. gr. vj. f. pulv. cap. ex Emulſ. ſem. Card. ben. mar. papav. alb. ana ʒij c. ſ. q. ▽ flor. pap. rhœad. f. emulſ. contra pleuritidem & peripneumoniam.

℞. Mandib. luc. piſc. ſuccin. præp. ana ℈j f. pulv. cap. ex ▽ flor. nymph. contra gonorrhœam & fluorem album.

DENS APRI.

A mandibulis lucii piſcis nil differt, & cum his ſimul datur. ℞. dent. apr. ʒij detur ex ☍ lini rec. expr. ℥iiij per vices ad expectorationem & concoquendam materiam in pleuritide.

In inflammationibus internis. ℞. Dent. apri. ʒß. ⊖ vol. C. C. ℈ß. ex ▽ papav. ſcabioſ. ana ℥j.

OS SEPIÆ.

Piſcis eſt, qui in pedibus habet ſuccum nigerrimum inſtar atramenti, quem Chinenſes cum brodio oryzæ vel alterius leguminis inſpiſſant & formant, & in univerſum orbem tranſmittunt, ſub nomine atramenti Chinenſis.

Eſt ſubſtantia teſtacea candida, levis, utrinque tumida, parte ſuperna duriuſcula, inferna fungoſa mollecula, leviter aſpera, friabilis, nata in dorſo piſcis Sepia dicti, ſaporis aliquantum ſalſi & ſubacris, odoris nullius.

Vi abſorbente & præcipitante potentius eſt quam Oculi ♋, in ſpecie laudantur in gonorrhœa, fluore albo & omnibus cæteris fluxibus, intenſiorem vim adſtringentem acquirit, ſi ei affundatur -♌-us ⊕li, eoque iterum abſtracto, reſiduus pulvis reſervetur uſui, qui à ℈j ad ʒß datur. Specificum eſt in quavis ſiſtenda gonorrhœa.

℞. Os ſep. borrac. calcin. ℥ß. Nuc. Moſch. ʒß. conſ. roſ. r. nymph. ana ℥j ſyr. nymph. roſar. ana q. ſ. f. Elect. in gonorrhœa.

℞. Os ſepiæ. C. C. uſt. pumic. præp. ana ℥iij Moſch. ℈ß. inſperg. -♌- roſar. q. ſ. f. pulv. ad dealbandos dentes.

OCULI CANCRI.

Sunt lapilli cruſtacei, duri, albicantes, glabri, orbiculati, piſo majores, una parte compreſſi & ſinuati, altera convexi, nati utrinque unus in ventriculo cancri fluviatilis aſtacei, ſaporis & odoris nullius evidentis. Scilicet circa dies caniculares languere incipiunt, non nutriuntur, exanimantur fere, in eorum ventriculo

culo reperitur materia quædam lactea, quæ ſenſim poſt 6 vel 8 dies concreſcit in lapillos, qui deinde ſenſim iterum conſumuntur, & ut probabile eſt in reſtaurationem cedunt, exuvias enim tum deponunt & reſumunt.

Incidendi diſcutiendi & abſorbendi virtute ſuperant cornu cervi, pellunt quoque urinam & ſudorem, in ſanguine congrumato diſſolvendo ſpecificam habent virtutem, in ſananda hectica peculiare quid obtinent.

℞. Ocul. ♋i ʒij Corall. præp. limat. ♂tis ana ʒiß. rad. ari ℈j f. pulv. in bulimo & fame canina ad abſorbenda acida.

℞. Ocul. ♋. ℈j. ⊖ abſ. ℈ß Nuc. Moſch. gr. vij f. pulv. digeſtivus diurnus, add. L. O. gr. j. f. pulv. digeſtivus nocturnus.

In febribus intermittentibus ℞. Oc. ♋. præp. ℈j Bez. min. vel ♁ diaph. vel arcan. dupl. Mynſ gr. xv. ⊖ abſynth. vel ammoniac. ℈j 🝆d. caryoph. vel junip. gt. j. M.

Famoſus ille pulvis contra caſum fit h.m. ℞. Ocul. ♋i ʒj ſperm. cet. mum. ſang. dracon. ana ʒß cape in acet. deſt. ℥j ▽ chærefol. ℥ij.

Familiaris mihi eſt in Noſocomio pulvis ſequens antihecticus. ℞. Ocul. ♋. rad. ir. flor. ana ℈j flor. 🜍is ℈j. antihect. Poter. ʒj L. O. gr. j.

SANGUIS HIRCI.

Eſt ſanguis hirci exſiccatus, quem Helmontius & alii prius plantis aromaticis, lithontripticis alendum voluere, hinc angendum inſigniter, & intra extremas anguſtias pone aurem incidendam eſſe arteriam & colli-

gendum

gendum ſanguinem, alii hujus teſticulis reſectis diligenter colligunt, ſed hoc perinde eſt, conſtat enim ſale volatili copioſo, terræ incluſo. Ob copioſum ſalem volatilem alcalicum & terram parciorem ſuperat oculos ♋i, præcipue virtute diſcutiendi, incidendi & urinam movendi.

℞. Sang. hirc. ʒß cape ex ▽ flor. papav. contra pleuritidem & peripnevmoniam, ex ▽ chærefol. ad ſanguinem congrumatum diſſolvendum.

℞. Ocul. ♋ ſang. hirc. ana ʒß cape ex aceto ad ſanguinem congrumatum reſolvendum.

MUMIA.

Eſt ſubſtantia reſinoſa, indurata, nigra, ſplendida ſaporis ſubacris, amaricantis & reſinoſi. Accedit virtute diſcutiendi & incidendi ad ſanguinem hirci, aſt ob reſinoſas ſimul ipſis adjunctas mulcens eſt anodynum.

℞. Oc. ♋i. ʒj. rhabarb. ſperm. cet. mum. ſang. hirc. ana ʒß f. pulv. ad ſanguinem congrumatum diſſolvendum, datur ex aq. chærefol. in empyemate. [Mumiam nunquam interne exhiberem.]

℞. Mum. ſuber. uſt. terr. ⊕li dulc. ana q. v. add. ſucc. plantag- q. ſ. f. liniment. in ſiſtendo fluxu hæmorrhoidum.

MARGARITÆ.

Sunt partes rotundæ vel anguloſæ, albæ pellucidæ, ſaporis terrei, excrementitiæ ſcirrhoſæ, conchæ, neque enim in omnibus reperiuntur, tunc enim ſunt inſipidæ, cum alias ſint ſapidiſſimæ, quando nullas habent perlas. Pro oculis cancrorum nihil habeat peculiare,

culiare, niſi quod ob ſubſtantiam molliorem in ventriculo magis convincibiles & ſubtiles, quare in phthiſi & hectica commendantur.

℞. Margar. præp. ʒß ſpec. diambr. ℈j cape ex Emulſ. ſeq.

℞. Amygd. dulc. ℥ß. aq. Meliſſ. ℥ij aq. roſar. cinam. ana ℥ß. f. emulſ. add. conf. alkerm. ʒß. exhibe contra virium laſſitudinem & languorem, in hectica, phthiſi, tabe.

℞. Margar. ℈j. ⊖ abſynth. ℈j piper. alb. gr. vij. f. pulv. contra febres intermittentes.

Margaritæ in calcem redactæ ſunt ſpecificum in quartana, quod mihi & Medicis Orientalibus per experientiam conſtitit. Ita : ℞. Calcis margarit. ʒß ſacch. ♄ni ℈j. ⊖✳ci ℈j L. O. gr. j. f. pulv. vel ℞. Marg. calcin. ℈ß. ⊖ abſ. gr xv. Nuc. Moſch. toſt. ℈ß. 🝆 caryoph. gt. j. [Mater perlarum eſt teſta valida & ponderoſa, extus rugoſa, & aſpera non ſtriata, intus lævis argenteo colore prædita, & hæc teſta ejuſdem naturæ & qualitatis eſſe videtur cum margaritis, niſi quod ſit ſubſtantiæ durioris & ſolidioris, hinc non neceſſe eſt, ut margaritæ in calcem pro pulvere antifebrili redigantur ſed ex matre perlarum optime fieri poteſt, eodem modo, ſicuti ex conchis vulgaribus paratur ſpecificum antifebrile Crollii]

BEZOAR ORIENTALIS
ET OCCIDENTALIS.

Orientalis eſt ſubſtantia lapidea, foris ſplendida, glabra, quaſique lævigata, coloris foris atro virentis, intus obſcure cinerei, forma modo rotunda, modo oblonga, tereti vel ovali, magnitudinis variæ, com-

pacta ex pluribus lamellis ſibi invicem incumbentibus, oſtendens in meditullio cavitatem notabilem, continentem paleas, pilos, fibrillas, aliaque ſimilia, nata in ventriculo caprarum ſylveſtrium Indicarum, ſaporis & odoris nullius evidentis.

Occidentalis eſt ſubſtantia lapidea, orientali fragilior & laxior, foris ſcabra & minus lævigata, coloris modo ex viridi nigricantis, modo cinerei, ex multis quaſi tunicis ſibi invicem incumbentibus compacta, forma ut plurimum rotunda, magnitudine varia, habens in meditullio cavitatem, includentem paleas, pilos &c. nata in ventriculis caprarum Peruanarum & Mexicanarum.

Non in omnibus reperiuntur, ſed partes excrementitiæ earum quæ morboſæ ſunt videntur, uti notavimus de margaritis. Eſt hoc proprium in India occidentali, quod vix animal reperiatur in quo non generentur lapilli tales, & omnes ſub hoc nomine ad nos deferuntur, paucioribus ⊖ibus volatilibus gaudet, quam orientalis.

Uterque cum margaritis & oculis ♋ convenit, ſed ſubſtantia ſubtilior eſt, & in melancholia, febribus malignis pro ſudorifero habetur, in virium languore ab acido præ cæteris laudatur. Datur à ℈j ad ʒß imo ʒj niſi pretium obſtaret. Refertur quoque inter anthelmintica, forte quatenus acida imbibit, quæ nidus ſunt vermium.

℞. Lap. bez. ℈j. abſynth. gr. xv. ☿ dulc. ℈ß f. pulv. contra vermes ex ▽ graminis vel hyperici vel decocte filicis.

℞. Lap. bez. ℈j ⊖ volat. ſuccin. gr. vij. f. pulv. dandus ex ⏝u lil. convall. contra epilepſiam.

℞. Lap.

℞. Lap. bez. ℈j Croc. ♂t. ℈ß. ⊖ vol. C. C. gr. v. Croc. or. gr. vij. f. pulv. in melancholia ex aq. fumar.

DENTALI ENTALI. UMBILICUS MARINUS.

Sunt conchæ maritimæ. Alcali volatili & terra gaudent & ad acida infringenda veluti oculi cancrorum exhibentur, ſed ob ſubſtantiam duriorem poſtponuntur oculis ♋, certis tamen in caſibus.

℞. Dental. tere in pulv. exhib. ʒß cum aq. deſt. cinamom. vel alia ad abſorbenda acida. Verſa in calcem fiunt ſpecificum contra quartanam, v. c.

℞. Calcis hujus ʒß. ⊖ abſynth. ℈ß. quibus addi poteſt ſacch. ♄ni L. O. exhibetur ex vino abſynthite.

* * *

PARTES ANIMALES SALE VOLATILI OLEOSO CUM ODORE FRAGRANTISSIMO CONSTANTES.

MOSCHUS.

Si optimus, debet eſſe 1. ex ferrugineo nigricantis coloris. 2. unctuoſus. 3. ſaporis grati & amaricantis. 4. odoris fragrantis non adulterati.

Eſt ſubſtantia grumoſa pinguis & unctuoſa, ſanguini congrumato ſimilis, collecta ex Dorcadis ſeu capræ Moſchiferæ Aſiaticæ folliculo, ſito in abdomine inter umbilicum & pudenda, ſaporis ſubacris amaricantis, odoris fragrantis & grati.

Ob volatilem ſalem ſpiritibus analogum obtinet in ſpecie virtutem analepticam, cardiacam, confortantem, ♀em promoventem, ſanguinem congrumatum diſſolventem & hyſtericam paſſionem tollentem.

℞. Moſch. ℈ß. ambr. gryſ. ℈iß. ſacch. cand. q. ſ. ☉ citr. gt. v. cinam. gt. ij. caryoph. gt. j. M. f. pulvis analepticus cardiacus ad ℈j cum vino vel ſyr. chermes exhibendus.

℞. Moſch. gr. iij detine ſub lingua, ſi ex accepto vulnere thoracis nullus ſanguis effluit, illicò ſanguis fluere incipiet.

℞. Ladan. ℥iij. ſtyr. cal. ℥j benz. ℥ij flor. benz. ℥j Moſch. ℈ß. ambr. gryſ. ℥ß. ☉ Cort. Citr. rec. expreſſ. gt. xxv. mucag. tragacanth. cum ▽ roſ. fact. q. ſ. f. pomum ambræ odoriferum analepticum.

℞. Moſch. gr. v. extract. caſtor. inſpiſſ. ℈ß. mac. gr. iiij f. pil. deaurentur fortiter ad promovendum conceptum.

℞. Coagul. lepor. gr. v. Moſch. gr. j. Elect. diaſatyr. ℥j ambr. gr. iij ☉ majoran. gt. iiij ſyr. gran. cherm. q. ſ. f. Elect. paulo ante coitum.

ZIBETHUM.

Eſt ſubſtantia unctuoſa flaveſcens pallida, melli vel butyro forma & conſiſtentia ſimilis, ſaporis ſubacris, odoris fragrantiſſimi, collecta ex folliculo animalis Zibethici vel majoris vel minoris, ſito intra anum & pudenda, in fœmina, in mare inter penem & teſtes.

Hujus uſus potius externus quam internus, quia multum pinguedinis habet nauſeoſæ, eſt nempe antepilepticum, antihyſtericum, nervinum & anodynum.

℞. Zibeth. gr. iij Caſtor. ℈j conſ. fl. anth. ſyr. matricar. q. ſ. f. Electuarium antihyſtericum.

℞. ☉

℞. 🝆 majoran. gr. iij Zibeth. ʒß. 🝆 pin. recenter expreſſ. q. ſ. Moſch. gr. v. f. unguentum quo inungatur penis.

℞. Zibeth. ʒß. 🝆 laur. expr. ℥ß. camph. ℈ß f. liniment. in colica.

℞. Moſch. gr. iij Zibeth. ʒß. 🝆 pin. recenter expreſſ. q. ſ. ambr. gryſ. ʒj f. unguentum quo inguina & lumbi ſunt inungenda ad ♀em ſtimulandam præcipue in feminis.

CASTOREUM.

Non ſunt teſticuli caſtoris, ſed excreſcentiæ quædam glanduloſæ inter pudenda, quæ in medio fovent materiam melleam, quæ eſt quaſi quinta eſſentia. Sic multa alia animalia iiſdem tumoribus, qui cum genitalibus nullum plane commercium habent, gaudent, uti capræ moſchiferæ, quæ tamen ſimplicem tumorem habent, felis Zibethi qui duplicem habent, quorum ex glandulis liquor etiam pinguis emanat. In Canada mures ſunt aquatici, qui valde conveniunt cum Caſtoribus, habent etiam tumorem unicum, qui exciſſus ſpirat moſchum fragrantiſſimum, ſed tamen nihil moſchi reperitur etiam in tumore ipſo.

Teſticuli caſtoris foras non propendent, ſed ſunt in ipſo abdomine.

Eſt ſubſtantia glanduloſa, oblonga, rotunda nata in inguine ſubtus teſticulis ovi magnitudine, veſtita pellicula craſſa & impleta liquore oleoſo & ceroſo tractu temporis ſeipſum ex ſiccante, ſaporis acris amaricantis nauſeoſi, odoris gravis.

Ratione ⊖is volatilis penetrantiſſimi pinguedini incluſi ſpecifice convenit in morbis ab acido auſtero

ortis, morbis ſoporoſis, paſſionibus colicis, hyſtericis, convulſivis, in virginum cachexia, in ſingultu, cardialgia, palpitatione, variolis & morbillis. Eſſentiam Caſtorei ex oleo ſpicæ diſſolut. in colica convulſiva cum ſucceſſu exhibuit D. Hermann. doſ. à gt. x. ad xx. Externe naribus admotum torpidos ſpiritus in morbis ſoporoſis excitat.

Datur à ℈j ad ʒß in majori doſi datum alvum ſolvit.

℞. Caſtor. ℈j. L O. gr. ij ☍ d. chamom. gt. ij ocul ♋. ʒß. M. f p. divid. in duas part. æq. ad omnis generis dolores ex acido.

℞. Caſtor. ℈j ſcammon. præp. gr. xv. f. pulv. dandus ex oxym. contra morbos ſoporoſos.

Ego mihi paravi nobiliſſimam eſſentiam ſequentem: ℞. Rad. ari. ʒij pyretr. ℥ß. caſtor. ʒij. ♎ ⊖✳ci q. ſ. f. eſſent. in aphonia, applica cum linteaminibus vel detineatur in ore. it. in tuſſi.

***

PARS ANIMALIS SALE VOLATILI CUM MULTA PINQUEDINE CONSTANS.

SPERMA CETI.

Optimum eſt quod eximitur ex certis cavitatibus proceſſuum cerebri Orcæ piſcis, parte poſtica utrinque unus, it. canibus Carchariis Indis, item ex aliis piſcibus majoribus Indiæ orientalis.

Eſt ſubſtantia gummoſa, alba, tenera, mollis, pinguis oleoſa, quaſi ex parvis raris furfuraceis ſquamis compoſita, balænarum, orcarum, præcipue ceti dentati

tati & aliorum piſcium vel poſt mortem effuſa & in maris ſuperficiem acta, vel ſtudio ex cranio ſectove cerebro exempta, beneficio lixivii ſæpius purificata, quouſque ad albedinem redigatur, ſaporis pinguis butyracei & viſcoſi, odoris grati debilis, quod in ſuperficiem maris eructatur, invenitur rarius.

Salem ſuum volatilem caſtoreo blandiorem ſubſtantiæ pingui habet incluſum, unde præter virtutem diſcutientem & diureticam obtinet ſingularem vim anodynam, hinc audit anticolicum, antipleuriticum, antihyſtericum, antiaſthmaticum, anticatarrhale, in ſanguine & lacte congrumato diſcutiendo præ cæteris ſpecificam habet virtutem.

Datur ℈j vel ʒß c. vitell. ovor.

℞. Sperm. Cet. ʒß. caſtor. gr. vij f. pulv. contra pleuritidem, exhibe in vitello ovorum, ſuper bibendo aquam chærefolii.

℞. Sperm. cet ʒß. ♉ diaph. gr. xv. cape contra lac congrumatum.

℞. Emplaſtr. d. ſperm. cet. applica mammis ad lactis diſcuſſionem.

Sequens cataplaſma in coagulato lacte & hinc mammis tumidis commendatur. ℞. Sperm. cet. ʒj croc. or. ʒß. cum mica panis & vino generoſo f. cataplaſma.

*
* *

ANIMALIA SALE VOLATILI BLANDO CONSTANTIA.

SCINCUS.

Eſt animal quadrupes oviparum, lacertæ ſimile, niſi quod dorſum habeat latius magis glabrum & minus

nus ſerratum, cute ſquamoſa albicante tectum, cœſia à capite ad caudam decurrente linea, in aquis degens, potiſſimum maris rubri, inteſtina rejiciuntur & exiccatur.

Sale volatili blando ſcatet, ob quod diureticum eſt, an vero habeat ſimul vim alexipharmacam, aphrodiſiacam & adverſus lepram uti veteres putarunt, experientiæ committendum, interne datur à nonnullis ejus caro ad excitandam ♀em.

℞. Carn. ſcinc. formicar. alat. ana ℥j Elect. diaſatyr. ℥iij ſyr. cherm. q. ſ. f. Elect. ad proritandam ♀em; add. ſemen eruc. & eryſ. pro ſtimulo.

Cineres cum ⊖e volatili conjuncti laudantur in hydrope & in nephritide.

LUMBRICI TERRESTRES.

Sunt animalia apoda, exoſſia, digitum longa, craſſitie pennæ anſerinæ, mollia & carnoſa, coloris dilute ſanguinei, collo rubicundis angulis cincto, ſaporis nullius evidentis terrei.

Conſtant ⊖e diuretico & diaphoretico, hinc in ſpecie ad arthritidem & colicos dolores commendantur, in hunc finem dantur intra, & extra applicantur.

℞. Lumbric. terr. viv. ℥j affund. vin. ſ. ▽ gramin. f. infuſum diureticum.

℞. Lumbric. terr. viv. q. v. applica parti arthriticæ & inprimis vaga afflictæ.

℞. Rad. helen. ℥iß. lign. ſaſſafr. ℥ij herb. capill. Mij lumbric. vivor. ℥iiij paſſul. min ℥iß coq. ex ▽ f. apozema ad morbos ſcorbuticos & arthriticos.

℞. ℴℴ lumbric. terr. factum per coctionem, adde ℴℴ proprium deſtillatum per retortam, applica partibus arthriticis.

ANIMAL

*
* *

ANIMAL CAUSTICUM ET ACERRIMUM.

CANTHARIDES.

Sunt animalia oblonga, alata parva, vaginipennia, aureo colore viridantia, ſaporis primo nullius, hinc acerrimi picei, cauſtici erodentis, odoris cum recentes ſunt fœtidiſſimi, inveniuntur maxime in regionibus calidioribus, Hiſpania, Gallia, etiam in Germania ſuper cynogloſſam, ſed earum virtus brevi perit, acerrimæ tamen etiam ſunt, ubi recentes, ut plurimum gaudent fraxino, ut & liguſtro.

Vis cauſtica conſiſtit in ⊖e, infuſa enim fiunt cauſtica ex vino & -ſ-u vini, ſed ſimplici tantum non rectificato, alias ⊖ non ſolvitur, & ſ. q. cantharidum, hinc forte cujuſdam error, qui ſumſit forſitan rectificatum -ſ-um vini, vel quantitatem cantharidum non ſatis largam, ut ſcripſerit infuſa ipſorum non ſatis eſſe cauſtica. Cæterum quæ Olaus Borrichius fecit experimenta ſunt optima. [conf. act. Hafn. Volum. IV. obſ.80. p.m. 185. & Vol. V. obſ.89. p.m. 216.]

Harum vim diureticam & ſimul cauſticam & veſicatoriam, non ab imaginariis illis aculeis, ſed ab △is ſeu ⊖is particulis acerrimis dependere evincunt tum autopſia, tum experientia, à Segero aliiſque tradita. [conf. etiam Ehrnfrid. Hagendorn. C. N. C. Dec. II. ann. 1. obſ. 163. p. 388. ubi de virtute liquoris ex cantharidibus prolecti agit.] Quamvis tamen earum uſus ob vim cauſticam renes & veſicam exulcerantem ſuſpectus eſt, tamen in morbis nonnullis ad urinam

movendam, ad corrigendam gonorrhœam virulentam exhiberi poteſt, incipiendo à doſi minori, ſcilicet à gr. ß & ſi fiat cum ægri levamine ad gr. j. vel ij.

Frequentior earum uſus eſt externus in excitandis veſicis, ad capitis, aurium & oculorum dolorem, it. ad ſuffuſiones & tumores œdematoſos.

Abſque ſumma neceſſitate vix dantur interne, nam comperi vel quartam grani partem aliquando renes inflammaſſe, unde diureſes copioſæ, dolorificæ, imo cruentæ. vid. Wedel. Amœn. Mat. Med. Vis nimia cauſtica temperatur lacte vaccino vel ℅ lini expreſſo.

℞. Canthar. gr. j. datur in Emulſ. ſemin. 4. frigid. maj. ad diureſin excitandam.

℞. Canthar. ℈j Vin. rhenan. ℥iiij f. infuſ. cujus cochl. ß vel j cum vino vel decocto diuretico poteſt permiſceri & pervices exhiberi, præcipue hydropicis ad urinam movendam quod commendatur à Willisio. [conf. Th. Bartholin. Cent. V. Hiſtor. 82. p. m. 159.]

Pro Veſicatorio dantur ſi recentiſſimæ gr. v. ſi annoſiores ℈ß. ad ℈j Veſicatoria conveniunt in oculorum inflammationibus, à lymphæ coagulo, in capitis dolore, morbis ſoporoſis, ſurditate. Fiunt pone aures, in nucha, dorſo, lumbis, brachiis, femore, ſura &c.

℞. Canthar. ℈j pulp. ficus q. ſ. f. Veſicator.

℞. Canthar ℈j ferment. piſt. ʒj.

℞. Canthar. ℈j. Empl. Citr. ʒß Thereb. q. ſ. f. veſicator. ſub forma emplaſtr.

℞. Infuſ. Cantharid. ℥ß. ▽ Nymph. portulac. gramin. ana ℥ij. ℅ thereb. gt. v. exhibe ſucceſſive hoc infuſum ceu extremum remedium in gonorrhœa virulenta

lenta inveterata cum ulceribus ſordidis, per penem etiam injicitur.

Hæc ſalia cauſtica minime infringuntur ab acidis, quidam poſſident ⌒um fœtidiſſimum Cantharidum, qui ſanguini affuſus fluiditatem ipſi reddit.

***

PARTES ANIMALES OLEOSIS CRASSIS CONSTANTES.

ICHTHYOCOLLA.

Eſt ſubſtantia ex albo flaveſcens glutinoſa excocta ex piſce Huſo dicto, cujus ventriculus, inteſtina, veſica coquuntur, quouſque in tenues poſſint diduci lamellas, poſt contorqueri & cogi in ſpiras, ſaporis glutinoſi, odoris nullius.

Pinguibus glutinoſis ſeu mucaginoſis cum paucis volatilibus conſtat, & propterea tanquam temperans & anodynum ſumitur, in ſpecie in dyſenteria, diarrhœa, temperanda & incraſſanda lympha acri. Jungitur decoctis & clyſteribus.

℞. Raſur. C. C. ℥iiij ichthyocoll. ʒij coq. c. ſ. q. ▽ commun. in conſiſtentiam gelatinæ. Hujus cape ℥j ▽ cinam. ℥ij ▽ Cichor. ℔j ſyr. violar. ℥j M. f. julapium in febribus ardentibus & malignis.

℞. Mucag. ichthyocoll. tragac. g. arab. ana ʒj. ſacch. alb. in ▽ puleg. ſolut. f. trochiſci contra acrem lympham, catarrhos ſalſos, inflammationem faucium & tuſſim.

Egregii fiunt clyſteres ad demulcendum acre dyſentericum. ℞. lact. chalybeat. ℥viij ichthyoc. ʒij coque ad ſolutionem, add. Balſ. Peruv. ʒj. vitell. ov. num. j.

***

ANIMAL SANGUISUGUM.

HIRUDINES.

Sunt animalcula apoda, exoſſia, longa, lubrica, lineis & punctis nigris variegata, ore triſulco; vivunt in paluſtribus.

Applicantur utero, podici, brachiis, pedibus, temporibus aliiſque partibus ad extrahendum ſanguinem, maxime in hæmorrhoidibus cœcis tumidis & dolentibus. Cum diſtentæ apparent ſanguine, nec ſua ſponte volunt decidere, aſperguntur ☍ ♀ri per deliquium, vel ſale quocunque.

***

PARS ANIMALIS TERREIS CONSTANS CUM PAUCO SALE VOLATILI.

FOLLICULI SERICI.

Sunt veſiculæ pallide flaveſcentes, ovi columbini magnitudine, textæ ex filo ſerico continuato ſæpe replicato, ex bombycum ore emiſſo, ſaporis & odoris nullius evidentis.

Ob paucum ⊖ volatile cardiaci habentur. Externe cinis eorum vulneribus mundificandis inſpergitur.

℞. Ciner. follic. ſeric. ℈j. ⊕l. Hungar. g. iij. inſperge vulneribus.

PARS

PARS TERTIA.
REGNUM MINERALE EXHIBENS.

MINERALIA SALE VOLATILI SALSO CONSTANTIA.

SAL AMMONIACUM.

EST ſubſtantia alba cryſtallina, ſalina, ex quintuplo urinæ humanæ, una parte ſalis communis, & dimidia fuliginis parata, ſaporis acris & urinoſi. Omne eſt factitium, ipſi Arabes rident nativum dari.

Ob ſal ſalſum obtinet vim diureticam, ſtomachicam, antifebrilem. Datur in ſubſtantia à ℈ß ad ʒß, jungitur variis ſpecificis.

℞. ⊖ ✳ci ℈j. ♀r. ⊕lat. gr. xv. ⊖ abſynth. ℈j f. pulvis digeſtivus.

℞. Sal. Ammon. Nitr. depur. ana ʒß. rad. liquir. ir. flor. ana ʒj f. pulv. contra phthiſin doſ. ad ℈j ex juſculo pingui.

℞. Sal. ammon. ʒß. Corall. ℈j. 8° d. junip. gt. ij M. f. pulv. antifebr. ad ij doſes.

℞. Gumm. ammon. ℥j. tragac. ʒj empl. d. mucilag. ʒij ſal. ammon. ʒiij. oo ſuccin. q. ſ. f. emplaſtrum imponendum ad tumores viſcerum.

In ophthalmia à Catarrhis. ℞. ▽ fœnic. ℥j ſal. ammon. gr. v. f. collyr. addi poteſt ſacch. ♄ni. Flores ſublimati proficui ſunt, in iiſdem affectibus, in quibus

Sal. Spiritus totus eſt alcalinus, excepto phlegmate, convenit hinc in omnibus morbis provenientibus ab acido, egregium eſt antifebrile.

℞. -ꝛ-us ſal. ammon. ʒij. ▽ Card. Bened. ℥iiij ſyr. abſynth. ℥ß. ⊖ abſynth. ℈ß L. O. gr. j. f. mixtura in febribus.

BORAX.

Eſt ſubſtantia pellucida glaciei inſtar, alumine durior, ex lapide nitroſo vel chryſocolla calcinata & in ▽ diſſoluta, cocta & in cryſtallos concreta, ſaporis acris ſalini & nitroſi. Scilicet in India orientali, eruuntur ex fodinis terræ nitroſæ, quæ leviter calcinatæ in pulverem tritæ coquuntur affuſa aqua, vel lixivio forti, hinc exponuntur ut concreſcant in cryſtallos, quæ rarius in India ulterius præparantur, ſed huc locorum tranſlata ulterius ſolvuntur, cryſtalliſantur, donec pelluciditatem adipiſcantur.

Urinam, menſes, fœtum potenter promovet, recipitur quoque inter compoſitiones aphrodiſiacas. Doſis à ℈j àd ʒß.

℞. Borac. ℈j, cinam. ℈ß. M. f. pulv. exhibetur in decocto puleg. vel roriſmar. ad remoram menſium, additur gt. j ꝏ cinamom.

℞. Myrrh. ℈j ſuccin. alb. gr. xv. Croc. or. gr. v. borac. gr. vij. f. pul. cap. & ſuperbibatur ▽ cinamomi.

13. ℞. Borac. ℈j Tinct. Caſtor. ʒj. ⊖ vol. ſuccin. ℈ß, ▽ Cinam. ℥j. puleg. ℥ij ſyr. cinam. vel menth. vel aurant. ℥j f. mixtura in uteri vitiis.

℞. Borac. Venet. ʒj. ⊖ vol. ſuccin. ℈ß. Croc. ♂t. aperitiv. ℈j. ꝏ cinam. gt. ij ſacch. cand. q. ſ. ad menſes provocandum.

℞. Elect.

℞. Elect. diaſatyr. ʒij borac. ℈j f. bolus ad ♀em ſtimulandam.

Borax calcinata ad expulſionem omnis ſalis & acidi volatilis, ſpecifice convenit in fluxibus alvi, ſeminis, & eſt nihil aliud quam terra ſtyptica, datur à ℈j ad ʒß in conſerva roſar. vel per ſe, vel congeneribus. ſic. ℞. oſſ. ſep. præp. borac. calcin. Nuc. moſch. toſt. ſacch. ♄ni. ℈j f. pulv. qui propinatur ex vino rubro, vel alio decocto adſtringente.

CHRYSOCOLLA CRYSTALLINA.

Eſt ſubſtantia cryſtallina alba, condenſata velut borax, facta ex chryſocolla nativa, diſſoluta & concreta in cryſtallos, ſaporis borace acrioris ſalini, odoris nullius. Principia eadem quæ boraci, niſi quod incluſa terræ copioſiori. In uſum cedit pictorum & aurifabrorum ad conſolidandum aurum & argentum.

***

MINERALIA SALE SALSO FIXO CONSTANTIA.

SAL COMMUNE.

Eſt vel marinum, vel fontanum, Sal marinum paratur coctione ignis vel ſolis, ubi in maritimis per paratas foveas & canales deducitur aqua marina, ubi accedente ſole quicquid humiditatis eſt avolat & conſumitur, reliquum à vento frigidiuſculo ultro inſpiſſatur & concreſcit in cryſtallos.

Vim habet diureticam incidentem, & ſimul eſt ciborum condimentum catholicum, ſtomachicum, ob

ob vim abſtergentem & potenter diſcutientem, additur ſtimuli loco clyſteribus.

℞. ⊖ decrepitat. piper. alb. ana, applica uvulæ prolapſæ & inflammatæ.

℞. Decoct. Emollient. ℥viij add. ⊖ comm. ʒij f. Clyſter.

℞. Sal. comm. ℈ß ſal. vol. ſuccin. gr. v. aq. chærefol. ℥iij ſyr. dialth. q. ſ. f. hauſtus diureticus.

Externe uſurpatur in doloribus colicis, artuum, capitis &c. ubi decrepitatum in ſacculis calefactum imponitur, forte quia humorem in poris iſtis in ſe ſuſcipit.

℞. Sal. commun. ℈j ſal. gemm. gr. xv. ſal. ſuccin. volat. ℈ß. ex vin. diuretic. ℥v. vel vj f. hauſtus antihydropicus egregius, add. roob. vel ſyr. juniper. ℥j.

SAL GEMMÆ.

Eſt ſubſtantia alba, cryſtallina ſalina, eruta ex terræ fodinis, ſaporis acris & ſalſi.

Sale communi potentius eſt, cæteroquin cum ipſo convenit, hincque hydropicis datur ʒß. Utplurimum tamen externe datur in clyſteribus, gargariſmatibus.

℞. ▽ portulac. plantag. vel ſucci ſemper viv. ana ℥iiij ſal. gemm. ʒij ſyr. violar. q. ſ. ad grat. pro gargariſm.

NITRUM.

Eſt ſubſtantia alba, cryſtallina ſalina, ex terra pingui ſalſa & nitroſa cum urina vel aqua in lixivium cocta, in cryſtallos concreta, ſaporis amaricantis & acris. Scilicet in tractibus Bengalenſibus inveniuntur terræ nitroſæ, ſaporis ſalſi acidi ſubacris, cui lixivium forte

ex

ex ſalicibus, vel ▽ vel urina affunditur, coquitur, inſpiſſatur & exponitur ut concreſcat in cryſtallos. Deinde repetita apud nos filtratione, concretione magis exaltantur, purificantur, tumque nitrum depuratum dicitur. Effloreſcit etiam aliquando ex rupibus & parietibus antiquis, ex alcali calcis & acido aëris.

Sale ſalſo mediæ naturæ inter volatile & fixum conſtat, gaudet virtute diuretica, refrigerante, temperat nimiam acredinem bilis, ſedat ſitim, & ſanguinis phlogoſes, præcipue in malignis, eſt quoque antaphrodiſiacum.

℞. Nitr. ſal. ammon. ana ʒij liquir. ir. florent. ana ʒß. f. pulv. antihecticus doſ. ℈j.

℞. Sem. agn. caſt. Nitr. depur. ana ʒij camph. ℈ß. L. O. gr. iij f. pulv. contr. nimiam ſalacitatem.

℞. Fol. ſalv. Mij ſem. naſturt. ʒij flor. malv. hortenſ. Miß. rad polypod. ʒiij coq. in ▽ fontan. ℞. Colatur. ℔ij add. Mel. roſ. ℥iß. Nitr. præp. ʒiij M. f. gargariſm. pro exeſione, inflammatione & affectu ſcorbutico gingivarum, faucium & laryngis.

In hæmorrhagia narium, nitrum in aqua ſolutum & naribus hauſtum profuit.

Lapis prunellæ nitro gratior eſt & temperatior, alias cum ipſo convenit, eſtque nihil aliud quam Nitrum cum ſulphure depuratum. doſ. ℈j.

℞. Lap. prunell. ℈j. ⊖ abſynth. gr. v. ocul. ♋. gr x ☍ junip. gt. ij M. f. pulv. exhibendus hora j ante paroxyſmum febrilem.

ALUMEN RUPEUM ET ROMANUM.

Eſt ſubſtantia modo alba modo ex albo rubicunda, pellucida, cryſtallina, ponderoſa, glaciei inſtar,

ex lapide aluminoſo rocha dicto calcinato & cum aqua cocto, in cryſtallos variæ magnitudinis & formæ concreta, ſaporis acris, urinoſi & adſtringentis.

Conſtat ⊖e ſalſo p. m. corroſivo, magis tamen ad acidum inclinante, tum volatili tum fixo, ut & terra ſtyptica, quæ elixiviato alcali & acido plane dulceſcit. Vim habet diureticam & antifebrilem, ſimulque relinquit in receſſu vim adſtrictoriam: Externe laudatur in ophthalmia & abſumendis carnibus fungoſis vulnerum & aliarum partium.

℞. Alum. nitr. Nuc. moſch. ana ℈j M. f. pulv. ſumendus hora j. ante paroxyſmum febrilem, ſi febris non cederet adde L. O. gr. j. & ſacch. ♄ni ℈ß ſic enim certiſſime febris profligabitur.

℞. Alum. ℈j lap. lyncis. ⊖ abſynth. ana ℈ß L. O. gr. j. f. pulv. in febribus intermittentibus.

℞. Alum. nuc. moſch. ana ʒß. M. f. pulv. in febribus intermittentibus infantum, pro duabus doſibus.

℞. Alum. crud. ʒj. Miv. cydon. ʒj f. bolus adſtringens præcipue contra vomitum ventriculi add. L. O. gr. j. Vel ℞. alum. ℈ß. terr. ſigill. Corall. præpar. ana ℈j.

℞. Alum. crud. ʒiß. flor. aquil. ſalv. ana Mij rad. ir. florent. ʒiiß Myrrh. ℈ij. Mell. roſat. q. ſ. f. linimentum ad dentium vacillationem.

℞. Alum. ſal. piper ana q. v. applica uvulæ inflammatæ.

ALUMEN PLUMOSUM.

Eſt ſubſtantia media inter terram & lapidem ex viridi caneſcens, pinguis, ex numeroſis tenuibus fibris plumoſis vel lanuginoſis ſibi invicem incumbentibus & ſa-

& facile ſeparabilibus compacta, eruta ex terræ fodinis ſubſtantiæ ſubadſtringentis ſubacris.

Transfertur ex Rhodo & Creta inſula magna in copia ad nos, habentur ibi montes integri. Differt ab Asbeſto, qui ex Japonia & China ad nos adfertur, ſimilis etiam materia circa Oxoniam reperitur, unde chartas ſcriptiles conficiunt.

Ob acutiſſimas ſuas particulas cuti inſperſum pruriginem excitat, in quem finem membris paralyticis ſolet applicari ad vellicandos nervos; & torpidum ſpirituum animalium motum corrigendum, cum tinctura Caſtorei, ſpiritu vini camphorato &c.

* * *

MINERALIA SALE SALSO VOLATILI ET CORROSIVO CONSTANTIA.

CALX VIVA.

Eſt ſubſtantia lapidoſa, ex albo cinerea, ex calcario lapide ignis violentia in calcem exuſta, ſaporis acris & mordacis.

Multo ſale alcalico prævalente & pauco acido, utroque tamen volatili & ſimul corroſivo cum multa terra conſtat; hinc in vulnerum acido deſtruendo, gangræna tollenda & ſphacelo compeſcendo, item in nervis vellicandis & paralyſi tollenda, ut & in ambuſtis & hydrocephalo externe uſurpatur.

Latet in calce viva admirandum contra calculum & podagram remedium, ita ut ab externo ejus uſu in podagra jam inveterata nodi & tophi feliciter diſſipati fuerint. vid. Stiſſer. diſſert. d. arthrit.

Recipitur à nonnullis inter injectiones contra gonorrhæam & clyſteres contra colicam cum alvo adſtricta.

℞. Raſ. lign. ſaſſafr. ℥j. ſem. aniſ. ʒiij paſſul. ℥ß. liquirit. ʒij. ▽ calc. vivæ ℔ij f. infuſum ex mente Williſii [Pharm. Rat. ſ. 4. c. 4. p.m. 98. & 99.] contra diabetem ex acido in ſanguine orto. Cujus doſ. cochl. j.

℞. Calc. viv. ℔ij. ♀r. crud. ℥iiij alum. ℥j coque cum ▽ fontan. ad coagulum uſque, cui affunde ſpir. vini rectificat. f. tinctura, cujus gt. xx. in urinæ ſuppreſſione, item nephritide dantur.

℞. Lixiv. calc. vivæ, impone capiti contra hydrocephalum.

℞. ▽ calc. vivæ ℥iiij aloës myrrh. ana ʒß. mell. ʒvj. f. injectio contra gonorrhœam.

Si juſta doſi exhibeatur egregium eſt in febribus deſperatis. v. c. ℞. Calc. viv. gr. x. ad xv. Ocul. ♋i ℈j f. pulv. nonnulli audaciores ad ℈j aſcendunt, cum ſacch. ♄ni ℈ß & L. O. gr. j, pro robuſtis. [Sed audacia imo temeritas erit in tali qui hoc remedium præſcribet.]

℞. Rad. ari. pyretr. carlin. ana ℥ß. ▽ calc. vivæ q. ſ. f. infuſum, quod petiis exceptum adplica membris paralyticis ad concitandum ſpirituum affluxum.

ARSENICUM ALBUM ET RUBRUM.

Album eſt ſubſtantia ponderoſa, albicans, gypſi inſtar condenſata, foris cryſtalli inſtar ſplendens, facta ex auripigmenti cum ſalis communis anatica portione, ſaporis acerrimi & exurentis.

Salibus

Salibus ⊖ſis, 🜍eiſque volatilibus acerrimis exedentibus conſtat, humanæ naturæ averſiſſimis. Extrinſecus tantum adhibetur ad abſumendas carnes, clavos, ſcirrhos, pilos, item ad ulcera ſerpentia & cancroſa. Interne non ſine ſummo cum periculo in febribus intermittentibus uſurpatur.

℞. Vitriol. calcin. ℥iiij arſenic. ſulphur. ana ℥j ſal. gemm. ℥ß miſc. & cum aceto in vaſe clauſo calcina, calx inſpergatur ulceribus cancroſis.

℞. Arſenic. alb. irroretur cum aceto, impon. clavis & ſcrophulis abſumendis, interim vicinæ partes muniantur emplaſtro de mucilaginibus.

Arſenicum rubrum ſeu ſandaracha Græcorum eſt ſubſtantia metallica condenſata, ponderoſa, ex flavo rufeſcens & ſplendens, facta ex auripigmento in ollam clauſam injecto & ad rubedinem calcinato ſeu cocto, ſaporis exurentis.

Uſurpatur à pictoribus, poteſt tamen in caſu neceſſitatis ſubſtitui arſenico albo.

Oleo per deliquium utimur in ♋o aperto.

AURIPIGMENTUM.

Cum arſenico omnino convenit, & eodem modo externe uſurpatur.

CINNABARIS FACTITIA.

Eſt ſubſtantia metallica, ſolida ponderoſa, rubicunda, numeroſis ſubinde decurſantibus fibris ſeu ſtriis, ſplendentibuſque micis interpolata, facta ex una parte ſulphuris & tribus mercurii vivi, vi ignis ſublimata.

Datur à gr. iij ad v, robuſtiſſimis tamen, & qui aliis mercurialibus non moventur, cum conſerva meliſſæ, theriaca. Sale enim volatili corroſivo tam ſurſum quam deorſum vehementer purgante conſtat. Frequentior ejus uſus eſt externus in ſuffimigiis ad excitandam ſalivationem, & curandam luem veneream, & ſcabiem Gallicam, calloſque abſumendos.

℞. Cinnabar. factit. gr. iij. ſucc. liquir. q. ſ. f. pil. ad vomitum & ſalivationem promovendam.

℞. Cinnab. factit. ℥ij vel iij adde ſi lubet thuris, ſtyrac. balſam. Tolut. ana ʒv. M. f. pulv. inſperge per vices carbonibus ignitis, quibus in ſella foraminata æger locatus & operculo tectus, capite excepto, ſuperſedeat, ita ut fumus totum corpus pervadat, & ſudores expectet, hac fumigatione per 4 ad 8 dies continuata prodit ſalivatio & ulcera cutis Gallica exſiccantur.

℞. Folior. nicotian. conciſ. ℥j. cinnab. factit. ℥ß. inde tubulo tabacali & hauri fumum ad ſalivationem.

Cinnabaris nativa alia eſt, quæ modo in ☉i modo in ☽ti fodinis reperitur, ex qua fit mercurius vivus; per ſe non utimur, nam à ventriculo ſolvi non poteſt.

MERCURIUS SUBLIMATUS.

Eſt ſubſtantia alba cryſtallina, ponderoſa, ſplendens, ſublimata ex part. j. ☿ vivi & duplo ⊕li calcinat. & ⊖is; ſaporis acerrimi & exurentis. Conſtat ⊖e ⊖ſo volatili acerrimo & corroſivo vomitum & ſedes vehementer cauſante, inſequente ſalivatione, quare interne rariſſime exhibetur, nam venenis acerrimis merito annumeratur, externe vero in ulceribus Gallicis & ſordidis mundificandis locum habet.

Adjun-

Adjungitur loco ſtimuli unguento Neapolitano ad ſalivationem promovendam. v. c. ℞. Ung. Mercurial. ℥ij ☿-♌-ti gr. v. f. unguentum pro excitando fluxu ſalivæ.

℞. ☿-♌-ti gr. ij includ. ſucc. glycirrh. ſ. q. f. pil. involvantur auro foliato, deglutiantur, & ſuperbibatur 🜄 deſtillata menthæ vel juſculum pingue largiori doſi ad vomitum promovendum & ſalivationem. N. hoc tantum robuſtiſſimis exhibendum eſſe, & qui blandioribus non moventur.

MERCURIUS PRÆCITATUS RUBER.

Fit ex affuſa aqua forti Mercurio vivo, qui hinc coit in magma, quod in igne reverberii ad rubedinem tendit. Mitior eſt multo quam mercurius ſublimatus, unde interne exhiberi poteſt, cum circumſpectione tamen, tum ad vomitum & ſedes, tum ad ſalivationem promovendam. Externe ſtimuli loco additur unguentis pro ſordidis ulceribus mundificandis.

℞. Merc. præc. rubr. gr. iij theriac. ʒß. f. bolus ad excitandum vomitum & ſalivationem.

℞. Pil. Maſtichin. ℈j Merc. præcip. gr. j. ℅ ſuccin. q. ſ. f. pilulæ in dolore capitis ex lue venerea.

℞. Ung. Mercurial. ℥iiij Merc. præc. rubr, ʒj f. unguentum ad promovendam ſalivationem.

Mercurius præcipitatus albus mitior eſt rubro.

MERCU-

MERCURIUS DULCIS.

Eſt ſubſtantia alba cryſtallina, ſplendens, ſublimata ex Mercurii ſublimati & Mercurii vivi anatica portione, ſaporis & odoris nullius.

Inter mercurialia temperatior eſt & blandior, adeoque & pueris tuto exhibetur, purgat blande per inferiora & cæteris purgantibus loco coadjuvantis adjungitur, imo & diagrydio pro correctorio, in lue ♀ea curanda ſingulis diebus vel biduis exhibetur ad excitandam ſalivationem, laudatur quoque in expellendis lumbricis.

Datur à ℈ß. ad ℈j & ʒß in bolo vel pilulis, & quibus ad promovendam ſalivationem datur cum antimonio diaphoretico vel ſale volatili.

℞. Merc. dulc. ℈ß. Ocul. ♋. gr. xv. ☍ cinam. gt. j. f. pulv. digeſtivus præmittendus purgantibus.

℞. Rad. Jalap. ʒj. Merc. dulc. ℈ij ſacch. cand. ℥v. ▽ flor. hyperic. q. ſ. f. rotulæ purgantes contra vermes.

℞. Merc. dulc. gr. xv. ♁ diaphor. ℈ß. f. pil. cum ſucco glycyrrh. vel f. bolus c. theriaca vel conſerva flor. anthos, ad ſalivationem ſingulis diebus exhib.

℞. Pil. Maſtich. ℈j Merc. dulc. gr. xv. balſ. peruv. q. ſ. vel extract. caſtor. f. pil. purgantes anticatarrhales.

℞. Merc. dulc. ℈j Myrrh. ℈ß. Caſtor. gr. v. ☍ ſabin. gr. ij troch. alhand. gr. j. f. pil. cum extracto caſtorei ad menſes promovendos.

Riverius in ſuis ſcriptis ſæpe meminit Calomelani & eſt hoc: ℞. ☿ dulc. ℈j diagryd. gr. vij f. pulv.

MERCU-

MERCURIUS VIRIDIS.

Fit ex Mercurio & cupro ſolutis in aqua forti, quæ deinde abſtrahitur, pulvis deinde aceto irroratus viriditatem induit.

Principium activum eſt ſalſum Emeticum, mitius tamen Mercurio ſublimato, ſpecificum eſt in gonorrhæa, ratione forte cupri propter terram vitriolicam, doſ. gr. iij ad v. quam non excedere debemus niſi ſalivationem excitandi animus ſit.

In principio gonorrheæ ℞. ☿ vir. gr. iij extract. cathol. ℈j thereb. q. ſ. f. pil. Deinde his ſuccedere poterit ſeq. ℞. rhabarb. ʒj. ☿ vir. gr. iij & ubi ſimul conjunctus ardor. ſacch. ♄ni ℈ß. thereb. ʒß. f. maſſa ex vitello ovi continuetur hoc, quouſque fœtor ceſſarit. Hinc adſtringatur Rhabarb. Nuc. Moſch. toſt. Borrac. calcin. terr. ⊖li dulc. ana ℈j. ☿ vir. ℈j. f. pil. pro ſingulis doſibus ☿ij virid. gr. j.

In ulceribus penis feci inde candelas cum felici ſucceſſu: ℞. Merc. virid. q. pl. 🝪 ♄ni q. ſ. ceræ par. f. candela cum ſuo elychnio, ut pro lubitu poſſit extrahi peni intruſa.

TURBITH MINERALE.

Fit ex 🝪 ⊖li & ☿io, ubi poſt abſtractionem in fundo remanet pulvis albiſſimus, qui deinde ▽ ablutus mox flaveſcentem induit colorem.

Principium activum eſt ſalſum, cauſticum, emeticum catharcticum, operatur paulo blandius ſublimato & præcipitato.

Datur à gr. iij ad v. v. g. ℞. Extr. Cathol. ℈j turb. min. gr. iij f. pil. vel cum conſerva flor. perſicor. bolus.

***

MINERALE SALE VOLATILI ALCALICIS AFFINI CONSTANS ABSQUE TERRA.

MERCURIUS VIVUS.

Eſt ſubſtantia metallica mobilis fluida, ponderoſa, ex innumeris tenuiſſimis globulis conflata, coloris ex cæſio albicantis, facta ex cinnabaris minera deſtillatione per deſcenſum. Invenitur & nativus, ſed hic tantum inter rara aſſervatur.

Alcalicæ Naturæ Mercurium eſſe, probat 1. ejus efferveſcentia cum acidis 2. quoniam in omnibus morbis ab acido ortis v. c. Lue ♀ea egregie convenit. Volatilitas autem ejus fit manifeſta ex eo, quod △i impoſitus facile in auras abire ſolet. Crudus vel interne vel externe uſurpatus ſalivationem excitat, & vermes necat, qui ejus auram ferre nequeunt. In morbis ♀eis curandis ſpecificum eſt. Ratione gravitatis ſuæ, in colica, iliaca paſſione, & alvi fœcibus induratiſſimis cum fructu exhibetur.

℞. Merc. viv. ℈ß ſucc. glycyrrh. q. ſ. f. pil. ſingulis diebus exhibendæ, donec appareant futuræ ſalivationis ſigna, pro lue ♀ea curanda.

℞. Merc. viv. ℈ß detur ex melle ad enecandos lumbricos.

℞. Merc. viv. ℥j affund. ▽ gramin. & pone in digeſtione, exhibe decantatum infantibus contra lumbricos.

℞. Saturni ʒij Mercurii ʒj liqueſcant ſuper igne, forma globulos, qui in colica paſſione à fœcibus induratis ſumuntur, ſuperbibendo ☍ amygd. d. ℥j & ultra.

℞. Merc.

℞. Merc. viv. ʒij pingued. porcin. ℥j f. ung. quo inungatur corpus ſingulis biduis ad ſalivationem excitandam, donec ſalivationis ſigna appareant, in robuſtis augendus ☿ ad ℥j, pro correctorio adde aliquid camphoræ, & ſi vis 🝆 laurin. idem unguentum necat quoque pediculos.

⁂

MINERALE SALE VOLATILI SULPHUREO CUM TERRA CONSTANS.

ANTIMONIUM CRUDUM.

Eſt ſubſtantia metallica ſolida, ponderoſa, obſcuri cœrulei plumbei coloris cum longioribus radiantibus ſtriis concoloribus & aliquando rufeſcentibus compacta, concreta ex ejus minera liquata, & in figuram pyramidalem ut plurimum effuſa, ſaporis & odoris nullius evidentis.

Partibus 🜍eo alcalinis cum multa terra minerali conſtat, ob terram acidum imbibentem & evertentem, datur crudum in epilepſia, orta præcipue à vitio ventriculi, & cruditatibus primarum viarum. Ob 🜍 volatile aliqui addunt decoctis in morbis chronicis.

℞. ♁ crud. gr. iij rad. pœon. ℈j f. pulv. contra epilepſiam, dandus ex 🜄u ceraſorum nigrorum.

℞. Candel. ex cera alba factam cui admiſce ♁ crud. vel hujus loco crocum metallorum, inde meatui urinario ad tollendas caruncula s.

*In Dyſenteria maligna multi aſſumunt ℈j. ad ʒj ex conſerva roſarum.

Potiora medicamenta quæ ex antimonio parantur deſcribuntur in Collectan. Chymic. Leydenſ. p.m. 23. uſque ad 54.

***

MINERALIA SALE VOLATILI PURGANTE CONSTANTIA CUM TERRA ADSTRINGENTE.

LAPIS LAZULI ET ARMENUS.

Lapis Armenus eſt ſubſtantia lapidea ad metallicam accedens duritiem, pallide cœrulea cum quadam viriditate ex parvis quaſi grumis & intercurſantibus quibuſdam argenteis micis compacta, eruta ex fodinis argenti.

Lapis Lazuli eſt ſubſtantia lapidea ad metallicam accedens duritiem, ſaturate cœrulea, cum ſplendentibus quibuſdam aureis micis, armenio lapide durior & compactior, eruta ex fodinis auri.

Uterque poſſidet virtutem purgantem & emeticam & ſpecifice in morbis ſoporoſis, mania, apoplexia, ut & in quartana ſingulariter commendatur.

Toſtus ſeu ignitus ex quo expulſa ſunt ſalia volalilia, totus adſtrictorius eſt.

Doſ. à ʒß ad ʒj ſed raro, interdum enim in minima doſi purgat, interdum in maxima doſi non operatur.

VITRUM ANTIMONII.

Eſt ſubſtantia ponderoſa ſolida vitri inſtar, hyacinthini obſcuri coloris, facta ex antimonio calcinato & fuſo, ſaporis nullius evidentis.

Conſtat

Conſtat ⊖e alcalino, qui per ſuperiora & inferiora contenta vehementer expurgat, præcipue bilem. Raro datur in ſubſtantia, frequenter in infuſo, ubi menſtrui ratio eſt habenda, non ipſius vitri antimonii.

℞. Vitr. antim. ℥j affund. vin. hiſpan. q. v. infuſi hujus filtrati capi ℥j vel ℥iß vel per ſe vel cum ſyr. vel 🜄 cinamomi, vel quovis alio correctorio, ad vomitum & ſedes movendos.

CROCUS METALLORUM

Mitior eſt quam vitrum ♁ij, datur eodem modo.

℞. Croc. metallor. gr. iij detur ex melle ſeu oxymell. ad vomitum.

℞. Croc. Metallor. q. v. affunde vin. Hiſpan. part. ij vel iij, filtrati exhibe ℥j vel ij eodem modo quo infuſum vitri ♁ij exhibetur. Hoc infuſum ſtimuli loco adjungitur clyſteribus contra colicam, & morbum apoplecticum, item ad colyria exhibetur.

In Colicis paſſionibus cum ſcybalis duriſſimis.

℞. Decoct. Emoll. & carminativ. ℥viij infuſ. ♁ij ℥ij ⊖ commun. ʒij f. Clyſt.

***

MINERALIA SALE ACIDO CUM PAUCO ALCALI CONSTANTIA.

VITRIOLUM ALBUM. HUNGARICUM. ROMANUM.

Vitriolum album eſt ſubſtantia ſalina alba, ⊖is inſtar concreta, facta ex decocto vel lixivio terræ vitriolicæ, ſaporis acris ſubacidi & adſtringentis.

Hungaricum eſt ſubſtantia metallica tranſparens ex viridi cœrulea, ex terræ vitriolicæ decocto vel lixivio in cryſtallos concreta, ſaporis cum albo ejuſdem.

Romanum ſeu Cyprium eſt ſubſtantia cryſtallina cœrulea, tranſparens &c.

Vitrioli genera omnia conſtant ſale acido tam volatili quam fixo p.m. corroſivo cum pauco alcali & terra metallica. In ſale fixo alcali juncto conſiſtit vis emetica, in terra vis adſtrictoria. Album mitius eſt ad vomitum. Cætera vitriola fortiora ſunt, & ad externum potiſſimum uſum.

℞. Vitriol. alb. ʒj cape ex ▽ Mulſ. ad ℥iiij vel cereviſia, vel vino ad vomitum ciendum.

℞. Vitriol. alb. gr. iij ▽ fœnic. ℥j f. colyr. ad oculorum ſuffuſionem.

℞. Vitriol. alb. inſperge vulneribus mundificandis.

℞. Vitriol. Hungar. applica ulceribus ſordidis, ejuſdem virtutis eſt & Vitriolum Romanum.

℞. Vitriol. Roman. ʒij. ▽ commun. ℔j coque, injice in veſicam, ad mundificandum ulcus colli veſicæ ad mentem Foreſti.

CHALCITIS.

Eſt ſubſtantia metallica ponderoſa dura, glabra, rubicunda, æri ſimilis, multis intercurſantibus ſplendidis fibrillis, eruta ex venis æris, ſaporis acris & adſtringentis.

Cum vitriolo convenit, aſt in externo uſu potentior eſt. Chalcitis quæ nunc in officinis habetur, nihil aliud eſt quam vitriolum uſtum, & ſervit in officinis ad inducendam theriacæ nigredinem.

VIRIDE ÆRIS.

Sale ſalſo cauſtico gaudet, & ad ulcera mundificanda uſurpatur, abſumit quoque excreſcentias carnis præter naturales.

℞. Mell. roſat. miſce c. virid. æris q. ſ. applica vulneribus ſordidis.

℞. Virid. æris. arſen. alb. ana q. v. excipe tantillo ceræ & olei, applica verrucis & clavis pedum tollendis, ſeu ſolve cum aceto & applica.

***

MINERALE OLEOSIS SEU PINGUIBUS CUM ACIDO VOLATILI CONSTANS.

SULPHUR.

Multis pinguibus & paucis acidis gaudet, ob pinguedinem temperat & mulcet, & proinde in phthiſi, tuſſi, colica aliiſque caſibus laudatur; infringit quamcunque acrimoniam nimiam præcipue bilis. Externe temperat & abſorbet quælibet acria, ſcabiem & puſtulas generantia.

Datur crudum in ſubſtantia ſeu ejus flores à ℈j ad ʒß. Præſcribitur & ſpecie fumi, quo aquæ perfumantur, & à 🜍is acido imprægnantur.

℞. Sulph. crud. ℈j. cape ex ovo ſorbili ad tuſſim, & phthiſin ſingulis diebus ſeu cape ex pulpa paſſularum & ſyrupo papaveris.

℞. 🜄 ſcabioſ. ℥vj aſthmat. ℥iij diacod. ℥j cape cum flor. 🜍is ℈j pro tuſſi add. rad. ir. florent. ℈j.

℞. Rad. ir. flor. rad. ari. 🜍is ana ℈j f. pulv. add. flor. benz. gr. vij.

℞. Flor.

℞. Flor. 🜍is ℈j Zedoar. gr. vij L. O. gr. j. elæoſacch. chamom. ℈j f. pulv. contra colicam.

℞. 🜍is q. v. M. c. 🝆 laurin. f. ung. pro ſcabie add. ſi opus ☿ viv.

℞. 🜍 ʒß. fulig. ʒß f. pulv. in doloribus poſt partum.

***

MINERALIA FIRMIORIBUS PARTICULIS CONSTANTIA.

AURUM FOLIATUM.

Sunt ſubſtantiæ lamellatæ tenuiſſimæ, ex auro inter bina coria malleo cuſo & in lamellas redacto, commendatur ad compeſcendam ſalivationem nimiam à ☿io ortam.

Eo involvuntur pilulæ, ingreditur quoque confectiones & pulveres.

℞. Aur. foliat. q. v. ſucc. glycyrrh. q. ſ. f. pil. ad cohibendam ſalivationem ex ☿io, his ſolet addi reſin. jalap. gr. v. troch. alhand. gr. j. ſi alvus adſtrictior.

℞. Aur. fol. num. v. ung. camphorat. ℥ij quo inungatur corpus ad nimiam ſalivationem compeſcendam.

℞. Aur. fol. q. v. ſacch. in 🜄 lavend. ſ. q. ſolut. ℥j f. rotulæ ad compeſcendam ſalivationem.

MINE-

***

MINERALIA TERREIS CONSTANTIA.

LAPIS CALAMINARIS.

Eſt ſubſtantia media inter lapidem & terram, conſiſtentiæ friabilis laxæ, modo ex rufo modo ex albo flaveſcens, effoſſa ex fodinis metallicis maxime cupri, ſaporis nullius evidentis. Hujus uſus eſt in exiccandis vulneribus & ulceribus præcipue intertrigine.

℞. Lap. calam. ʒß croc. ♂t. ʒj Cret. albiſſ. Ceruſſ. ana ℈j f. pulv. pro intertrigine infantum.

℞. Lap. calam. q. v. miſc. c. 🝆 lini, applica herpeti.

CRYSTALLUS.

Eſt ſubſtantia ſolida, denſa, albicans, glaciei inſtar pellucida, dura, eruta è terræ fodinis, vel ſummis montium jugis, ſaporis & odoris nullius.

Ob potentem vim abſorbendi acida laudatur in epilepſia, & à nonnullis in cardialgia à bile æruginoſa. Datur ad ʒj.

℞. Cryſtall. præpar. ʒj. -∩- ceraſ. n. ℥j. ▽ flor. til. ℥ij cape pro epilepſia, ſed plus tribuendum vehiculo quam pulveri.

℞. Cryſtall. præpar. ʒj C. C. Corall. ana ʒß. L. O. gr. ij f. pulv. pro ij doſ. in cardialgia.

℞. Cryſtall. præpar. Croc. ♂t. ana ʒß. 🝆 d. lign. ſaſſafr. gt. j. f. pulv. contra icterum, ſi ſit à bile tenuiore & acriore.

℞. Cryſtall. præpar. ℈j lign. aloës ℈ß conſ. anthos ʒß. f. bolus in fluore albo.

Denique in defectu lactis mulierculæ vel interne ſumunt, vel externe appendunt dorſo. ℞. Cryſtall. præpar. lac. ☽æ ana ℈j. ▽ fœnicul. ℥ij f. hauſtus.

FRAGMENTA LAPIDUM PRETIOSORUM.

Ad acida primarum viarum abſorbenda efficacia ſunt, hinc forte eſt, quod in lypothymia, palpitatione cordis & aliis affectibus laudantur.

GRANATUS.

Sunt lapilli ex fuſco rubicundi, ſeu ex rufo nigricantes, parvi anguloſi, granis mali punici ſimiles, collecti ex variis terræ lapidoſæ locis.

Iis in diarrhœis deſperatis feliciter uſus fuit Hermannus. h.m. ℞. Granat. ʒj conſerv. roſar. ℥j f. bolus.

HYACINTHUS.

Sunt lapilli ex flavo rubicundi, pellucidi, ignis inſtar fulgentes, granatis rubedine dulciores, collecti ex variarum regionum variis fluviis vel eorum ripis.

Confectio de Hyacintho cardiaca eſt, quatenus abſorbet acida, quæ ſpiritus poſſunt tenere ligatos.

RUBINUS.

Sunt lapilli pellucidi rubentes, laccæ granulatæ ſimiles, collecti ex arenis ad ripas fluminum, de collo geſtatus ſubinde colorem perdit, ut & corallia ab effluviis acidis, quæ in ſe recipiunt, unde lixivio alcalico iis, uti coralliis color reſtituitur.

SAPPHI-

SAPPHIRUS.

Sunt lapilli pellucido diaphani, modo cœrulei, modo pallidi, collecti vel ex arena ad fluminum ripas, vel ex terris lapidoſis.

Oleum ſ. Q. E. Sapphiri commendatur in morbis malignis, ſed nota ſi flores chamomillæ deſtillantur in vaſe cupreo cum pauxillo ⊖is ammoniaci vel ꝏ thereb. tum eandem exhibebunt tincturam, quam falſo Sapphiri nominant.

SARDIUS.

Eſt lapis opacus, colore Hepati ſimilis, in ſuperficie variis ſubinde albicantibus vel etiam cœruleis ſtriis notatus, collectus ex ſaxis quibus adnaſcitur.

Hæmorrhagiam ſiſtit manu ſolum ſi retinetur.

SMARAGDUS.

Sunt lapilli pellucidi, virides, reliquis gemmis molliores, collecti ad fluminum ripas.

Contra pavores nocturnos infantum laudatur à Timæo.

TOPASIUS.

Sunt lapilli pellucidi, tranſparentes, aurei coloris, ad Solem radios de ſe ſpargentes, collecti vel ex terris ladidoſis vel ex arenis.

OSTEOCOLLA.

Eſt ſubſtantia media inter terram & lapidem, alba, friabilis, multis tuberculis exaſperata, cruſtacea, oblonga, ex baſi craſſiori in obtuſum deſinens mucro-

 nem,

nem, excreſcens ex arenis, aliiſque locis lapidoſis, ſaporis nullius evidentis terrei.

Præter facultates terreis communes obtinet ſingulare quid in generando callo, laudatur & in fluore albo & febribus intermittentibus.

℞. Oſteocoll. ʒj cape ex hauſtu vini, vel ex aceto ad conſolidanda oſſa & generandum callum.

℞. Oſteocoll. magiſter carniol. corall. alb. Smaragd. ana q. v. exhibe ʒj ex vino ſaſſafr. contra fluorem album.

℞. Oſteocoll. ebor. ſuccin. præpar. ana ʒiß. conſ. anthos f. bolus.

TUTIA.

Eſt ſubſtantia metallica ponderoſa, cruſtacea, corticis inſtar, intus lævis ex flavo cinerea, extus numeroſis exaſperata tuberculis, ex cœruleo pallido cinerea, congeſta è fuligine æris in fornacibus metallicis cum cadmia liquati & vel fornacibus vel ferreis piſtillis, quibus æs agitatur adhæret.

In inflammatione oculorum & hæmorrhoidum, ut & in vulneribus curandis & ulceribus ſiccandis exhibetur, exhibetur externe tantum.

POMPHOLYX.

Eſt pulvis tenuiſſimus farinaceus albicans, favillæ inſtar, ſublimatus ex cadmia & ære coctis in fornace, ſaporis nullius evidentis.

Convenit omnimode cum tutia.

℞. Tut. ʒß os ſep. ℈j ſacch. cand. ʒj. ⊖l. alb. ℈j f. pulv. ſubtiliſſimus inſpergendus oculis inflammatis. vel add. ▽ plantag. fœnicul. roſar. & fac. colyr.

℞. Tut.

℞. Tut. inſperge vulneribus ſordidis ad ea mundificanda.

℞. Ung. de tut ℥j. ſacch. ♄ni ʒij. ☿ viv. ʒj. Camph. ʒß opii ʒj. 🝆 lin. q. ſ. f. liniment. ad hæmorrhoidum dolorem, intrudendum penicillo convenienti in anum.

Similia medicamenta parantur etiam ex NIHILO ALBO quod eſt ſpodium Græcorum. Nam ſpodium Arabum eſt ebur uſtum.

PUMEX.

Eſt ſubſtantia lapidea levis, poroſa, cinerea, nata ex ſcoriis montium ignivomorum in mare projectis à ⊖ibus eroſis & poroſis redditus & in littus propulſis, ſaporis & odoris nullius evidentis.

Ob ſcabras particulas ſordes dentium abſumentes ingreditur quoque pulveres dentifricios, ad dealbandos dentes. Nonnulli etiam hunc lapidem præparatum, ut Timæus, ad nimios alvi fluxus ſiſtendos propinant.

℞. Pumic. ℥ß. Corall. rubr. albor. ana ʒiß alum. ʒj rad. ir. florent. ℥j f. pulv. dentifricius, adde olea deſtill. & ambram pro odore.

℞. Off. ſep. C. C. uſt. pumic. Corall. r. ana ʒj Nuc. Moſch. ʒß. f. pulv. irroretur tinctura coccinell. vel ipſis in ſubſtantia, vel ſantalo rubro ad colorem rubrum conciliandum, adde moſch. & ambram & ſi cauſſa ſimul peccat ſcorbutica add. ⊖ ammoniac. ʒj vel 🝆 cochlear.

Externe uſurpatur in intertrigine, exulceratione cutis & ulceribus manantibus. ℞. Pumic. lap. calamin. ana ʒj. tere in pollinem & inſperge.

Eodem modo uſurpatur SPONGIÆ LAPIS, qui naſcitur fruticis inſtar ex ſaxis, conchis, aliiſque marinis.

***

MINERALIA PARTICULIS TERREIS CUM ALCALI CONSTANTIA.

CORALLIA RUBRA, ET ALBA.

Rubrum, eſt ſubſtantia lapidea ramoſi fruticis inſtar excreta rubicunda, per longitudinem exiguis ſtriis donata, nata in ſcopulis, lignis conchis aliiſque ſubmarinis.

Album eſt ſubſtantia levis poroſa, ac cortice ſuo quo tegitur molleculo ferrugineo liberata &c.

Nigrum frequenter in China reperitur.

Corallia ſemen non habent, quod enim intra poros coralliorum albidum & viſcoſum reperitur, aqua potius marina eſt.

Falſum eſt Corallia ſub aquis eſſe mollia, inducere duritiem ab aëre, erroris cauſa hæc videtur, teguntur cortice molli poroſo, qui ad tactum facile cedit, imo ab ipſis aliquando tempeſtatibus diſcutitur, ſed rejecto cortice contrarium videmus.

Utrumque in terra, acida abſorbente, paucum alcali includit reſpectu Oculorum ♋i; laudatur in ſiſtendo vomitu, fame canina, fluxu alvi, in ſiſtendis menſibus, palpitatione cordis, febribus &c.

℞. Corall. ʒj. ⊖ abſynth. ʒß ſucc. limon. ʒij. ▽ Cinam. ℥ij f. potio ad Vomitum ſiſtendum, add. L. O. gr. j.

℞. Co-

℞. Corall. ʒj Cryſtall. præpar. ʒß C.C. uſt. ʒß L.O. gr. j. f. pulv. divid. in ij doſes ad cardialgiam.

℞. Corall. præpar. ℈j ſal. ammoniac. ʒß. 🝆 junip. gt. ij L. O. gr. j. f. pulv. antifebrilis.

Tinctura Corallorum à nonnullis ſequenti modo paratur. ℞. Corallor. in pollin. ſutiliſſ. tritor. q. pl. affunde 🝆 thereb. junip. vel butyr. relinque in digeſtione, ſic omnis rubedo extrahitur ex coralliis, decantant menſtruum, & huic affundunt ⁀um vini, qui omnem tincturam in ſe ſuſcipit, ſed objiciunt tincturam hanc etiam ab his menſtruis poſſe provenire, quod ut eluderem ſequenti proceſſu modo: Recepi butyrum non ſalitum, huic toties affudi ⁀um vini, quouſque butyrum orbatum omni colore, butyrum affudi coralliis, atque ſic omnem extraxit rubedinem, quam in ſe deinde recepit omnem affuſum alcohol vini & ſic veram obtinui tincturam corallorum.

UNICORNU FOSSILE.

Eſt ſubſtantia media inter terram & lapidem, exuſtis oſſibus ſimilis, cortice exteriore duro, flaveſcente aut ſubcinereo vel nigricante, medulla molli alba friabili, linguæ firmiter adhærente, ſaporis adſtringentis & exſiccantis.

Convenit cum coralliis, ſpecialiter laudatur in variolis & morbillis cum fluxu alvi; præſcribitur ut corallia.

℞. Unicorn. foſſil. ʒj C.C. ʒij terr. ſigill. ʒß. f. pulv. polychreſtus diaphoreticus conveniens in malignis, add. ⊖ C. C it. camphor. vel L.O.

℞. Unicorn. foſſil. cortic. ovor. ana ℈j œleoſacch. cortic. aurant. f. pulv. ad fluxum menſium

BOLUS.

BOLUS ARMENA: VULGARIS: ALBA.

Bolus Armena eſt ſubſtantia terrea ex flavo pallide rubicunda, ponderoſa, pinguis, facile friabilis, ex terræ fodinis effoſſa, ſaporis adſtringentis & pinguis.

Vulgaris eſt ſubſtantia terrea rubicunda &c. ſaporis terrei & adſtringentis.

His accedit alba, differt tantum colore.

Optima vero Armena, cujus terræ pinguedo cum pauco ſale volatili, ♂tialibus accedit, hinc ſudorem movet, & ſimul adſtringit, & contra eoſdem affectus, ad quos corallia, exhibetur.

℞. Bol. armen. diaſcord. Fracaſt. ana ℥j ſyr. Card. bened. q. ſ. f. bolus, in malignis.

℞. Bol. armen. ℥j conſerv. roſar. q. ſ. f. bolus contra ſputum ſanguinis.

℞. Bol. armen acet. deſt. q. ſ. f. catapl. applicandum fronti, ad ſiſtendas hæmorragias.

℞. Bol. arm. Cret. alb. Croc. ♂t. ana f. pulv. ad intertriginem.

TERRA LEMNIA ET SIGILLATA.

Eſt ſubſtantia argillacea ſebacea, pinguis, ex flavo rubeſcens, eruta ex terræ viſceribus in inſula Lemno & in placentulas coacta, ſaporis pinguis & adſtringentis.

Terra ſigillata eſt ſubſtantia argillacea ſebacea &c. cujus variæ dantur ſpecies.

Cum bolo ſymbolizant, qua tamen ſunt pinguiores & præſtantiores.

℞. Terr.

℞. Terr. ſigill. troch. d. charab. ana ℥ß. Corall. r. ℈j L. O. gr. ij f. pulv. vel cum conſerv. roſ. bolus.

℞. Terr. ſigill. ℈j L. O. gr. ij ℴℴ cinam. gt. j miv. cydon. q. ſ. f. bolus ad vomitum nimium.

℞. Terr. ſigill. affund. ⏝ ⊕li & ⦶tri donec in pultem abeat, ſicca, & poſt repone in cellam ut abeat in liquorem, qui liquor ſtipticus adpellatur, hujus cape ℈j vel per ſe vel ex ſyrupo cydoniorum, ad omnem hæmorrhagiam & fluores, utere quoque externe ad hæmorrhagias.

PLUMBUM.

Eſt ſubſtantia metallica ſolida, flexibilis tamen, ex certa minera in furnis fuſa, & per canales in proxima vaſa effuſa. Utimur tantum ejus præparatis.

LITHARGYRIUM AURI. ARGENTI.

Auri eſt ſubſtantia metallica, modo in glebas modo in pulverem redacta, ex fulvo ſplendeſcens, cum ſplendeſcentibus quibuſdam aureis micis, cocta in fornacibus ex ♄no, dum ☉ eo purificatur, imo etiam argentum, dum diutius tractatur in igne.

Argenti eſt ſubſtantia metallica ponderoſa, cinerea, cum micis ☽teis, cocta ex ♄no dum ☽ eo purificatur.

Uſurpatur externe ad exſiccanda ulcera it. in intertrigine, inflammatione.

℞. Litharg. 🜂 tragacanth. ana ʒj. 🜕 roſat. q. ſ. cum tantillo aceti f. liniment. ad anum exulceratam.

℞. Litharg. ceruſſ. ſacch. ♄ni ana q. v. affund. acet. c. 🜄 ſolani & tantillo roſar. f. lavamentum, applica inflammationibus, adde his pulveribus aliquid therebinthinæ cum vitello ovi ſubactæ, inde vulneribus.

In officinis habetur emplaſtrum tetrapharmacum, quod ulceribus manantibus, & tumoribus maturandis imponitur.

CERUSSA.

Eſt ſubſtantia alba condenſata, friabilis, ponderoſa, ex plumbi lamellis ſupra acetum ſuſpenſis & eroſis, in glebas coacta.

Convenit cum Lithargyrio & uſui externo tantum dicatur.

℞. Ceruſſ. ℥ij. 🜕 roſac. q. ſ. f. unguent. applica hæmorrhoidibus inflammatis.

Nota de omnibus ♄eis, quod ſpecifica ſint in omnibus herniis & rupturis, maxime à peritonæo relaxato, ubi coquuntur in 🜄 ſimplici vel 🜕 therebintinæ.

MINIUM.

Eſt pulvis rubicundus ponderoſus, factus ex ♄no igne reverberatorio & continua agitatione in rubicun-

cundum calcem uſtulatus, cujus pondus augetur, ſi perite tractetur; ex libris enim xi ♄ni obtinentur Minii libræ xij.

Ceruſſæ reſpondet & ad ulcera gallica & ambuſta it. lentigines & lichenes uſurpatur.

℞. Minii, ☿ præcipitat. ana ℥iiij. 🝆 commun. ℥ij Ceruſſ. Ceræ ana ℔ß M. f. emplaſtrum in tumoribus & ulceribus gallicis.

℞. Empl. d. min. ℥j ſacch. ♄ni ʒj. 🝆 roſac. q. ſ. f. emplaſtr. refrigerans in ardore & aliis malis.

SACCHARUM SATURNI.

Nihil aliud eſt quam 🜔 aceti in naturam ſubdulcem converſum, fit ex omnibus præparatis ♄eis; deſtillatum fundit ſpiritum ardentem, non acidum, caput mortuum fuſum redit in ♄num, cum quo eadem principia habet. Interne datur à ℈j ad ʒß.

℞. Sacch. ♄ni ℈j Ocul. ♋ ʒß. 🝆 d. caryoph. g. j. M. f. pulv. optimus in febribus.

In ardore urinæ. ℞. Sacch. ♄ni ℈ß Nuc. Moſch. gr. xv. Ocul. ♋. mandib. luc. piſc. ℈j. M.

FERRUM ET CHALYBS.

Chalybs eſt ferro compactior & ſolidior factus ex ferro ſæpius candefacto & in ▽ extincto. Ferrum ob ſubſtantiam minus ſolidam frequentius datur. Gaudet terra metallica, acida propterea abſorbente & ſale alcalico acida deſtruente, hincque

utroque nomine obſtructiones reſerat, & inde ortos morbos præcipue chronicos curat.

Datur pro ſcopo aperiente tum in ſubſtantia, tum in infuſo. Crudum & in ſubtiliſſimum pollinem redactum, præfertur omnibus cæteris præparatis, nam cum aceto ſi præparatur, eo tam copioſe ſaturatur, ut tantum adſtrictoriam habeat virtutem.

℞. Ferr. candefact. extingue aliquoties in vino Malvatico vel alio generoſo f. vinum chalybeatum.

℞. Limat. ferri recenter facti ℥j Zedoar ʒvj Vin. Hiſpan. q. ſ. f. infuſum in morbis chronicis, præcipue in hydrope imminente.

℞. Limat. ferri ʒj caryoph. arom. gr. v. conſ. flor. anth. ſyr. d. cinam. f. electuarium deobſtruens in morbis chronicis.

℞. Limat. ferri ℈j reſin. jalap. gr. x. ⊖ 🜿ri ℈ß 🝆 d. caryoph. gr. ij troch. alhand. gr. iij f. pulv. purgans acida.

Si excrementa non tingantur ex ferri uſu nigredine, non facile continuandum, remanet enim in Ventriculo.

CROCUS MARTIS.

Eſt pulvis ex flavo nigricans, factus ex ferro vel per calcinationem vel ſolutionem per acida, saporis adſtringentis. Eſt vel aperiens vel adſtringens. Aperiens fit quando lamina aut limatura ferri madefacta exponitur aëri vel rori, aut aqua conſpergitur,

gitur, ut rodatur in calcem. Adſtringens fit affuſo ⌒u acido, uterque conſtat terra metallica puriſſima & proprio ⊖ alcalico aperiente, quod tamen potentius in aperiente, uterque in receſſu relinquit vim adſtrictoriam.

LAPIS HÆMATITES.

Eſt ſubſtantia metallica mediæ conſiſtentiæ inter lapidem & terram, ponderoſa, rubicunda ex ſtriis vel lamellis ſibi invicem incumbentibus compacta, eruta ex terræ fodinis ferro gravidis, ſaporis adſtringentis.

Differt à ſuperioribus quod ſubſtantia ſit magis poroſa, ſubtilior & magis convincibilis in ventriculo. In ſpecie contra hæmorrhagias laudatur tam interne quam externe.

℞. Lap. hæmat. ʒß. Corall. ʒij ſacch. ♄ni ℥ij L. O. ℈ß. f. pulvis ad 9 vel 10 doſes in hæmorrhagiis.

℞. Lap. hæmat. ſacch. ♄ni liq. terr. ſigillatæ M. applica hæmorrhagiis externis.

Optimum Hermanni remedium in hæmorrhagiis fuit ſequens: ℞. Lap. hæmat. ʒj. ⊕l. alb. ℈ß phlegm. ⊕li q.ſ. excipe ſtuppis & formetur turunda: In narium hæmorrhagia ſufficit ⊕l. ad gr. ij ſumtum.

MAGNES.

Eſt ſubſtanttia lapidea ad metallicam accedens, ferruginea, polos mundi oſtendens, effoſſa ex variis terræ mineris ferro gravidis.

Ferrum virtute æmulatur & in ſubſtantia ad ʒß in omnibus morbis chronicis ab obſtructione pendentibus exhibetur.

℞. Magn. ʒj. 🜿r. ⊕lat. gr. vij 🝆 cinam. gt. j ſacch. cand. q. v. f. pulv. deobſtruens in morbis chronicis.

℞. Crem. 🜿ri. ℥ß. Magn. ʒj. 🜔 fraxin. gr. xv. ſacch. cand. ʒj f. pulv. aperitivus & diureticus in omnibus morbis chronicis & incipiente hydrope.

INDEX

INDEX MATERIÆ MEDICÆ.

A.

ABrotanum mas 214
Abſynthium vulgare 216
ponticum 217
Acaciæ ſuccus 296
Acorus indicus 5
vulgaris 4
Adianthum album 237
nigrum 237
Agallochum 109
Agaricus 120
Agrimonia 218
Alcanna 82
Alchymilla 236
Alcis ungula 300
Alhandal 161
Alkekengi baccæ 153
Aloe. lignum 107
Aloe. Caballina 283
Hepatica 284
ſuccotrina 284
Althæa radix 77
ſemen 194
herba 231
Alumen plumoſum 322
romanum 321
rupeum 321
Ambra liquida 277
Ammios ſemen 131
Amomum 123
Ammoniacum gummi 259
ſal 317
Amygdalæ amaræ 181
dulces 179
Anacardium 184
Anchuſa Monſpelienſium 82
Anethum 135
Angelica 12
Anime gummi 264
Aniſum Chinenſe 133
vulgare 132
Anonis 49
Anthophylli 122

Anthos

Anthos flores 238
Antimonium crudum 331
vitrum 332
Apium paluſtre 33
ſemen 142
Apri dens 301
Aquilegiæ ſemen 195
Arabicum gummi 291
Ariſtolochia & ſpecies 53
Armenus lapis 332
Armoracia 35
Arſenicum album
rubrum 324
Artemiſia 213
Arum 36
Aſa dulcis 271
fœtida 262
Aſarum 56
Aſparagus radix 48
ſemen 156
Aſplenium 237
Atriplex ſemen 175
Aurantiorum cortex 91
Auricula Judæ 119
Auripigmentum 325
Aurum foliatum 336

B.

Balanus Myrepſica 170
Balauſtiorum flores 252
Balſamum de Copaiva 276
Peruvianum 275
Tolutanum nigr. 275
Bardana. radix 46
ſemen 152. 195
Been fructus 170
Benzoë 271
Betonica 219
Bezoar Occidentalis
Orientalis 305
Biſtorta 82
Bitumen Judaicum 292
Blatta Byzantina 300
Boletus cervi 118
Bolus alba
armena
vulgaris 344
Bombacis ſemen 176
Borrago 249
Botrys 218
Borax 318
Boviſta 121
Bryonia 72
Bugloſſa 249
Burſa paſtoris 235

C.

Cacao fructus 185
Calaminaris lapis 337
Calamintha 210
Calamus aromaticus 6
Calendula 248
Calx viva 323
Camphora 254

Canca-

Cancamum 264
Cancrorum oculi 302
Cannabis ſemen 178
Cantharides 313
Capillares herbæ 237
Capparis cortex 102
Caranna 270
Cardamomum 127
Carduus Benedictus herba 225
ſemen 151
Carduus Mariæ 152
Caricæ 186
Carlina 30
Carpobalſamum 140
Carthamus ſemen 170
Carvi ſemen 136
Caryophyllata 84
Caryophylli aromatici 122
Caſſia fiſtula 173
caryophyllata 95
lignea 95
Caſtanearum cortex 103
Caſtoreum 309
Catechu 293
Cataputiæ ſpecies 168. 169
Cauda equina 233
Centaurium minus 234
Cera 293
Ceraſorum nuclei 183
Ceratonia 191
Ceruſſa 346
Ceterach 237
Chalcitis 334
Chalybs 347
Chamædrys 220
Chamæpithys 217
Chamæmeli flores 240
Cheiri flores 248
Chermes grana 143
China Chinæ utraque 98. 99
radix 41
Chocolata 185
Chryſocolla 319
Cicer. ſemen 196
Cichorium 47
Cinæ ſemen 151
Cinamomum 93
Cinnabaris factitia 325
Citri cortex 92
Citrulli ſemen 192
Cnicus ſemen 170
Cochlearia 222
Coffee 157
Colchicum 59
Colocynthis 161
Colophonium 292
Colubrinum lignum 113
Conſolida major 79
Saracenica 234
Contrayerva 84
Corallia alba
rubra 342
Coriandrum 145
Cornu cervi 298

Yy

Cortex

Cortex Winteranus 96
Coſtus 20
Cranium humanum 299
Crocus Martis 348
Metallorum 333
Orientalis 241
Cryſtallus 337
Cubebæ 126
Cucumis aſininus 68
Cucumeris ſemen 192
Cucurbitæ ſemen 192
Cuminum 136
Cupreſſi nux 203
Curcuma 27
Cuſcuta 224
Cyani flores 249
Cyclamen 58
Cydoniorum ſemen 204
Cynogloſſa 88
Cyperus longus 20
rotundus 20

D.

Dactyli 186
DaucusCreticus &vulgaris 140
Dens apri 301
Dentali 307
Diagrydium 284
Dictamnus albus 53
Creticus 213
Doronicum 76

E.

Ebulus 73
Ebur 299
Elaterium 69. 281
Elemi gumi 269
Entali 307
Epithymum 224
Equiſetum 233
Eryngium 44
Eryſimum 223
Eſula 69
Eupatorium 212
Euphorbium 281
Euphraſia 221

F.

Fabarum ſemen 197
Ferrum 347
Ficus 186
Filix 55
Fœniculum radix 32
ſemen 133
Fœnum græcum 195
Fraxinella 53
Fraxini cortex 101
Fumaria herba 225
ſemen 150
Fungi 118
Sambuci 119

Galanga

G.

Galanga major 8
minor 8
Galbanum 261
Gallæ 202
Geniſtæ ſemen 155
Genſingh 16
Gentiana 51
Gialappa 64
Glandes 202
Glans unguentaria 170
Glycyrrhiza radix 78
ſuccus 290
Goſſypii ſemen 176
Graminis radix 40
Grana paradiſi 130
Granatorum cortex 103
Granatus lapis 338
Grana tilli 165
Gratiola 228
Guajaci cortex 97
lignum 106
Gummi Arabicum 291
Gutta gamba 283

H.

Hæmatites lapis 349
Halicacabi baccæ 153
Hedera terreſtris 221
Helenium 25
Helleborus albus 71
niger 70
Hermodactyli 59
Hirci ſanguis 303
Hirudines 316
Hordei ſemen 177
Horminum 208
Hyacinthus lapis 338
Hyoſciami ſemen 200
Hypericum herba 222
ſemen 199
Hypociſtidis ſuccus 297
Hyſſopus 212

I.

Jalappa 64
Ichthyocolla 315
Imperatoria 11
Iris florentina 26
Jujubæ 188
Juncus odoratus 214
Juniperi baccæ 137

K.

Kermes grana 143
Kinkina utraque 98

L.

Lacca gummi 267
Lactucæ ſemen 193

 Lada-

Ladanum 278
Lavendulæ flores 239
Laureolæ baccæ 158
Lauri baccæ 138
folia 218
Lazuli lapis 332
Leucojum luteum 248
Leviſticum radix 14
ſemen 134
Lignum Moluccenſe 117
ſanctum 106
vulnerarium 101
Lilium album 79
convallium 246
Lingua cervina 238
Linum ſemen 198
Liquiritia radix 75
ſuccus 290
Liquor ſtipticus 345
Lithargyrium auri
argenti 345
Lucii piſcis mandibulæ 301
Lumbrici terreſtres 312
Lupini ſemen 197
Lupulus 226

M.

Macis 124
Magnes 350
Majorana 205
Malicorium 103
Malva herba 232
ſemen 194
roſea ſeu hortenſis 250
Mandibulæ lucii piſcis 301
Mandragora 89
Manna 286
Margaritæ 304
Marrubium 216
Maſtiche 265
Matricaria 213
Mechoacanna 65
Mel 90
Meliloti flores 247
Meliſſa 207
Melonum ſemen 192
Mentha aquatica 210
criſpa 209
Menthaſtrum 210
Mercurialis 230
Mercurius vivus 330
dulcis 328
præcipitatus ruber 327
ſublimatus 326
viridis 329
Meſpilum 203
Meum 14
Mezereum 158
Milium ſolis 156
Minium 346
Moſchus 307
Mumia 304

Muſcus quernus 233
Myrobalanorum quinque ſpecies 171
Myrrha 279
Myrti baccæ 203

N.

Napi ſemen 149
Naſturtium 223
 ſemen 147
Nepeta 210
Nephriticum lignum 111
Nicotiana 229
Nigella 146
Ninzin 15
Nux cupreſſi 203
 moſchata 124
 vomica 114.164
Nymphæa flores 252
 radix 86

O.

Oculi cancri 302
Olea odorata parandi modus 171
Oleris atri ſemen 149
Olibanum 266
Ononis 49
Opium 257
Opobalſamum 277
Opopanax 288
Orchis 42
Origanum 205
Orobi ſemen 196
Oſmunda 56
Os ſepiæ 302
Oſteocolla 339

P.

Papaver erraticum 251
Papaveris ſemen 201
Parietaria 231
Parthenium 213
Paſſulæ majores
 minores 189
Pavana 117
Pentaphyllum 85
Perſicaria 223
Perſicorum nuclei 183
Petaſites 29
Petroſelinum radix 34
 ſemen 142
Peucedanum 52
Phyllitis 238
Piloſella 236
Pimpinella ſanguiſorba 85.221
 ſaxifraga 39.148
Pini nuclei 182
Piper album 128
 longum 130
 nigrum 128
Piſtacia 182

Pix 272
Plantago 193
Plumbum 345
Pœonia flos 251
  radix 87
  ſemen 201
Polytrichum aureum 237
Polium 212
Polygonum 237
Polypodium 74
Pompholyx 340
Portulacæ ſemen 193
Praſſium 216
Primula veris 246
Pruna damaſcena 188
Pſyllium 160
Pulegium 211
Pulmonaria 234
Pumex 341
Pyretrum 38
Pyrola 236

Q.

Quercus cortex 104
  folia 236
Quinque folium 85

R.

Raphanus ſemen 150
  ruſticanus 35
Rhabarbarum 60
Rhaponticum 63
Rhodium lignum 110
Ribeſium 220
Ricini ſpecies 168
Roſarum flores 248
Roſæ rubræ 252
Roſmarinus flores 238
  herba 208
Rubia tinctorum 50
Rubinus 338
Ruſcus radix 49
  ſemen 156
Ruta herba 215
  ſemen 142
Ruta muraria 237

S.

Sabina 215
Saccharum 289
  ♄ni 347
Sagapenum 263
Sal commune 319
  gemmæ 320
Salſa parilla 41
Salvia herba 209
  flos 240
Sambucus flores 247
  ſemen 176
Sandaracha 265

Sanguis

Sanguis draconis 294
hirci 303
Sanicula 236
Sandalum album
citrinum 109
rubrum 115
Sapphirus 339
Sarcocolla 287
Sardius 339
Saſſafras 105
Satureja 206
Satyrium 42
Scammonium 284
Scariola 193
Schœnanthus 214. 243
Scordium 226
Scilla 37
Scincus 311
Sclarea 208
Scolopendrium 237
Scorzonera 44
Sebeſten 188
Senna 227
Sepiæ os 302
Seſeli ſemen 141
Serici folliculi 316
Serpentaria Virginiana 19
Serpillum 206
Siler ſemen 141
Siliqua dulcis 191
Sinapi 148
Smaragdus 339
Soldanella 228
Sophia chirurgorum 146
Solidago Saracenica 234
Sperma ceti 310
Spicæ flores 239
Spica nardi 22
Staphis agria 159
Stœchatos flores 243
Succinum 268
Sulphur 335
Sumach ſemen 202
Symphitum 79

T.

Tacamahaca 270
Tamarindi 174
Tamariſcus 102
Tanacetum 143. 216
Tartarus albus
ruber 297
Terebinthina 273
Terra Japonica 293
Lemnia &
Sigillata 344
Thea 219
Thlaſpi ſemen 147
Thymus 206
Thus 266
Tiliæ flores 246
Tinctura Corallorum 343
Topaſius 339

Tormen-

Tormentilla 83
Tragacantha 292
Tunicæ flores 243
Turbith 66
minerale 329
Tuſſilago flores 81
herba 232
Tutia 340

V.

Vainiglia 144
Valeriana 18
Verbaſcum flores 250
herba 232
Verbena 235
Veronica 220
Vincetoxicum 23
Violaria 231
flores 244
Virga aurea 234
Viride æris 335
Vitriolum 333
Viſcus quernus 116
Vitrum antimonii 332
Umbilicus marinus 307
Ungula alcis 300
Unicornu foſſile 343
verum 300
Urticæ ſemen 152

W.

Winteranus cortex 96

Z.

Zedoaria 6
Zibethum 308
Zinziber 9
Ziziphi 188

SYLLABUS

SYLLABUS REMEDIORUM, JUXTA PARTES CONSTITUTIVAS.

EX REGNO VEGETABILI.

SALE VOLATILI OLEOSO conſtant. Radices acor vulgaris: Acori indici. Calamus aromaticus. Zedoaria Galanga major & minor. Zinziber. p. 3. ad 10. Cortices aurantiorum. Citri. Cinamomi. Caſſiæ ligneæ & caryophyllatæ. Cortex Winteranus. Guajaci. p. 91 ad 98.

SALE VOLATILI DULCEDINE JUNCTA conſtant. Radices Imperatoriæ ſeu Oſtrutii. Angelicæ. Mei. Leviſtici. Ninzin. p. 11 ad 17.

SALE VOLATILI OLEOSO AROMATICO, MIXTIS PARTICULIS TERREIS ET SALINIS FIXIS. Radices valerianæ. Serpentariæ Virginianæ. Coſti. Cyperi longi & rotundi. Spicæ nardi. Vincetoxici. Helenii. Iridis Florentinæ. Curcumæ. Petaſitidis. Carlinæ. p. 17 ad 31.

SALE AROMATICO BLANDO SEU MITIORE, CUM VIRTUTE DIURETICA APERIENTE ET BLANDE SUDORIFERA. Radices fœniculi. Apii paluſtris. Petroſelini. p. 32 ad 34.

SALE VOLATILI AROMATICO SPIRITUOSO PENETRANTE ET PAUCO ACIDO VOLATILI. Radices armoraciæ ſeu raphani ruſticani. Ari. quæ conſtat ſale volatili acerrimo abſque oleo. Scillæ. Pyrethri. conſtat ſale acri fixo abſque oleo. p. 34 ad 38.

SALE ACRI ABSQUE OLEO TAM FIXO QUAM VOLATILI. Radices pimpinellæ ſaxifragæ. Graminis. p. 38 ad 40.

SALE BLANDO BALSAMICO. Radices Salſæparillæ. Chinæ. p. 41. 42.

SALE VOLATILI BLANDO CUM TENUI VISCOSITATE TEMPERATA. Orchis ſeu Satyrium. p. 42. 43. Boletus cervi. p. 118.

SALE BLANDO FIXO, TENUI VISCOSITATI INCLUSO. Radices Scorzoneræ. Eryngii. Bardanæ. Cichorii. p. 44. ad 47.

SALE BLANDO DIURETICO. Radices Aſparagi. Ruſci. Ononidis. Rubiæ tinctorum. p. 48 ad 50. Lignum Nephriticum. p. 111.

SALE AMARO FIXO. Radices Gentianæ. p. 50. Sale fixo amariſſimo cortex Chinæ chinæ. p. 99.

SALE FIXO CUM PAUCO VOLATILI RESINÆ INCLUSO. Radices Peucedani. Fraxinellæ. Ariſtolochiæ longæ, rotundæ, fabaceæ. Filicis. Oſmundæ. p. 52 ad 56.

Radices purgantes SALE VOLATILI GAUDENTES. Aſarum. Cyclamen. Hermodactyli. Rhabarbarum. Rhaponticum. p. 56 ad 64.

Radices purgantens SALE FIXO RESINÆ INCLUSO conſtantes. Gialappa. Mechoacanna. Turbith. p. 64 ad 67. Agaricus. p. 120.

Radices SALE FIXO ACRI CONSTANTES & vehementiſſime purgantes. Cucumis aſininus. Eſula major & minor. Helleborus niger & albus. Bryonia. p. 68 ad 73.

Radices SALE PURGANTE BLANDO conſtantes & vulgo laxantes. Ebulus. Polypodium. p. 73. 74.

PARTICULIS BLANDE VISCOSIS CUM DULCEDINE conſtant. Radices Glycyrrhizæ. Doronici. p. 75. 77.

AQUO-

AQUOSIS CRASSIS SEU MUCILAGINOSIS conſtant. Radices Althææ. Liliorum alborum. Symphiti ſeu conſolidæ majoris. p. 77 ad 80.

MUCAGINE TENUIORE PAUCIS FIXIS SALINIS. Radices Tuſſilaginis. p. 81.

PARTICULIS TERREIS conſtant. Radices Alcannæ. Biſtortæ. Tormentillæ. Contrayervæ. Caryophyllatæ. Pimpinellæ ſanguiſorbæ. Pentaphylli. Nymphæa. p. 80 ad 87. Cortices Caſtaneorum. Granatorum. Quercus. p. 103. ſeq. Santalum rubrum. Viſcus quernus. p. 115. 116. Fungus orbicularis ſeu boviſta. p. 121.

TERREIS ET SIMUL MUCAGINOSIS VOLATIBILIBUS conſtant. Radices pæoniæ. Cynogloſſæ. Mandragoræ. p. 87. ad 90.

SALE FIXO DIURERICO. Cortex fraxini. Tamariſci. Capparis. 101. ſeq.

SALE OLEOSO AROMATICO. Lignum ſaſſafras. Guajacum & ſanctum. Aloës. Agallochum. Santalum citrinum & album. Rhodium. p. 105 ad 111.

SALIBUs FIXIS AMARISSIMIS. Lignum colubrinum. p. 113.

SALE PURGANTE TAM FIXO QUAM VOLATILI. Pavana ſeu lignum Moluccenſe. p. 117.

TERREIS SUBAUSTERIS conſtat. Auricula judæ ſeu fungus ſambuci. p. 119.

SALE VOLATILI OLEOSO AROMATICO. Cariophylli. Anthophylli. Amomum. Macis. Nux Moſchata. Cubebæ. Cardamomum. Piper album, nigrum, longum. Grana Paradiſi. Semen Ammios. Aniſi. & Aniſi Chinenſis. Fœniculi. Leviſtici. p. 122 ad 134.

SALE VOLATILI AROMATICO CUM BLANDA VISCOSITATE. Semen Anethi. Cumini. Carvi. Baccæ Juniperi. Lauri. Carpobalſamum. Semen Dauci. Selelios.

Sileris. Petroſelini. Apii. Rutæ. Tanaceti. p. 125. ad 134.
SALE AROMATICO BLANDO CONFORTANTE. Grana Kermes. Vainiglia. Semen Coriandri. Nigellæ. p. 143 ad 146.
SALIBUS ACRIBUS FIXIS. Semen Sophiæ Chirurgorum. Thlaſpi. Naſturtii. Saxifragiæ. p. 146 ad 148.
SALE VOLATILI ACRI CUM OLEO CRASSO. Semen Sinapi. Napi. Oleris atri. Raphani. p. 148 ad 150.
SALE FIXO AMARO. Semen Fumariæ. Cinæ. Cardui Benedicti. Cardui Mariæ. Bardanæ. Urticæ. p. 150 ad 153.
SALE BLANDO DIURETICO. Semen Alkekengi ſeu Halicacabi. Geniſtæ. Milii Solis. Ruſci. Aſparagi. Coffee. p. 153 ad 157.
SALE PURGANTE ACERRIMO. Semen Mezerei ſeu Laureolæ. Staphiſagriæ. Pſyllium, habet ſalem blandum purgantem. p. 158 ad 161.
SALE ACRI PURGANTE TAM VOLATILI QUAM EIXO. Colocynthis ſeu alhandal. Nux vomica. p. 161 ad 165.
SALE PURGANTE TAM FIXO QUAM VOLATILI OLEO CRASSO INCLUSO. Grana tilli. Ricinus Medius. Ricinus Americanus. Cataputia minor. Semen Cnici ſeu Carthami. Balanus mirepſica ſeu nux been. Myrobalanorum quinque ſpecies. Caſſiæ fructus. Tamarindi. Semen Atriplicis. Semen Sambuci. à p. 165 ad 176.
BLANDA VISCOSITATE conſtant. Semen Goſſypii ſeu bombacis. Hordeum. Semen Cannabis. p. 176 ad 178.
SUBDULCIBUS OLEOSIS ET TEMPERATIS conſtant. Amygdalæ dulces & amaræ. Piſtacia. Nuclei Pini, Ceraſorum, Perſicorum. Anacardium. Cacao. p. 179 ad 186.
DULCIBUS VISCOSIS conſtant. Dactyli. Ficus ſeu Caricæ. Jujubæ ſeu Ziziphi. Sebeſten. Pruna Damaſcena. Paſſulæ

majores

majores & minores ſeu Corinthiacæ. Siliqua dulcis ſeu Ceratonia. p. 186 ad 191.

PARTICULIS AQUOSIS ET CRASSIS CUM OLEO PAUCO. Semina quatuor frigida majora & minora. p. 192. & 193.

PARTICULIS CRASSIS AQUOSIS SEU MUCAGINOSIS ABSQUE OLEOSE TAM SUBITO MANIFESTANTE conſtant. Semen Altheæ; Malvæ. Aquilegiæ. Bardanæ. Fœnum Græcum. Semen Orobi; Ciceris; Lupini; Fabarum; Lini; Hyperici; [eſt reſinoſum.] p. 194 ad 199.

MUCAGINE VOLATILI prædita ſunt ſemen Hyoſciami, Pœoniæ, Papaveris. p. 200. 201.

TERREIS conſtant. Gallæ. Glandes. Semen Sumach. Meſpilum. Nux Cupreſſi. Baccæ Myrti. Semen Cydoniorum. (terreas in mucaginoſis obtinet.) p. 202 ad 204

PLANTÆ SALE VOLATILI OLEOSO SEU AROMATICO conſtant. Herbæ Origanum. Majorana. Serpillum. Thymus. Satureja. Meliſſa. Horminum ſclarea dictum. p. 205. ad 208.

PLANTÆ SALE AROMATICO VOLATILI CUM QUIBUSDAM FIXIS CONSTANTES. Roſmarinus. Salvia. Mentha Criſpa & Aquatica. Menthaſtrum. Nepetha. Calamintha. Pulegium. Eupatorium. Hyſſopus. Polium. Dictamnus Creticus. Matricaria ſeu Parthenium. Artemiſia. p. 205 ad 214

PLANTÆ SALE AROMATICO VOLATILI OLEOSO CUM MULTO FIXO CONSTANTES. Schœnanthus ſeu Juncus odoratus. Abrotanum mas. Ruta. Sabina. Marrubium ſeu Praſſium. Tanacetum. Abſynthium vulgare & Ponticum. p. 214 ad 217.

PLANTÆ SALE AROMATICO CUM PAUCA RESINA conſtantes. Chamæpithys. Botrys. Lauri folia. p. 217. 218.

PLANTÆ SALE BLANDO CUM PAUCO VOLATILI CONSTANTES. Agrimonia. Thea. Betonica. Chamædrys. Ribeſium. p. 218. 220.

PLANTÆ SALE FIXO BLANDO CUM QUIBUSDAM TERREIS CONSTANTES. Veronica. Hedera Terreſtris. Pimpinella. Euphraſia. Hypericum. p. 220 ad 222.

PLANTÆ SALE AROMATICO OLEOSO PENETRANTISSIMO CUM ACIDO VOLATILI CONSTANTES. Cochlearia. Naſturtium. p. 222.

PLANTÆ SALIBUS FIXIS SEU AMARIS CONSTANTES CUM PAUCO VOLATILI SALE. Centaurium minus. Carduus Benedictus. Fumaria. Scordium. Lupulus. p. 224. ad 227.

PLANTÆ PURGANTES. Senna. Soldanella. Gratiola. Nicotiana. p. 227 ad 230.

PLANTÆ CONSTANTES SALE NITROSO LAXANTE. Mercurialis. Parietaria. Violaria. p. 230. 231.

PLANTÆ CONSTANTES PARTICULIS OLEOSIS AQUOSIS CRASSIS. Althæa. Malva. Tuſſilago. Verbaſcum. p. 231. 232.

Plantæ TERREIS CONSTANTES. Equiſetum ſeu Cauda Equina. Muſcus quernus. Pulmonaria. Solidago. Saracenica. ſeu Virga aurea. Burſa Paſtoris. Verbena. Folia quercus. Pyrola. Sanicula. Alchymilla. Piloſella. Polygonum. p. 233 ad 237.

PLANTÆ PARTICULIS TERREIS TEMPERATIS ET BLANDIS VOLATILIBUS CONSTANTES. Herbæ capillares. Phyllitis ſeu Lingua Cervina. p. 237.

FLORES SALIBUS VOLATILIBUS OLEOSIS AROMATICIS CONSTANTES. Flores Anthos. Lavendulæ. Salviæ. Chamæmeli. Crocus Orientalis. Flores Stœchados. Schœnanthos. Tunicæ. Violarum. Roſarum pallidarum. Tiliæ. Lilium convallium. Primula veris. p. 238 ad 246.

Flores

Flores MUCAGINE TENUI pollentes. Meliloti. Sambuci. p. 247.

Flores SALE FIXO CUM PAUCIS VOLATILIBUS CONSTANTES. Cheiri. Calendulæ. p. 248.

Flores AQUEIS CONSTANTES. Borrago. Bugloſſa. Cyanus. (aliis terreis, ſed toti in mucilaginem abeunt.) p. 249.

Flores MUCAGINE CRASSA CONSTANTES. Malvæ roſeæ ſeu hortenſis. Verbaſcum. Pæonia. Papaver erraticum. p. 250. 251.

Flores TERREIS CONSTANTES. Balauſtiorum. Nympheæ. Roſarum rubrarum. p. 252.

SALINIS VOLATILIBUS OLEOSIS CUM MUCILAGINOSIS TENUISSIMIS SUBTILIBUS conſtant. Camphora. Opium. p. 254 ad 259.

Particulis SALINIS VOLATILIBUS OLEOSIS CUM VIRTUTE LAXANDI conſtant. Ammoniacum. Galbanum. Bdellium. Aſa fœtida. Sagapenum. p. 259 ad 264.

Reſinæ SPISSIORES ET SOLIDIORES ARIDIORES SALE OLEOSO CONSTANTES. Gummi anime & Cancamum. Sandaracha. Maſtiche. Thus ſeu Olibanum. Lacca. Succinum. Elemi. Caranna & Tacamahaca. Benzoe ſeu Aſa Dulcis. Pix. Styrax Calamita. p. 264. ad 273.

Reſinæ LIQUIDÆ SALE VOLATILI OLEOSO CONSTANTES. Therebintina. Balſamum Peruvianum & Tolutanum nigrum. Balſamum Peruvianum album. Balſamum de Copaiva. Ambra liquida. Opobalſamum. Ladanum. Styrax liquida. p. 273 ad 278.

Reſina SALE TAM VOLATILI QUAM FIXO CUM INTENSISSIMA AMARITIE CONSTANS. Myrrha. p. 279.

Reſinæ GUMMOSÆ ET SUCCI PURGANTES FORTIORES ET MITIORES. Euphorbium. Elaterium. Gutta Gamba. Aloe. Manna. Sarcocolla. Opopanax. p. 281 ad 288.

Succi CONCRETI MUCAGINOSI DULCES. Saccharum. Succus Glycyrrhicæ. Mel. p. 289 ad 291.
GUMMOSA AQUOSA CRASSA. Gummi Arabicum. p. 291. Tragacantha. p. 291. ſeq.
Reſinæ OLEO CRASSO conſtantes. Bitumen Judaicum. Colophonium. Cera. p. 292. ſeq.
Reſinæ & ſucci TERREIS conſtantes. Sanguis Draconis. Terra Japonica ſeu Catechu. Succus acaciæ. Hypociſtidis ſuccus. p. 294 ad 297.
Succus SALE SALSO conſtans, nonnihil laxante. Tartarus albus & ruber. p. 297. ſeq.

EX REGNO ANIMALI.

PArtes Animales SALE VOLATILI OLEOSO CUM MULTIS TERREIS CONSTANTES. Cornu Cervi. Sanguis hirci. Mumia. Margaritæ. Bezoar orientalis & occidentalis. Dentali. Entali. Umbilicus marinus. p. 298 ad 307.
Partes animales SALE VOLATILI OLEOSO CUM ODORE FRAGRANTISSIMO conſtantes. Moſchus. Zibethum. Caſtoreum. p. 307 ad 310.
Pars animalis SALE VOLATILI CUM MULTA PINGUEDINE conſtans. Sperma. Ceti. p. 310.
Animalia SALE VOLATILI BLANDO conſtantia. Scincus. Lumbrici terreſtres. p. 311. ſeq.
Animal CAUSTICUM ET ACERRIMUM. Cantharides. p. 313.
Partes animales OLEOSIS CRASSIS conſtantes. Ichthyocolla. p. 315.
Animal SANGUISUGUM. Hirudines. p. 316.
Pars animalis TERREIS CUM PAUCO SALE VOLATILI conſtans. Folliculi ſerici. p. 316.

EX

EX REGNO MINERALI.

Mineralia SALE VOLATILI SALSO conſtantia. Sal Ammoniacum. Borax. Chryſocolla Cryſtallina. p. 317 ad 319.

Mineralia SALE FIXO SALSO conſtantia. Sal commune. Sal gemmæ. Nitrum. Alumen rupeum & romanum. Alumen plumoſum. p. 319 ad 323.

Mineralia SALE SALSO VOLATILI ET CORROSIVO conſtantia. Calx viva. Arſenicum album & rubrum. Auripigmentum. Cinabaris factitia & nativa. [nullam habet corroſivitatem] Mercurius ſublimatus. Mercurius præcipitatus ruber. Mercurius dulcis. Mercurius viridis. Turbith Minerale. p. 323 ad 329.

Minerale SALE VOLATILI ALCALICIS AFFINI ABSQUE TERRA conſtans. Mercurius vivus. p. 330.

Minerale SALE VOLATILI SULPHUREO CUM TERRA conſtans. Antimonium crudum. p. 331.

Mineralia SALE VOLATILI PURGANTE CUM TERRA ADSTRINGENTE conſtantia. Lapis Lazuli & Armenus. Vitrum Antimoniii. Crocus Metallorum, p. 332. ſeq.

Mineralia SALE ACIDO CUM PAUCO ALCALI conſtantia. Vitriolum album. Hungaricum & Romanum. Chalcitis. Viride æris. p. 332 ad 335.

Minerale OLEOSIS SEU PINGUIBUS CUM ACIDO VOLATILI conſtans. Sulphur. p. 335. ſeq.

Mineralia FIRMIORIBUS PARTICULIS conſtantia. Aurum foliatum. Argentum foliatum. p. 336.

Mineralia TERREIS conſtantia. Lapis calaminaris. Cryſtallus. Fragmenta lapidum pretioſorum. Granatus. Hyacinthus. Rubinus. Sapphirus. Sardius. Smaragdus. Oſteocolla

colla. Tutia. Pompholyx. Nihil album Pumex. Spongiæ lapis. p. 337 ad 342.

Mineralia PARTICULIS TERREIS CUM ALCALI conſtantia. Corallia rubra & alba. Unicornu foſſile. Bolus armena vulgaris & alba. Terra Lemnia & Sigillata. p. 342 ad 345.

Metallum SOLIDUM FLEXILE tamen. PLUMBUM. Ex eo præparata ſunt. Lithargyrium auri & argenti. Ceruſſa. Minium. Saccharum Saturni. p. 345 ad 347.

Metallum SOLIDUM TERRA conſtans. Ferrum. Chalybs. p. 349.

Subſtantia LAPIDEA AD METALLICAM FERRI accedens Magnes. p. 350.

Spec — 25— 26 — 35— 60 — 198 — 216 —
51— 53 — 215 —